

10-27-00

A

<b>UTILITY PATENT APPLICATION TRANSMITTAL</b>  <i>(Only for new nonprovisional applications under 37 CFR 1.53(b))</i>	Attorney Docket No.	BB1043 US NA DIV
	First Named Inventor or Application Identifier	
	Jonathan Edward Lightner et al.	
	Express Mail Label No.	EL073740691US
	Express Mailing Date	October 26, 2000

JC806 U.S. PTO  
09/697379

10/26/00

**APPLICATION ELEMENTS**

See MPEP chapter 600 concerning utility patent application contents.

ADDRESS TO: Assistant Commissioner for Patents  
Box Patent Application  
Washington, DC 20231

1. ☒ Fee (Authority to charge deposit account below.)  
(Submit an original, and a duplicate for fee processing)
2. ☒ Specification [Total Pages **161**]  
(preferred arrangement set forth below)
- Descriptive title of the invention
  - Cross References to Related Applications (if needed)
  - Statement Regarding Fed sponsored R & D (if needed)
  - Reference to Microfiche Appendix (if filed)
  - Background of the Invention
  - Brief Summary of the Invention
  - Brief Description of the Drawings (if filed)
  - Detailed Description
  - Claim(s)
  - Abstract of the Disclosure
3. ☐ Drawing(s) (35 USC 113) [Total Sheets **0**]
4. ☒ Oath or Declaration [Total Pages **1**]
- a. ☐ Newly executed (original or copy)
  - b. ☒ Copy from a prior application (37 CFR 1.63(d))  
(for continuation/divisional with Box 14 completed)
  - i. ☐ **DELETION OF INVENTORS**  
Signed Statement below at 15 deleting inventor(s) named in the prior application, see 37 CFR 1.63(d)(2) and 1.33(b).
5. ☒ Incorporation by Reference (useable if Box 4b is checked)  
The entire disclosure of the prior application, from which a copy of the oath or declaration is supplied under Box 4b, is considered as being part of the disclosure of the accompanying application and is hereby incorporated by reference therein.

6. ☐ Microfiche Computer Program (Appendix)
7. Nucleotide and/or Amino Acid Sequence Submission (if applicable, all necessary)
- a. ☒ Computer Readable Copy
  - b. ☒ Paper Copy (identical to computer copy)
  - c. ☒ Statement verifying identity of above copies

**ACCOMPANYING APPLICATION PARTS**

8. ☐ Power of Attorney
9. ☐ Information Disclosure Statement (IDS)/Cover Letter plus PTO-1449 ☐ Copies of IDS Citations
10. ☒ Preliminary Amendment
11. ☒ Return Receipt Postcard (MPEP 503)  
(Should be specifically itemized)
12. ☐ Certified Copy of Priority Document(s)  
(if foreign priority is claimed)
13. ☐ Other:

**14. If a CONTINUING APPLICATION, check appropriate box and supply the requisite information:**
☐ Continuation    ☒ Divisional    ☐ Continuation-in-part (CIP)    of prior Application No.: 09/133,962

15. ☐ **DELETION OF INVENTOR(S) STATEMENT:** This application is being filed by less than all the inventors named in the prior application. In accordance with 37 CFR 1.63(d)(2) and 1.33(b), the Assistant Commissioner is requested to delete the name(s) of the following person or persons who are not inventors of the invention being claimed in this application:
16. ☒ Amend the specification by inserting before the first line the sentence:  
-- This is a division of Application No. 09/133,962 filed August 14, 1998, now pending. --
17. ☐ Cancel in this application original claims \_\_\_\_ of the prior application before calculating the filing. (At least one original independent claim must be retained for filing purposes.)
18. ☐ Priority of foreign Application No. \_\_\_\_\_ filed on \_\_\_\_\_ in \_\_\_\_\_  
\_\_\_\_\_ is claimed under 35 U.S.C. 119.  
(country)

CLAIMS	(1) FOR	(2) NUMBER FILED	(3) NUMBER EXTRA	(4) RATE	(5) CALCULATIONS
	<b>TOTAL CLAIMS</b> (37 CFR 1.16(e))	67 - 20 =	47	x \$ 18 =	\$ 846.00
	<b>INDEPENDENT CLAIMS</b> (37 CFR 1.16(b))	18 - 3 =	15	x \$ 80 =	\$ 1200.00
	<b>MULTIPLE DEPENDENT CLAIM(S) (if applicable)</b>			+ \$ 270 =	270.00
				<b>BASIC FEE</b> (37 CFR 1.16(a))	+ \$ 710.00
				<b>TOTAL =</b>	\$ 3026.00

19. The Commissioner is hereby authorized to credit overpayments or charge the following fees to Deposit Account No. 04-1928:

a. ☒ Fees required under 37 CFR 1.16.

b. ☒ Fees required under 37 CFR 1.17.

20. ☐ Other:

#### 21. CORRESPONDENCE ADDRESS

NAME	Lynne M. Christenbury				
ADDRESS	E. I. du Pont de Nemours and Company				
	Legal - Patents				
	1007 Market Street				
CITY	Wilmington	STATE	Delaware	ZIP CODE	19898
COUNTRY	U.S.A.	TELEPHONE	302-992-5481	FAX	302-892-1026

#### 22. SIGNATURE OF ATTORNEY OR AGENT REQUIRED

NAME	Lynne M. Christenbury	REG. NO.:	30,971
SIGNATURE	<i>Lynne M. Christenbury</i>		
DATE	26 October 2000		

EXPRESS MAIL LABEL NO. EL073740691US

PATENT  
IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

IN THE APPLICATION OF:

JONATHAN EDWARD LIGHTNER  
JOHN JOSEPH OKULEY

CASE NO.: BB-1043 US NA DIV

APPLN. NO.: UNKNOWN

GROUP ART UNIT: UNKNOWN

FILED: CONCURRENTLY HERewith

EXAMINER: UNKNOWN

FOR: GENES FOR MICROSOMAL DELTA-12  
FATTY ACID DESATURASES AND  
RELATED ENZYMES FROM PLANTS

Date: OCTOBER 26, 2000

Assistant Commissioner for Patents  
Washington, DC 20231

Sir:

Preliminary Amendment

This is submitted to facilitate prosecution of the above-identified application.

In the Claims

Kindly cancel claims 2-8, 11, 12, 14 and 21-23.

Kindly amend the following claims:

9. (amended) Oil obtained from the seeds of [the plants of Claim 8] a soybean plant having a chimeric gene comprising a nucleic acid fragment encoding a fatty acid desaturase with an amino acid identity of 50% or greater to the polypeptide encoded by any one of the sequences set forth in SEQ ID NOS: 1, 3, 5, 7, 9, 11 or 15.

10. (amended) A method of producing seed oil containing altered levels of [unsaturated] hydroxylated fatty acids comprising:

- (a) transforming a plant cell of an oil-producing species with a chimeric gene [of Claim 5] comprising a nucleic acid fragment encoding a delta-12 hydroxylase;
- (b) growing fertile plants from the transformed plant cells of step (a);

(c) screening progeny seeds from the fertile plants of step (b) for the desired levels of [unsaturated] hydroxylated fatty acids; and

(d) processing the progeny seed of step (c) to obtain seed oil containing altered levels of [unsaturated] hydroxylated fatty acids.

15. (amended) A method for altering fatty acids composition in seeds comprising:

(a) making a cross between a mutant line with altered fatty acid composition with a plant [containing] comprising [the] a chimeric gene [of Claim 7] capable of causing altered levels of fatty acids in a transformed plant cell, said chimeric gene comprising a nucleic acid fragment selected from the group consisting of:

(i) an isolated nucleic acid fragment comprising a nucleic acid sequence encoding a plant delta-12 desaturase or a delta-12 hydroxylase wherein said isolated nucleic acid fragment hybridizes to one of the nucleotide sequences set forth in SEQ ID NOS:1, 3, 5, 7, 9, 11, and 15 under one of the following sets of conditions:

(a) hybridization in 50 mM Tris, pH 7.6, 6X SSC, 5X Denhardt's, 0.5% sodium dodecyl sulfate (SDS), 100 µg denatured calf thymus DNA and 50 °C and wash twice with 2X SSC, 0.5% SDS at room temperature for 15 min each, then wash twice with 0.2X SSC, 0.5% SDS at room temperature for 15 min each and then wash twice with 0.2X SSC, 0.5% SDS at 50 °C for 15 min each;

(b) hybridization in 6X SSPE, 5X Denhardt's solution, 0.5% sodium dodecyl sulfate (SDS), 5% dextran sulfate, 100 µg denatured calf thymus DNA at 50 °C and wash twice with 2X SSC, 0.5% SDS at room temperature for 15 min each, then wash twice with 0.2X SSC, 0.5% SDS at room temperature for 15 min each and then wash twice with 0.2X SSC, 0.5% SDS at 50 °C for 15 min each; or

(c) hybridization in 50% formamide, 5X SSPE, 1% sodium dodecyl sulfate (SDS), 1% Denhardt's Reagent, 100 µg denatured salmon sperm DNA at 42 °C and wash twice with 2X SSPE, 0.2% SDS at 42° C for 15 min each, then wash twice with 0.2X SSPE, 0.2% SDS at 55 °C for 30 min each;

(ii) the isolated nucleic acid fragment of (i) wherein the encoded polypeptide has an amino acid identity of 60% or greater to the polypeptide encoded by any one of the sequences set forth in SEQ ID NOS:1, 3, 5, 7, 9, 11, or 15; or

(iii) the isolated nucleic acid fragment of (i) wherein the nucleic acid identity is 90% or greater to any one of the sequences set forth in SEQ ID NOS:1, 3, 5, 7, 9, 11, or 15;

(b) growing fertile plants from seeds obtained from the cross; and

(c) screening progeny seeds from the fertile plants of step (b) for seeds containing altered fatty acids.



16. (amended) A method for reducing polyunsaturated fatty acids in rapeseed oil comprising:

(a) making a cross between a rapeseed variety with increased oleic acid content or reduced linolenic acid content with a plant [containing] comprising [the] a chimeric gene [of Claim 7] capable of causing altered levels of fatty acids in a transformed plant cell, said chimeric gene comprising a nucleic acid fragment selected from the group consisting of:

(i) an isolated nucleic acid fragment comprising a nucleic acid sequence encoding a plant a delta-12 desaturase or a delta-12 hydroxylase wherein said isolated nucleic acid fragment hybridizes to one of the nucleotide sequences set forth in SEQ ID NOS:1, 3, 5, 7, 9, 11, and 15 under one of the following sets of conditions:

(a) hybridization in 50 mM Tris, pH 7.6, 6X SSC, 5X Denhardt's, 0.5% sodium dodecyl sulfate (SDS), 100 µg denatured calf thymus DNA and 50 °C and wash twice with 2X SSC, 0.5% SDS at room temperature for 15 min each, then wash twice with 0.2X SSC, 0.5% SDS at room temperature for 15 min each and then wash twice with 0.2X SSC, 0.5% SDS at 50 °C for 15 min each;

(b) hybridization in 6X SSPE, 5X Denhardt's solution, 0.5% sodium dodecyl sulfate (SDS), 5% dextran sulfate, 100 µg denatured calf thymus DNA at 50 °C and wash twice with 2X SSC, 0.5% SDS at room temperature for 15 min each, then wash twice with 0.2X SSC, 0.5% SDS at room temperature for 15 min each and then wash twice with 0.2X SSC, 0.5% SDS at 50 °C for 15 min each; or

(c) hybridization in 50% formamide, 5X SSPE, 1% sodium dodecyl sulfate (SDS), 1% Denhardt's Reagent, 100 µg denatured salmon sperm DNA at 42 °C and wash twice with 2X SSPE, 0.2% SDS at 42° C for 15 min each, then wash twice with 0.2X SSPE, 0.2% SDS at 55 °C for 30 min each;

(ii) the isolated nucleic acid fragment of (i) wherein the encoded polypeptide has an amino acid identity of 60% or greater to the polypeptide encoded by any one of the sequences set forth in SEQ ID NOS:1, 3, 5, 7, 9, 11, or 15; or

(iii) the isolated nucleic acid fragment of (i) wherein the nucleic acid identity is 90% or greater to any one of the sequences set forth in SEQ ID NOS:1, 3, 5, 7, 9, 11, or 15;

(b) growing fertile plants from seeds obtained from the cross; and

(c) screening progeny seeds from the fertile plants of step (b) for seeds containing reduced polyunsaturated fatty acids.

18. (amended) A method for reducing saturated fatty acids in rapeseed seeds comprising:

09679-10268

(a) making a cross between a rapeseed variety with increased oleic acid content with a plant [containing] comprising [the] a chimeric gene [of Claim 7] capable of causing altered levels of fatty acids in a transformed plant cell, said chimeric gene comprising a nucleic acid fragment selected from the group consisting of:

(i) an isolated nucleic acid fragment comprising a nucleic acid sequence encoding a encoding a delta-12 desaturase or a delta-12 hydroxylase wherein said nucleic acid fragment hybridizes to one of the nucleotide sequences set forth in SEQ ID NOS:1, 3, 5, 7, 9, 11, and 15 under one of the following sets of conditions:

(a) hybridization in 50 mM Tris, pH 7.6, 6X SSC, 5X Denhardt's, 0.5% sodium dodecyl sulfate (SDS), 100 µg denatured calf thymus DNA and 50 °C and wash twice with 2X SSC, 0.5% SDS at room temperature for 15 min each, then wash twice with 0.2X SSC, 0.5% SDS at room temperature for 15 min each and then wash twice with 0.2X SSC, 0.5% SDS at 50 °C for 15 min each;

(b) hybridization in 6X SSPE, 5X Denhardt's solution, 0.5% sodium dodecyl sulfate (SDS), 5% dextran sulfate, 100 µg denatured calf thymus DNA at 50 °C and wash twice with 2X SSC, 0.5% SDS at room temperature for 15 min each, then wash twice with 0.2X SSC, 0.5% SDS at room temperature for 15 min each and then wash twice with 0.2X SSC, 0.5% SDS at 50 °C for 15 min each; or

(c) hybridization in 50% formamide, 5X SSPE, 1% sodium dodecyl sulfate (SDS), 1% Denhardt's Reagent, 100 µg denatured salmon sperm DNA at 42 °C and wash twice with 2X SSPE, 0.2% SDS at 42° C for 15 min each, then wash twice with 0.2X SSPE, 0.2% SDS at 55 °C for 30 min each;

(ii) the isolated nucleic acid fragment of (i) wherein the encoded polypeptide has an amino acid identity of 60% or greater to the polypeptide encoded by any one of the sequences set forth in SEQ ID NOS:1, 3, 5, 7, 9, 11, or 15; or

(iii) the isolated nucleic acid fragment of (i) wherein the nucleic acid identity is 90% or greater to any one of the sequences set forth in SEQ ID NOS:1, 3, 5, 7, 9, 11, or 15;

(b) growing fertile plants from seeds obtained from the cross; and

(c) screening progeny seeds from the fertile plants of step (b) for seeds containing reduced saturated fatty acids.

19. (amended) A method for reducing polyunsaturated fatty acids in soybean oil comprising:

(a) making a cross between a soybean variety with increased oleic acid content or reduced linolenic acid content with a plant [containing] comprising [the] a chimeric gene [of Claim 7] capable of causing altered levels of fatty acids in a

transformed plant cell, said chimeric gene comprising a nucleic acid fragment selected from the group consisting of:

(i) an isolated nucleic acid fragment comprising a nucleic acid sequence encoding a plant a delta-12 desaturase or a delta-12 hydroxylase wherein said isolated nucleic acid fragment hybridizes to one of the nucleotide sequences set forth in SEQ ID NOS:1, 3, 5, 7, 9, 11, and 15 under one of the following sets of conditions:

(a) hybridization in 50 mM Tris, pH 7.6, 6X SSC, 5X Denhardt's, 0.5% sodium dodecyl sulfate (SDS), 100 µg denatured calf thymus DNA and 50 °C and wash twice with 2X SSC, 0.5% SDS at room temperature for 15 min each, then wash twice with 0.2X SSC, 0.5% SDS at room temperature for 15 min each and then wash twice with 0.2X SSC, 0.5% SDS at 50 °C for 15 min each;

(b) hybridization in 6X SSPE, 5X Denhardt's solution, 0.5% sodium dodecyl sulfate (SDS), 5% dextran sulfate, 100 µg denatured calf thymus DNA at 50 °C and wash twice with 2X SSC, 0.5% SDS at room temperature for 15 min each, then wash twice with 0.2X SSC, 0.5% SDS at room temperature for 15 min each and then wash twice with 0.2X SSC, 0.5% SDS at 50 °C for 15 min each; or

(c) hybridization in 50% formamide, 5X SSPE, 1% sodium dodecyl sulfate (SDS), 1% Denhardt's Reagent, 100 µg denatured salmon sperm DNA at 42 °C and wash twice with 2X SSPE, 0.2% SDS at 42° C for 15 min each, then wash twice with 0.2X SSPE, 0.2% SDS at 55 °C for 30 min each;

(ii) the isolated nucleic acid fragment of (i) wherein the encoded polypeptide has an amino acid identity of 60% or greater to the polypeptide encoded by any one of the sequences set forth in SEQ ID NOS:1, 3, 5, 7, 9, 11, or 15; or

(iii) the isolated nucleic acid fragment of (i) wherein the nucleic acid identity is 90% or greater to any one of the sequences set forth in SEQ ID NOS:1, 3, 5, 7, 9, 11, or 15;

(b) growing fertile plants from seeds obtained from the cross; and

(c) screening progeny seeds from the fertile plants of step (b) for seeds containing reduced polyunsaturated fatty acids.

20. (amended) A method for reducing saturated fatty acids in soybean seeds comprising:

(a) making a cross between a soybean variety with increased oleic acid content with a plant [containing] comprising [the] a chimeric gene [of Claim 7] capable of causing altered levels of fatty acids in a transformed plant cell, said chimeric gene comprising a nucleic acid fragment selected from the group consisting of:

096974.1.02608

(i) an isolated nucleic acid fragment comprising a nucleic acid sequence encoding a plant enzyme encoding a delta-12 desaturase or a delta-12 hydroxylase wherein said nucleic acid fragment hybridizes to one of the nucleotide sequences set forth in SEQ ID NOS:1, 3, 5, 7, 9, 11, and 15 under one of the following sets of conditions:

(a) hybridization in 50 mM Tris, pH 7.6, 6X SSC, 5X Denhardt's, 0.5% sodium dodecyl sulfate (SDS), 100 µg denatured calf thymus DNA and 50 °C and wash twice with 2X SSC, 0.5% SDS at room temperature for 15 min each, then wash twice with 0.2X SSC, 0.5% SDS at room temperature for 15 min each and then wash twice with 0.2X SSC, 0.5% SDS at 50 °C for 15 min each;

(b) hybridization in 6X SSPE, 5X Denhardt's solution, 0.5% sodium dodecyl sulfate (SDS), 5% dextran sulfate, 100 µg denatured calf thymus DNA at 50 °C and wash twice with 2X SSC, 0.5% SDS at room temperature for 15 min each, then wash twice with 0.2X SSC, 0.5% SDS at room temperature for 15 min each and then wash twice with 0.2X SSC, 0.5% SDS at 50 °C for 15 min each; or

(c) hybridization in 50% formamide, 5X SSPE, 1% sodium dodecyl sulfate (SDS), 1% Denhardt's Reagent, 100 µg denatured salmon sperm DNA at 42 °C and wash twice with 2X SSPE, 0.2% SDS at 42° C for 15 min each, then wash twice with 0.2X SSPE, 0.2% SDS at 55 °C for 30 min each;

(ii) the isolated nucleic acid fragment of (i) wherein the encoded polypeptide has an amino acid identity of 60% or greater to the polypeptide encoded by any one of the sequences set forth in SEQ ID NOS:1, 3, 5, 7, 9, 11, or 15; or

(iii) the isolated nucleic acid fragment of (i) wherein the nucleic acid identity is 90% or greater to any one of the sequences set forth in SEQ ID NOS:1, 3, 5, 7, 9, 11, or 15;

(b) growing fertile plants from seeds obtained from the cross; and

(c) screening progeny seeds from the fertile plants of step (b) for seeds containing reduced saturated fatty acids.

27. (amended) Oil obtained from the plants of Claims [21-26] 24-26.

Please add the following new claims:

28. A method of producing seed oil containing altered levels of unsaturated fatty acids comprising:

(a) transforming a plant oil-producing species with a chimeric gene comprising the nucleic acid fragment encoding all or a portion of a plant delta-12 desaturase;

(b) growing fertile plants from the transformed plant cells of step (a);

(c) screening progeny seeds from the fertile plants of step (b) for the desired levels of unsaturated fatty acids;

(d) processing the progeny seed of step (c) to obtain seed oil containing altered levels of unsaturated fatty acids.

29. The product of the method of Claim 28.

30. Canola oil obtained from the seeds of a plant having a chimeric gene capable of causing altered levels of fatty acid in a transformed canola plant cell, said chimeric gene comprising an isolated nucleic acid fragment comprising a sequence encoding a fatty acid desaturase with an amino acid identity of 50% or greater to the polypeptide encoded by any one of the sequences set forth in SEQ ID NOS: 1, 3, 5, 7, 9, 11 or 15, operably linked to a suitable regulatory sequence.

31. The soybean oil of Claim 9 obtained from the seeds of a plant having a chimeric gene capable of causing altered levels of fatty acid in a transformed soybean plant cell, said chimeric gene comprising an isolated nucleic acid fragment comprising a sequence encoding a fatty acid desaturase wherein the amino acid identity is 60% or greater to the polypeptide encoded by any one of the sequences set forth in SEQ ID NOS: 1, 3, 5, 7, 9, 11 or 15, operably linked to a suitable regulatory sequence.

32. The canola oil of Claim 30 obtained from the seeds of a plant having a chimeric gene capable of causing altered levels of fatty acid in a transformed canola plant cell, said chimeric gene comprising an isolated nucleic acid fragment comprising a sequence encoding a fatty acid desaturase wherein the amino acid identity is 60% or greater to the polypeptide encoded by any one of the sequences set forth in SEQ ID NOS: 1, 3, 5, 7, 9, 11 or 15, operably linked to a suitable regulatory sequence.

33. An isolated nucleic acid fragment comprising a nucleic acid sequence selected from the group consisting of:

(a) a nucleic acid sequence encoding a fatty acid desaturase or a fatty acid desaturase-related plant enzyme with an amino acid identity of 50% or greater to the polypeptide encoded by any one of the sequences set forth in SEQ ID NOS:1, 3, 5, 7, 9, 11, or 15, or

(b) a nucleic acid sequence or a part thereof which is useful in antisense inhibition or sense suppression of endogenous desaturase activity in a transformed plant wherein the nucleic acid has an identity of 80% or greater to any one of the sequences set forth in SEQ ID NOS:1, 3, 5, 7, 9, 11, or 15 or a part thereof.

34. The isolated nucleic acid fragment of Claim 33 wherein the amino acid identity is 60% or greater to the polypeptide encoded by any one of the sequences set forth in SEQ ID NOS:1, 3, 5, 7, 9, 11, or 15.

35. The isolated nucleic acid fragment of Claim 33 wherein the nucleic acid identity is 90% or greater to any one of the sequences set forth in SEQ ID NOS:1, 3, 5, 7, 9, 11, or 15.

36. The isolated nucleic acid fragment of any of Claim 33, 34 or 35 wherein said fragment is isolated from an oil-producing plant species.

37. A chimeric gene comprising the nucleic acid fragment of Claim 33, 34 or 35 operably linked to suitable regulatory sequences.

38. A plant comprising in its genome the chimeric gene of Claim 37.

39. Seeds obtained from the plant of Claim 38.

40. A method of producing seed oil containing altered levels of unsaturated fatty acids comprising:

- (a) transforming a plant cell of an oil-producing species with a chimeric gene of Claim 37;
- (b) growing fertile plants from the transformed plant cells of step (a);
- (c) screening progeny seeds from the fertile plants of step (b) for the desired levels of unsaturated fatty acids; and
- (d) processing the progeny seed of step (c) to obtain seed oil containing altered levels of unsaturated fatty acids.

41. The isolated nucleic acid fragment of Claim 33, 34, or 35 comprising a nucleic acid sequence encoding a plant microsomal delta-12 fatty acid desaturase.

42. A nucleic acid fragment isolated from an oil-producing plant species wherein said fragment is selected from the group consisting of:

- (i) an isolated nucleic acid fragment comprising a nucleic acid sequence encoding an enzyme which catalyzes a reaction at carbon positions 6 and 7 numbered

from the methyl end of an 18 carbon long fatty acyl chain, wherein positions 6 and 7 correspond to carbon positions 12 and 13 numbered from the carbonyl carbon of an 18 carbon long fatty acyl chain and further wherein the amino acid sequence comprising said enzyme contains at least one of the following amino acid sequences selected from the group consisting of: AIPPHCF, AWXXYW, HECGH, LLVPY, WKYSHR, and SHRRHH;

(ii) an isolated nucleic acid fragment encoding an enzyme which catalyzes a reaction at carbon positions 6 and 7 numbered from the methyl end of an 18 carbon long fatty acyl chain wherein positions 6 and 7 correspond to carbon positions 12 and 13 numbered from the carbonyl carbon of an 18 carbon long fatty acyl chain wherein said isolated nucleic acid fragment encodes a protein comprising any one of the amino acid sequences set forth in SEQ ID NOS:2, 4, 6, 8, 10 or 12;

(iii) an isolated nucleic acid fragment encoding an enzyme which catalyzes a reaction at carbon positions 6 and 7 numbered from the methyl end of an 18 carbon long fatty acyl chain, wherein positions 6 and 7 correspond to carbon positions 12 and 13 numbered from the carbonyl carbon of an 18 carbon long fatty acyl chain wherein said isolated nucleic acid fragment hybridizes to the isolated nucleic acid fragment of (ii) under one of the following sets of conditions:

(a) hybridization in 50 mM Tris, pH 7.6, 6X SSC, 5X Denhardt's, 0.5% sodium dodecyl sulfate (SDS), 100 µg denatured calf thymus DNA and 50°C and wash twice with 2X SSC, 0.5% SDS at room temperature for 15 min each, then wash twice with 0.2X SSC, 0.5% SDS at room temperature for 15 min each and then wash twice with 0.2X SSC, 0.5% SDS at 50 °C for 15 min each;

(b) hybridization in 6X SSPE, 5X Denhardt's solution, 0.5% sodium dodecyl sulfate (SDS), 5% dextran sulfate, 100 µg denatured calf thymus DNA at 50°C and wash twice with 2X SSC, 0.5% SDS at room temperature for 15 min each, then wash twice with 0.2X SSC, 0.5% SDS at room temperature for 15 min each and then wash twice with 0.2X SSC, 0.5% SDS at 50 °C for 15 min each; or

(c) hybridization in 50% formamide, 5X SSPE, 1% sodium dodecyl sulfate (SDS), 1% Denhardt's Reagent, 100 µg denatured salmon sperm DNA at 42 °C and wash twice with 2X SSPE, 0.2% SDS at 42° C for 15 min each, then wash twice with 0.2X SSPE, 0.2% SDS at 55 °C for 30 min each.

43. A chimeric gene comprising the nucleic acid fragment of Claim 42 operably linked to suitable regulatory sequences.

44. A plant comprising in its genome the chimeric gene of Claim 43.

45. A method of producing seed oil containing altered levels of unsaturated fatty acids comprising:

- (a) transforming a plant cell of an oil-producing species with a chimeric gene of Claim 43;
- (b) growing fertile plants from the transformed plant cells of step (a);
- (c) screening progeny seeds from the fertile plants of step (b) for the desired levels of unsaturated fatty acids; and
- (d) processing the progeny seed of step (c) to obtain seed oil containing altered levels of unsaturated fatty acids.

46. A method for altering fatty acids composition in seeds comprising:

- (a) making a cross between a mutant line with altered fatty acid composition with a plant containing the chimeric gene of Claim 37;
- (b) growing fertile plants from seeds obtained from the cross; and
- (c) screening progeny seeds from the fertile plants of step (b) for seeds containing altered fatty acid levels.

47. A method for altering fatty acids composition in seeds comprising:

- (a) making a cross between a mutant line with altered fatty acid composition with a plant containing the chimeric gene of Claim 43;
- (b) growing fertile plants from seeds obtained from the cross; and
- (c) screening progeny seeds from the fertile plants of step (b) for seeds containing altered fatty acid levels.

48. A method for reducing polyunsaturated fatty acids in rapeseed oil comprising:

- (a) making a cross between a rapeseed variety with increased oleic acid content or reduced linolenic acid content with a plant containing the chimeric gene of Claim 37;
- (b) growing fertile plants from seeds obtained from the cross; and
- (c) screening progeny seeds from the fertile plants of step (b) for seeds containing reduced polyunsaturated fatty acids.

49. A method for reducing polyunsaturated fatty acids in rapeseed oil comprising:



(a) making a cross between a rapeseed variety with increased oleic acid content or reduced linolenic acid content with a plant containing the chimeric gene of Claim 43;

(b) growing fertile plants from seeds obtained from the cross; and

(c) screening progeny seeds from the fertile plants of step (b) for seeds containing reduced polyunsaturated fatty acids.

50. The method of Claim 46 or 47 wherein the cross in (a) is between a progeny plant derived from a seed comprising a Brassica variety having an oleic acid content of about 69% to 77%, based upon total extractable oil and belonging to a line in which the said oleic acid content has been stabilized for both the generation to which the seed belongs and its parent generation.

51. A method for reducing saturated fatty acids in rapeseed seeds comprising:

(a) making a cross between a rapeseed variety with increased oleic acid content with a plant containing the chimeric gene of Claim 37;

(b) growing fertile plants from seeds obtained from the cross; and

(c) screening progeny seeds from the fertile plants of step (b) for seeds containing reduced saturated fatty acids.

52. A method for reducing saturated fatty acids in rapeseed seeds comprising:

(a) making a cross between a rapeseed variety with increased oleic acid content with a plant containing the chimeric gene of Claim 49;

(b) growing fertile plants from seeds obtained from the cross; and

(c) screening progeny seeds from the fertile plants of step (b) for seeds containing reduced saturated fatty acids.

53. A method for reducing polyunsaturated fatty acids in soybean oil comprising:

(a) making a cross between a soybean variety with increased oleic acid content or reduced linolenic acid content with a plant containing the chimeric gene of Claim 37;

(b) growing fertile plants from seeds obtained from the cross; and

(c) screening progeny seeds from the fertile plants of step (b) for seeds containing reduced polyunsaturated fatty acids.

54. A method for reducing polyunsaturated fatty acids in soybean oil comprising:

096979-103600

(a) making a cross between a soybean variety with increased oleic acid content or reduced linolenic acid content with a plant containing the chimeric gene of Claim 43;

(b) growing fertile plants from seeds obtained from the cross; and

(c) screening progeny seeds from the fertile plants of step (b) for seeds containing reduced polyunsaturated fatty acids.

55. A method for reducing saturated fatty acids in soybean seeds comprising:

(a) making a cross between a soybean variety with increased oleic acid content with a plant containing the chimeric gene of Claim 37;

(b) growing fertile plants from seeds obtained from the cross; and

(c) screening progeny seeds from the fertile plants of step (b) for seeds containing reduced saturated fatty acids.

56. A method for reducing saturated fatty acids in soybean seeds comprising:

(a) making a cross between a soybean variety with increased oleic acid content with a plant containing the chimeric gene of Claim 43;

(b) growing fertile plants from seeds obtained from the cross; and

(c) screening progeny seeds from the fertile plants of step (b) for seeds containing reduced saturated fatty acids.

57. Oil obtained from the seeds of a soybean plant having an oleic acid content of at least 75%.

58. Oil obtained from the seeds of a soybean plant having an oleic acid content of at least 70% and a linoleic acid content less than 14%.

59. The oil of Claim 58 in which the linoleic acid content is less than 10%.--

In the Specification

Kindly amend the specification by adding after the title at line 3:

--This is a divisional of application number 09/133,962, pending, filed August 14, 1998, which is a continuation of application number 08/262,401, filed June 20, 1994, now abandoned, which was a national filing continuation-in-part of PCT/US93/09987, filed October 15, 1993, which was a continuation-in-part of 07/977,339, filed November 17, 1992, now abandoned.--

09/133,962, pending, filed August 14, 1998, which is a continuation of application number 08/262,401, filed June 20, 1994, now abandoned, which was a national filing continuation-in-part of PCT/US93/09987, filed October 15, 1993, which was a continuation-in-part of 07/977,339, filed November 17, 1992, now abandoned.--

Kindly amend the specification as follows:

Page 13, line 5: please delete "1394" and substitute therefor --1426--.

Page 13, line 7: please delete "99 to 101" and substitute therefor --130 to 132-

-.  
-

Page 13, lines 7-8: please delete "1248 to 1250" and substitute therefor --1282 to 1284--.

Page 13, line 10: please delete "99 to 1250" and substitute therefor --130 to 1284--.

Page 13, lines 10-11: please delete "1 to 98 and 1251 to 1394" and substitute therefor --1 to 129 and 1285 to 1426--.

Page 13, line 13: please delete "383" and substitute therefor --384--.

Page 13, lines 14-15: please delete "99 to 1250" and substitute therefor --130 to 1284--.

Page 36, line 8: please delete "1394" and substitute therefor --1426--.

Page 36, lines 10: please delete "99 to 101" and substitute therefor --130 to 132--and delete "1248 to 1250" and substitute therefor --1282 to 1284--.

Page 36, line 13: please delete "99 to 1250" and substitute therefor --130 to 1284--and delete "1 to 98 and 1251 to 1394" and substitute therefor --1 to 129 and 1285 to 1426--.

Page 36, lines 21: please delete "both"; after "nucleotide" insert --level--; and after "and" insert --90% at the--.

Page 36, line 22: please delete "levels" and insert --level--.

#### Remarks

This case is a divisional application under 37 CFR §1.53(b).

Claim 9 has been amended to recite oil obtained from the seeds of a soybean plant having a chimeric gene comprising a nucleic acid fragment encoding a fatty acid desaturase with an amino acid identity of 50% or greater to the polypeptide encoded by any one of the sequences set forth in SEQ ID NOS: 1, 3, 5, 7, 9, 11 or 15. Support for this clarification can be found in the specification on pages 44 - 48 and in the claims as originally filed. Thus, no new matter has been added.

Claim 10 has been clarified by replacing the term "unsaturated" with --hydroxylated and it has been rewritten to be in independent form. Support for the term "hydroxylated" can be found in the specification on page 9 at lines 8-18 of the specification. Thus, no new matter has been added.

096974576960

Claims 15, 16, 18, 19 and 20 have been amended by reciting alternate hybridization conditions. Support for the three different sets of alternative hybridization and wash conditions can be found in the specification on page 35 at lines 27-32, on page 70, on page 73 at lines 5-11 and on pages 74 and 75. Thus, no new matter has been added.

New claims 28 - 59 have been added.

Claim 28 is similar to original claim 10 except that it is written in independent form.

Support for claim 30 can be found in the specification on pages 44-48 and in Example 3.

Support for claims 31, 32, 34 can be found in the specification on pages 44-48 and, in particular, on page 46.

Support for claim 33 can be found in claim 1 as originally filed and in the specification on pages 54-55.

Support for claim 35 can be found in claim 3 as originally filed.

Support for claims 37-39 can be found throughout the specification and claims as originally filed.

Support for claim 40 can be found in claim 10 as originally filed.

Support for claim 41 can be found on page 6 of the specification.

Support for claim 42 can be found in Table 7 on pages 47 and 48 of the specification and on page 35 at lines 27-32, on page 70, on page 73 at lines 5-11 and on pages 74 and 75.

Support for claims 43-56 can be found in the specification and claims as originally filed.

Support for claims 57-59 can be found in the specification in Table 13A on page 100-119.

Thus, it can be seen that no new matter has been added.

Substituted herewith pursuant to 37 CFR §1.825(a) are substitute sheets for the sequence listing provided as set forth in 37 CFR §1.821 (c). These sheets and a substitute copy of the computer readable form are enclosed herewith. The computer readable form corresponds identically to the substitute sequence listing pages submitted with this response. Support for this amendment can be found in the specification.

Specifically, the sequence listings are the same except for SEQ ID NOS: 3 and 4 which are being clarified due to sequencing errors. Support for this clarification can be found on page 36 at lines 24-27 which shows that the material has now been correctly sequenced, pCF2-165D, and was deposited with the ATCC on October 16,

009201" 5225950

Application No.: UNKNOWN  
Docket No.: BB1043 US NA DIV


1992 under the provisions of the Budapest Treaty and bears accession number ATCC 69094. Accordingly, no new matter has been added.

The clarifications in the specification were necessitated due to the clarifications in the sequence listings.

Enclosed herewith along with this Preliminary Amendment is an Information Disclosure Statement setting forth all references which had been cited by Applicants or the Examiner in connection with Serial No. 09/133,912 and some additional information as well as a petition for a three (3) month extension of time. Also, a copy of U.S. Patent No. 5,534,425 which issued to Fehr et al. on July 9, 1996 accompanies this preliminary amendment.

Please charge any fees or credit any overpayment of fees which are required in connection with the filing of this Preliminary Amendment, Information Disclosure Statement and Petition for Extension of Time to Deposit Account No. 04-1928 (E. I. du Pont de Nemours and Company).

Respectfully submitted,



Lynne M. Christenbury  
Attorney for Applicants  
Registration No. 30,971  
Telephone: 302-992-5481

Enclosures

0969779-102600

TITLEGENES FOR MICROSOMAL DELTA-12 FATTY ACID  
DESATURASES AND RELATED ENZYMES FROM PLANTSFIELD OF THE INVENTION

5       The invention relates to the preparation and use of  
nucleic acid fragments encoding fatty acid desaturase  
enzymes to modify plant lipid composition. Chimeric  
genes incorporating such nucleic acid fragments and  
suitable regulatory sequences may be used to create  
10   transgenic plants with altered levels of unsaturated  
fatty acids.

BACKGROUND OF THE INVENTION

Plant lipids have a variety of industrial and  
nutritional uses and are central to plant membrane  
15   function and climatic adaptation. These lipids  
represent a vast array of chemical structures, and these  
structures determine the physiological and industrial  
properties of the lipid. Many of these structures  
result either directly or indirectly from metabolic  
20   processes that alter the degree of unsaturation of the  
lipid. Different metabolic regimes in different plants  
produce these altered lipids, and either domestication  
of exotic plant species or modification of agronomically  
adapted species is usually required to economically  
25   produce large amounts of the desired lipid.

Plant lipids find their major use as edible oils in  
the form of triacylglycerols. The specific performance  
and health attributes of edible oils are determined  
largely by their fatty acid composition. Most vegetable  
30   oils derived from commercial plant varieties are  
composed primarily of palmitic (16:0), stearic (18:0),  
oleic (18:1), linoleic (18:2) and linolenic (18:3)  
acids. Palmitic and stearic acids are, respectively,  
16- and 18-carbon-long, saturated fatty acids. Oleic,  
35   linoleic, and linolenic acids are 18-carbon-long,

009697379 102600

5

TABLE 1

- Canola
- Soybean
- Corn
- Peanut
- Safflower
- Sunflower
- Cotton

10

20

5 development of disagreeable odors and flavors  
(Gailliard, 1980, Vol. 4, pp. 85-116 In: Stumpf, P. K.,  
Ed., The Biochemistry of Plants, Academic Press, New  
York). The levels of poly-unsaturates may be reduced by  
hydrogenation, but the expense of this process and the  
0 concomitant production of nutritionally questionable  
trans isomers of the remaining unsaturated fatty acids  
reduces the overall desirability of the hydrogenated oil  
(Mensink et al., New England J. Medicine (1990) N323:  
439-445). Similar problems exist with soybean and corn  
5 oils.

20 nutritional supplements, for example in baby foods.

25 high (85%) oleic sunflower and low (2%) linolenic flax  
(Knowles, (1980) pp. 35-38 In: Applewhite, T. H., Ed.,  
World Conference on Biotechnology for the Fats and Oils  
Industry Proceedings, American Oil Chemists' Society).  
Similar commercial progress with the other plants shown  
30 in Table 1 has been largely elusive due to the difficult  
nature of the procedure and the pleiotropic effects of  
the mutational regime on plant hardiness and yield  
potential.

The biosynthesis of the major plant lipids has been  
35 the focus of much research (Browse et al., Ann. Rev.



Plant Physiol. Mol. Biol. (1991) 42:467-506). These studies show that, with the notable exception of the soluble stearyl-acyl carrier protein desaturase, the controlling steps in the production of unsaturated fatty acids are largely catalyzed by membrane-associated fatty acid desaturases. Desaturation reactions occur in plastids and in the endoplasmic reticulum using a variety of substrates including galactolipids, sulfolipids, and phospholipids. Genetic and physiological analyses of Arabidopsis thaliana nuclear mutants defective in various fatty acid desaturation reactions indicates that most of these reactions are catalyzed by enzymes encoded at single genetic loci in the plant. The analyses show further that the different defects in fatty acid desaturation can have profound and different effects on the ultra-structural morphology, cold sensitivity, and photosynthetic capacity of the plants (Ohlrogge, et al., Biochim. Biophys. Acta (1991) 1082:1-26). However, biochemical characterization of the desaturase reactions has been meager. The instability of the enzymes and the intractability of their proper assay has largely limited researchers to investigations of enzyme activities in crude membrane preparations. These investigations have, however, demonstrated the role of delta-12 desaturase and delta-15 desaturase activities in the production of linoleate and linolenate from 2-oleoyl-phosphatidylcholine and 2-linoleoyl-phosphatidylcholine, respectively (Wang et al., Plant Physiol. Biochem. (1988) 26:777-792). Thus, modification of the activities of these enzymes represents an attractive target for altering the levels of lipid unsaturation by genetic engineering.

Nucleotide sequences encoding microsomal delta-9 stearyl-coenzyme-A desaturases from yeast, rat, and

09697375-102600

mice have been described (Stukey, et al., J. Biol. Chem. (1990) 265:20144-20149; Thiede, et al., J. Biol. Chem. (1986) 261:13230-13235; Kaestner, et al., J. Biol. Chem. (1989) 264:14755-1476). Nucleotide sequences

5 encoding soluble delta-9 stearoyl-acyl carrier protein desaturases from higher plants have also been described (Thompson, et al., Proc. Natl. Acad. Sci. U.S.A. (1991) 88:2578-2582; Shanklin et al., Proc. Natl. Acad. Sci. USA (1991) 88:2510-2514). A nucleotide sequence from

10 coriander plant encoding a soluble fatty acid desaturase, whose deduced amino acid sequence is highly identical to that of the stearoyl-acyl carrier protein desaturase and which is responsible for introducing the double bond in petroselinic fatty acid (18:1, 6c), has

15 also been described [Cahoon, et. al. (1992) Proc. Natl. Acad. Sci. U.S.A. 89:11184-11188]. Two fatty acid desaturase genes from the cyanobacterium, Synechocystis PCC6803, have been described: one encodes a fatty acid desaturase, designated *des A*, that catalyzes the

20 conversion of oleic acid at the sn-1 position of galactolipids to linoleic acid [Wada, et al., Nature (1990) 347:200-203]; another encodes a delta-6 fatty acid desaturase that catalyzes the conversion of

25 linoleic acid at the sn-1 position of galactolipids to  $\gamma$ -linolenic acid (18:2, 6c,9c) [WO 9306712]. Nucleotide sequences encoding higher plant membrane-bound microsomal and plastid delta-15 fatty acid desaturases have also been described [WO 9311245]; Arondel, V. et. al. (1992) Science 258:1353-1355]. There is no report

30 of the isolation of higher plant genes encoding fatty acid desaturases other than the soluble delta-6 and delta-9 desaturases and the membrane-bound (microsomal and plastid) delta-15 desaturases. While there is extensive amino acid sequence identity between the

35 soluble desaturases and significant amino acid sequence

009207 546950

identity between the higher plant microsomal and plastid delta-15 desaturases, there is no significant homology between the soluble and the membrane-bound desaturases. Sequence-dependent protocols based on the sequences encoding delta-15 desaturases have been unsuccessful in cloning sequences for microsomal delta-12 desaturase. For example, nucleotide sequences of microsomal or plastid delta-15 desaturases as hybridization probes have been unsuccessful in isolating a plant microsomal delta-12 desaturase clone. Furthermore, while we have used a set of degenerate oligomers made to a stretch of 12 amino acids, which is identical in all plant delta-15 desaturases and highly conserved (10/12) in the cyanobacterial *des A* desaturase, as a hybridization probe to isolate a higher plant nucleotide sequence encoding plastid delta-12 fatty acid desaturase, this method has been unsuccessful in isolating the microsomal delta-12 desaturase cDNAs. Furthermore, there has been no success in isolating the microsomal delta-12 desaturase by using the polymerase chain reaction products derived from plant DNA, plant RNA or plant cDNA library using PCR primers made to stretches of amino acids that are conserved between the higher plant delta-15 and *des A* desaturases. Thus, there are no teachings which enable the isolation of plant microsomal delta-12 fatty acid desaturases or plant fatty acid desaturase-related enzymes. Furthermore, there is no evidence for a method to control the the level of delta-12 fatty acid desaturation or hydroxylation in plants using nucleic acids encoding delta-12 fatty acid desaturases or hydroxylases.

The biosynthesis of the minor plant lipids has been less well studied. While hundreds of different fatty acids have been found, many from the plant kingdom, only a tiny fraction of all plants have been surveyed for

009201" 52E26950

their lipid content (Gunstone, et al., Eds., (1986) The Lipids Handbook, Chapman and Hall Ltd., Cambridge).

Accordingly, little is known about the biosynthesis of these unusual fatty acids and fatty acid derivatives.

- 5 Interesting chemical features found in such fatty acids include, for example, allenic and conjugated double bonds, acetylenic bonds, trans double bonds, multiple double bonds, and single double bonds in a wide number of positions and configurations along the fatty acid
- 10 chain. Similarly, many of the structural modifications found in unusual lipids (e.g., hydroxylation, epoxidation, cyclization, etc.) are probably produced via further metabolism following chemical activation of the fatty acid by desaturation or they involve a
- 15 chemical reaction that is mechanistically similar to desaturation. Many of these fatty acids and derivatives having such features within their structure could prove commercially useful if an agronomically viable species could be induced to synthesize them by introduction of a
- 20 gene encoding the appropriate desaturase. Of particular interest are vegetable oils rich in 12-hydroxyoctadeca-9-enoic acid (ricinoleic acid). Ricinoleic acid and its derivatives are widely used in the manufacture of lubricants, polymers, cosmetics, coatings and
- 25 pharmaceuticals (e.g., see Gunstone, et al., Eds., (1986) The Lipids Handbook, Chapman and Hall Ltd., Cambridge). The only commercial source of ricinoleic acid is castor oil and 100% of the castor oil used by the U.S. is derived from beans grown elsewhere in the
- 30 world, mainly Brazil. Ricinoleic acid in castor beans is synthesized by the addition of an hydroxyl group at the delta-12 position of oleic acid (Galliard & Stumpf (1966) J. Biol. Chem. 241: 5806-5812). This reaction resembles the initial reaction in a possible mechanism
- 35 for the desaturation of oleate at the delta-12 position

009697379 102600

to linoleate since dehydration of 12-hydroxyoctadeca-9-enoic acid, by an enzyme activity analogous to the hydroxydecanoyl dehydrase of *E. coli* (Cronan et al. (1988) J. Biol. Chem. 263:4641-4646), would result in the formation of linoleic acid. Evidence for the hydroxylation reaction being part of a general mechanism of enzyme-catalyzed desaturation in eukaryotes has been obtained by substituting a sulfur atom in the place of carbon at the delta-9 position of stearic acid. When incubated with yeast cell extracts the thiostearate was converted to a 9-sulfoxide (Buist et al. (1987) Tetrahedron Letters 28:857-860). This sulfoxidation was specific for sulfur at the delta-9 position and did not occur in a yeast delta-9-desaturase deficient mutant (Buist & Marecak (1991) Tetrahedron Letters 32:891-894). The 9-sulfoxide is the sulfur analogue of 9-hydroxyoctadecastearate, the proposed intermediate of stearate desaturation.

Hydroxylation of oleic acid to ricinoleic acid in castor bean cells, like microsomal desaturation of oleate in plants, occurs at the delta-12 position of the fatty acid at the sn-2 position of phosphatidylcholine in microsomes (Bafor et al. (1991) Plant Physiol 280:507-514). Furthermore, castor oleate delta-12 hydroxylation and plant oleate microsomal delta-12 desaturation are both inhibited by iron chelators and require molecular oxygen [Moreau & Stumpf (1981) Plant Physiology 67:672-676; Somerville, C. (1992) MSU-DOE Plant Research Laboratory Annual Report]. These biochemical similarities in conjunction with the observation that antibodies raised against cytochrome b<sub>5</sub> completely inhibit the activities of both oleate delta-12 desaturation in safflower microsomes and oleate delta-12 hydroxylase in castor microsomes [Somerville, C. (1992) MSU-DOE Plant Research Laboratory Annual

Report] comprise strong evidence that the hydroxylase and the desaturase are functionally related. It seems reasonable to assume, therefore, that the nucleotide sequence encoding a plant delta-12 desaturase would be  
5 useful in cloning the oleate hydroxylase gene from castor by sequence-dependent protocols. For example, by screening a castor DNA library with oligomers based on amino acid regions conserved between delta-12 desaturases, or regions conserved between delta-12 and  
10 other desaturases, or with oligomers based on amino acids conserved between delta-12 desaturases and known membrane-associated hydroxylases. It would be more efficient to isolate the castor oleate hydroxylase cDNA by combining the sequence dependent protocols with a  
15 "differential" library approach. One example of such a difference library would be based on different stages of castor seed development, since ricinoleic acid is not synthesized by very young castor seeds (less than 12 DAP, corresponding to stage I and stage II seeds in  
20 the scheme of Greenwood & Bewley, Can. J. Bot. (1982) 60:1751-1760), in the 20 days following these early stages the relative ricinoleate content increases from 0% to almost 90% of total seed fatty acids (James et al. Biochem. J. (1965) 95:448-452, Canvin. Can. J. Biochem.  
25 Physiol. (1963) 41:1879-1885). Thus it would be possible to make a cDNA "difference" library made from mRNA present in a stage when ricinoleic acid was being synthesized at a high rate but from which mRNA present in earlier stages was removed. For the earlier stage  
30 mRNA, a stage such as stage II (10 DAP) when ricinoleic acid is not being made but when other unsaturated fatty acids are, would be appropriate. The construction of libraries containing only differentially expressed genes is well known in the art (Sargent. Meth. Enzymol. (1987)  
35 152:423-432). Assembly of the free ricinoleic acid, via

ricinoleoyl-CoA, into triacylglycerol is readily catalyzed by canola and safflower seed microsomes (Bafor et al., Biochem J. (1991) 280:507-514, Wiberg et al. 10th International Symposium on the Metabolism, Structure & Function of Plant Lipids (1992), Jerba, Tunisia) and ricinoleic acid is removed from phosphatidylcholine by a lipase common to all oilseeds investigated. Thus, expression of the castor bean oleate hydroxylase gene in oil crops, such as canola seeds and soybeans, would be expected to result in an oil rich in triglycerides containing ricinoleic acid.

#### SUMMARY OF THE INVENTION

Applicants have discovered a means to control the nature and levels of unsaturated fatty acids in plants. Nucleic acid fragments from cDNAs or genes encoding fatty acid desaturases are used to create chimeric genes. The chimeric genes may be used to transform various plants to modify the fatty acid composition of the plant or the oil produced by the plant. More specifically, one embodiment of the invention is an isolated nucleic acid fragment comprising a nucleotide sequence encoding a fatty acid desaturase or a fatty acid desaturase-related enzyme with an amino acid identity of 50%, 60%, 90% or greater respectively to the polypeptide encoded by SEQ ID NOS:1, 3, 5, 7, 9, 11, or 15. Most specifically, the invention pertains to a gene sequence for plant microsomal delta-12 fatty acid desaturase or desaturase-related enzyme. The plant in this embodiment may more specifically be soybean, oilseed Brassica species, Arabidopsis thaliana, castor, and corn.

Another embodiment of this invention involves the use of these nucleic acid fragments in sequence-dependent protocols. Examples include use of the fragments as hybridization probes to isolate nucleotide

09697379 102600

5 desaturases or fatty acid desaturase-related enzymes.

10 NOS:1, 3, 5, 7, 9, or 15 encoding fatty acid desaturases  
or SEQ ID NOS:11 encoding a desaturase or desaturase-  
related enzyme operably-linked in suitable orientation  
to suitable regulatory sequences. Preferred are those  
chimeric genes which incorporate nucleic acid fragments  
15 encoding microsomal delta-12 fatty acid desaturase or  
desaturase-related enzymes.

20 a plant cell with a chimeric gene described above;  
(b) growing sexually mature plants from the transformed  
plant cells of step (a); (c) screening progeny seeds  
from the sexually mature plants of step (b) for the  
desired levels of unsaturated fatty acids, and  
25 (d) processing the progeny seed of step (c) to obtain  
seed oil containing altered levels of the unsaturated  
fatty acids. Preferred plant cells and oils are derived  
from soybean, rapeseed, sunflower, cotton, cocoa,  
peanut, safflower, coconut, flax, oil palm, and corn.

30 Preferred methods of transforming such plant cells would include the use of Ti and Ri plasmids of Agrobacterium, electroporation, and high-velocity ballistic bombardment.

35 breeding to obtain altered levels of oleic acids in the



seed oil of oil producing plant species. This method involves (a) making a cross between two varieties of oil producing plant species differing in the oleic acid trait; (b) making a Southern blot of restriction enzyme digested genomic DNA isolated from several progeny plants resulting from the cross; and (c) hybridizing the Southern blot with the radiolabelled nucleic acid fragments encoding the fatty acid desaturases or desaturase-related enzymes.

The invention is also embodied in a method of RFLP mapping that uses the isolated microsomal delta-12 desaturase cDNA or related genomic fragments described herein.

The invention is also embodied in plants capable of producing altered levels of fatty acid desaturase by virtue of containing the chimeric genes described herein. Further, the invention is embodied by seed oil obtained from such plants.

#### BRIEF DESCRIPTION OF THE SEQUENCE DESCRIPTIONS

The invention can be more fully understood from the following detailed description and the Sequence Descriptions which form a part of this application. The Sequence Descriptions contain the three letter codes for amino acids as defined in 37 C.F.R. 1.822 which are incorporated herein by reference.

SEQ ID NO:1 shows the 5' to 3' nucleotide sequence of 1372 base pairs of the Arabidopsis thaliana cDNA which encodes microsomal delta-12 desaturase. Nucleotides 93-95 and nucleotides 1242-1244 are, respectively, the putative initiation codon and the termination codon of the open reading frame (nucleotides 93-1244). Nucleotides 1-92 and 1245-1372 are, respectively, the 5' and 3' untranslated nucleotides.

009975 548590

5 <sup>1476</sup> SEQ ID NO:3 shows the 5' to 3' nucleotide sequence  
of 1394 base pairs of the Brassica napus cDNA which  
encodes microsomal delta-12 desaturase in plasmid  
pCF2-165d. Nucleotides 99 to 101 and nucleotides 1248  
to 1250 are, respectively, the putative initiation codon  
and the termination codon of the open reading frame  
10 (nucleotides 99 to 1250). Nucleotides 1 to 98 and 1251  
to 1394 are, respectively, the 5' and 3' untranslated  
nucleotides.

SEQ ID NO:5 shows the 5' to 3' nucleotide sequence of 1369 base pairs of soybean (Glycine max) cDNA which encodes microsomal delta-12 desaturase in plasmid pSF2-169K. Nucleotides 108 to 110 and nucleotides 1245 to 1247 are, respectively, the putative initiation codon and the termination codon of the open reading frame (nucleotides 108 to 1247). Nucleotides 1 to 107 and 1248 to 1369 are, respectively, the 5' and 3' untranslated nucleotides.

SEQ ID NO:7 shows the 5' to 3' nucleotide sequence of 1790 base pairs of corn (Zea mays) cDNA which encodes microsomal delta-12 desaturase in plasmid pFad2#1. Nucleotides 165 to 167 and nucleotides 1326 to 1328 are, respectively, the putative initiation codon and the termination codon of the open reading frame (nucleotides 164 to 1328). Nucleotides 1 to 163 and 1329 to 1790

are, respectively, the 5' and 3' untranslated nucleotides.

SEQ ID NO:8 is the 387 amino acid protein sequence deduced from the open reading frame (nucleotides 164 to 1328) in SEQ ID NO:7.

SEQ ID NO:9 shows the 5' to 3' nucleotide sequence of 673 base pairs of castor (Ricinus communis) incomplete cDNA which encodes part of a microsomal delta-12 desaturase in plasmid pRF2-1C. The sequence encodes an open reading frame from base 1 to base 673..

SEQ ID NO:10 is the 219 amino acid protein sequence deduced from the open reading frame (nucleotides 1 to 657) in SEQ ID NO:9.

SEQ ID NO:11 shows the 5' to 3' nucleotide sequence of 1369 base pairs of castor (Ricinus communis) cDNA which encodes part of a microsomal delta-12 desaturase or desaturase-related enzyme in plasmid pRF197C-42. Nucleotides 184 to 186 and nucleotides 1340 to 1342 are, respectively, the putative initiation codon and the termination codon of the open reading frame (nucleotides 184 to 1347). Nucleotides 1 to 183 and 1348 to 1369 are, respectively, the 5' and 3' untranslated nucleotides.

SEQ ID NO:12 is the 387 amino acid protein sequence deduced from the open reading frame (nucleotides 184 to 1342) in SEQ ID NO:11.

SEQ ID NO:13 is the sequence of a set of 64-fold degenerate 26 nucleotide-long oligomers, designated NS3, made to conserved amino acids 101-109 of SEQ ID NO:2, designed to be used as sense primers in PCR to isolate novel sequences encoding microsomal delta-12 desaturases or desaturase-like enzymes.

SEQ ID NO:14 is the sequence of a set of 64-fold degenerate and 26 nucleotide-long oligomers, designated NS9, which is made to conserved amino acids 313-321 of

SEQ ID NO:2 and designed to be used as antisense primers in PCR to isolate novel sequences encoding microsomal delta-12 desaturases or desaturase-like enzymes.

SEQ ID NO:15 shows the 5' to 3' nucleotide sequence of 2973 bp of Arabidopsis thaliana genomic fragment containing the microsomal delta-12 desaturase gene contained in plasmid pAGF2-6. Its nucleotides 433 and 2938 correspond to the start and end, respectively, of SEQ ID NO:1. Its nucleotides 521 to 1654 are the 1134 bp intron.

SEQ ID NO:16 is the sequence of a set of 256-fold degenerate and 25 nucleotide-long oligomers, designated RB5a, which is made to conserved amino acids 318-326 of SEQ ID NO:2 and designed to be used as antisense primers in PCR to isolate novel sequences encoding microsomal delta-12 desaturases or desaturase-like enzymes.

SEQ ID NO:17 is the sequence of a set of 128-fold degenerate and 25 nucleotide-long oligomers, designated RB5b, which is made to conserved amino acids 318-326 of SEQ ID NO:2 and designed to be used as antisense primers in PCR to isolate novel sequences encoding microsomal delta-12 desaturases or desaturase-like enzymes.

#### DETAILED DESCRIPTION OF THE INVENTION

Applicants have isolated nucleic acid fragments that encode plant fatty acid desaturases and that are useful in modifying fatty acid composition in oil-producing species by genetic transformation.

Thus, transfer of the nucleic acid fragments of the invention or a part thereof that encodes a functional enzyme, along with suitable regulatory sequences that direct the transcription of their mRNA, into a living cell will result in the production or over-production of plant fatty acid desaturases and will result in increased levels of unsaturated fatty acids in cellular lipids, including triacylglycerols.

09697379-102600

Transfer of the nucleic acid fragments of the invention or a part thereof, along with suitable regulatory sequences that direct the transcription of their antisense RNA, into plants will result in the inhibition of expression of the endogenous fatty acid desaturase that is substantially homologous with the transferred nucleic acid fragment and will result in decreased levels of unsaturated fatty acids in cellular lipids, including triacylglycerols.

Transfer of the nucleic acid fragments of the invention or a part thereof, along with suitable regulatory sequences that direct the transcription of their mRNA, into plants may result in inhibition by cosuppression of the expression of the endogenous fatty acid desaturase gene that is substantially homologous with the transferred nucleic acid fragment and may result in decreased levels of unsaturated fatty acids in cellular lipids, including triacylglycerols.

The nucleic acid fragments of the invention can also be used as restriction fragment length polymorphism (RFLP) markers in plant genetic mapping and plant breeding programs.

The nucleic acid fragments of the invention or oligomers derived therefrom can also be used to isolate other related fatty acid desaturase genes using DNA, RNA, or a library of cloned nucleotide sequences from the same or different species by well known sequence-dependent protocols, including, for example, methods of nucleic acid hybridization and amplification by the polymerase chain reaction.

#### Definitions

In the context of this disclosure, a number of terms shall be used. Fatty acids are specified by the number of carbon atoms and the number and position of the double bond: the numbers before and after the colon

009201 545950

refer to the chain length and the number of double bonds, respectively. The number following the fatty acid designation indicates the position of the double bond from the carboxyl end of the fatty acid with the "c" affix for the *cis*-configuration of the double bond. For example, palmitic acid (16:0), stearic acid (18:0), oleic acid (18:1,9c), petroselinic acid (18:1, 6c), linoleic acid (18:2,9c,12c),  $\gamma$ -linolenic acid (18:3, 6c,9c,12c) and  $\alpha$ -linolenic acid (18:3, 9c,12c,15c).

Unless otherwise specified 18:1, 18:2 and 18:3 refer to oleic, linoleic and linolenic fatty acids. Ricinoleic acid refers to an 18 carbon fatty acid with a *cis*-9 double bond and a 12-hydroxyl group. The term "fatty acid desaturase" used herein refers to an enzyme which catalyzes the breakage of a carbon-hydrogen bond and the introduction of a carbon-carbon double bond into a fatty acid molecule. The fatty acid may be free or esterified to another molecule including, but not limited to, acyl-carrier protein, coenzyme A, sterols and the glycerol moiety of glycerolipids. The term "glycerolipid desaturases" used herein refers to a subset of the fatty acid desaturases that act on fatty acyl moieties esterified to a glycerol backbone. "Delta-12 desaturase" refers to a fatty acid desaturase that catalyzes the formation of a double bond between carbon positions 6 and 7 (numbered from the methyl end), (i.e., those that correspond to carbon positions 12 and 13 (numbered from the carbonyl carbon) of an 18 carbon-long fatty acyl chain. "Delta-15 desaturase" refers to a fatty acid desaturase that catalyzes the formation of a double bond between carbon positions 3 and 4 (numbered from the methyl end), (i.e., those that correspond to carbon positions 15 and 16 (numbered from the carbonyl carbon) of an 18 carbon-long fatty acyl chain. Examples of fatty acid desaturases include, but are not limited

00697379 102600

would be particularly useful. The consensus sequences will also take into account conservative substitutions known to one skilled in the art, such as Lys/Arg, Glu/Asp, Ile/Val/Leu/Met, Ala/Gly, Gln/Asn, and Ser/Thr.

5 Amino acid sequences as short as four amino acids long can successfully be used in PCR [Nunberg et. al. (1989) Journal of Virology 63:3240-3249]. Amino acid sequences conserved between delta-12 desaturases (SEQ ID NOS:2, 4, 6, 8, and 10) may also be used in sequence-dependent

10 protocols to isolate fatty acid desaturases and fatty acid desaturase-related enzymes expected to be more related to delta-12 desaturases, such as the oleate hydroxylase from castor bean. Particularly useful are conserved sequences in column 3 (Table 7), since they

15 are also conserved well with delta-15 desaturases (column 4, Table 7).

Determining the conserved amino acid sequences from diverse desaturases will also allow one to identify more and better consensus sequences that will further aid in

20 the isolation of novel fatty acid desaturases, including those from non-plant sources such as fungi, algae (including the desaturases involved in the desaturations of the long chain n-3 fatty acids), and even cyano-bacteria, as well as other membrane-associated

25 desaturases from other organisms.

The function of the diverse nucleotide fragments encoding fatty acid desaturases or desaturase-related enzymes that can be isolated using the present invention can be identified by transforming plants with the

30 isolated sequences, linked in sense or antisense orientation to suitable regulatory sequences required for plant expression, and observing the fatty acid phenotype of the resulting transgenic plants. Preferred target plants for the transformation are the same as the

35 source of the isolated nucleotide fragments when the

goal is to obtain inhibition of the corresponding endogenous gene by antisense inhibition or cosuppression. Preferred target plants for use in expression or overexpression of the isolated nucleic acid fragments are wild type plants or plants with known mutations in desaturation reactions, such as the Arabidopsis mutants fadA, fadB, fadC, fadD, fad2, and fad3; mutant flax deficient in delta-15 desaturation; or mutant sunflower deficient in delta-12 desaturation. Alternatively, the function of the isolated nucleic acid fragments can be determined similarly via transformation of other organisms, such as yeast or cyanobacteria, with chimeric genes containing the nucleic acid fragment and suitable regulatory sequences followed by analysis of fatty acid composition and/or enzyme activity.

#### Overexpression of the Fatty Acid

##### Desaturase Enzymes in Transgenic Species

The nucleic acid fragment(s) of the instant invention encoding functional fatty acid desaturase(s), with suitable regulatory sequences, can be used to overexpress the enzyme(s) in transgenic organisms. An example of such expression or overexpression is demonstrated by transformation of the Arabidopsis mutant lacking oleate desaturation. Such recombinant DNA constructs may include either the native fatty acid desaturase gene or a chimeric fatty acid desaturase gene isolated from the same or a different species as the host organism. For overexpression of fatty acid desaturase(s), it is preferable that the introduced gene be from a different species to reduce the likelihood of cosuppression. For example, overexpression of delta-12 desaturase in soybean, rapeseed, or other oil-producing species to produce altered levels of polyunsaturated fatty acids may be achieved by expressing RNA from the full-length cDNA found in p92103, pCF2-165D, and



09697379 "102600

pSF2-169K. Transgenic lines overexpressing the delta-12 desaturase, when crossed with lines overexpressing delta-15 desaturases, will result in ultrahigh levels of 18:3. Similarly, the isolated nucleic acid fragments encoding fatty acid desaturases from Arabidopsis, rapeseed, and soybean can also be used by one skilled in the art to obtain other substantially homologous full-length cDNAs, if not already obtained, as well as the corresponding genes as fragments of the invention. These, in turn, may be used to overexpress the corresponding desaturases in plants. One skilled in the art can also isolate the coding sequence(s) from the fragment(s) of the invention by using and/or creating sites for restriction endonucleases, as described in Sambrook et al., (Molecular Cloning, A Laboratory Manual, 2nd ed. (1989), Cold Spring Harbor Laboratory Press).

One particularly useful application of the claimed inventions is to repair the agronomic performance of plant mutants containing ultra high levels of oleate in seed oil. In Arabidopsis reduction in linoleate in phosphatidylcholine due to a mutation in microsomal delta-12 desaturase affected low temperature survival [Miquel, M. et. al. (1993) Proc. Natl Acad. Sci. USA 90:6208-6212]. Furthermore, there is evidence that the poor agronomic performance of canola plants containing ultra high (>80%) levels of oleate in seed is due to mutations in the microsomal delta-12 desaturase genes that reduce the level of linoleate in phosphotidylcholine of roots and leaves. That is, the mutations are not seed-specific. Thus, the root and/or leaf-specific expression (that is, with no expression in the seeds) of microsomal delta-12 desaturase activity in mutants of oilseeds with ultra-high levels of oleate in seed oil

will result in agronomically-improved mutant plants with ultra high levels of oleate in seed oil.

#### Inhibition of Plant Target

##### Genes by Use of Antisense RNA

5        Antisense RNA has been used to inhibit plant target genes in a tissue-specific manner (see van der Krol et al., *Biotechniques* (1988) 6:958-976). Antisense inhibition has been shown using the entire cDNA sequence (Sheehy et al., *Proc. Natl. Acad. Sci. USA* (1988) 85:8805-8809) as well as a partial cDNA sequence (Cannon et al., *Plant Molec. Biol.* (1990) 15:39-47). There is also evidence that the 3' non-coding sequences (Ch'ng et al., *Proc. Natl. Acad. Sci. USA* (1989) 86:10006-10010) and fragments of 5' coding sequence, 15 containing as few as 41 base-pairs of a 1.87 kb cDNA (Cannon et al., *Plant Molec. Biol.* (1990) 15:39-47), can play important roles in antisense inhibition.

20        The use of antisense inhibition of the fatty acid desaturases may require isolation of the transcribed sequence for one or more target fatty acid desaturase genes that are expressed in the target tissue of the target plant. The genes that are most highly expressed are the best targets for antisense inhibition. These genes may be identified by determining their levels of 25 transcription by techniques, such as quantitative analysis of mRNA levels or nuclear run-off transcription, known to one skilled in the art.

30        The entire soybean microsomal delta-12 desaturase cDNA was cloned in the antisense orientation with respect to either soybean b-conglycinin, soybean KTi3, and bean phaseolin promoter and the chimeric gene transformed into soybean somatic embryos that were previously shown to serve as good model system for soybean zygotic embryos and are predictive of seed 35 composition (Table 11). Transformed somatic embryos

09697279-102600

showed inhibition of linoleate biosynthesis. Similarly, the entire Brassica napus microsomal delta-12 desaturase cDNA was cloned in the antisense orientation with respect to a rapeseed napin promoter and the chimeric gene transformed into B. napus. Seeds of transformed B. napus plants showed inhibition of linoleate biosynthesis. Thus, antisense inhibition of delta-12 desaturase in oil-producing species, including corn, Brassica napus, and soybean resulting in altered levels of polyunsaturated fatty acids may be achieved by expressing antisense RNA from the entire or partial cDNA encoding microsomal delta-12 desaturase.

#### Inhibition of Plant

##### Target Genes by Cosuppression

The phenomenon of cosuppression has also been used to inhibit plant target genes in a tissue-specific manner. Cosuppression of an endogenous gene using the entire cDNA sequence (Napoli et al., The Plant Cell (1990) 2:279-289; van der Krol et al., The Plant Cell (1990) 2:291-299) as well as a partial cDNA sequence (730 bp of a 1770 bp cDNA) (Smith et al., Mol. Gen. Genetics (1990) 224:477-481) are known.

The nucleic acid fragments of the instant invention encoding fatty acid desaturases, or parts thereof, with suitable regulatory sequences, can be used to reduce the level of fatty acid desaturases, thereby altering fatty acid composition, in transgenic plants which contain an endogenous gene substantially homologous to the introduced nucleic acid fragment. The experimental procedures necessary for this are similar to those described above for the overexpression of the fatty acid desaturase nucleic acid fragments except that one may also use a partial cDNA sequence. For example, cosuppression of delta-12 desaturase in Brassica napus and soybean resulting in altered levels of

polyunsaturated fatty acids may be achieved by expressing in the sense orientation the entire or partial seed delta-12 desaturase cDNA found in pCF2-165D and pSF2-165K, respectively. Endogenous genes can also  
 5 be inhibited by non-coding regions of an introduced copy of the gene [For example, Brusslan, J. A. et al. (1993) Plant Cell 5:667-677; Matzke, M. A. et al., Plant Molecular Biology 16:821-830]. We have shown that an Arabidopsis gene (SEQ ID NO:15) corresponding to the  
 10 cDNA (SEQ ID NO:1) can be isolated. One skilled in the art can readily isolate genomic DNA containing or flanking the genes and use the coding or non-coding regions in such transgene inhibition methods.

Analysis of the fatty acid composition of roots and  
 15 seeds of Arabidopsis mutants deficient in microsomal delta-12 desaturation shows that they have reduced levels of 18:2 as well as reduced levels of 16:0 (as much as 40% reduced level in mutant seeds as compared to wild type seeds) [Miquel and Browse (1990) in Plant  
 20 Lipid Biochemistry, Structure, and Utilization, pages 456-458, Ed. Quinn, P. J. and Harwood, J. L., Portland Press. Reduction in the level of 16:0 is also observed in ultra high oleate mutants of B. napus. Thus, one can expect that ultra high level of 18:1 in  
 25 transgenic plants due to antisense inhibition or co-supression using the claimed sequences will also reduce the level of 16:0.

One particularly useful application of the claimed invention is to combine the high oleate trait of the  
 30 transformed seeds with mutations for altered fatty acid compositions to obtain novel fatty acid compositions and/or improved agronomy. Fatty acid compositions, such as ultra-high oleate levels (EPO 323753 and see below), resulting from mutagenesis alone, such as by combining  
 35 mutations, may be affected in agronomy or seed

096979-10260

germination. Transgene and mutant genes may be combined either by making a genetic cross between the two lines or by transforming the chimeric genes of the inventions into the mutant lines as known to one skilled in the art.

As examples, the high oleic acid trait of the invention can be combined with high stearate mutations, such as soybean A6 [Hammond, E. G. and Fehr, W. R. (1983)] or low linolenic acid mutations and soybean mutants A5, A23, A16 and C1640 [Fehr, W. R. et al. (1992) in Crop Science 32:903-906]. Oils produced from such combinations will provide improved feedstocks for production of margarines, shortenings, spray coating and frying oils and will eliminate or reduce the need for hydrogenation. Furthermore, these oils would provide a health benefit for consumers, for example by reducing or eliminating trans fatty acids which have recently been found to be associated with high risk to cardiovascular diseases.

It can also be combined with other high oleic acid mutants that will result in even higher oleic acid content. Examples of high oleate mutants include soybean lines A5 and N782245 [Martin, B. A. and Rinne, R. W. (1985) Crop Science 25:1055-1058] and rapeseed lines containing about 69% to 77% oleic acid. A mutant (IMC 129) was derived by chemical mutagenesis of Westar and is presently in commercial production in the U.S. This variety has yields comparable to Westar, a leading Canadian variety. Its high oleate trait can be attributed to a mutation found in the coding region (Glu<sub>106</sub> to Lys<sub>106</sub>) one of the two delta-12 desaturase structural genes found in Westar. Re-mutagenesis of this line resulted in a second mutation in the other delta-12 desaturase structural gene and even higher seed oleic acid content in the double mutant. However the

double mutant has significantly higher levels of oleic acid in its leaves and roots than Westar, i.e., the mutant phenotype is not seed specific. The double mutant has extremely poor agronomy characteristics, which appear to be related to the high oleate in its leaves and/or roots. Thus, one can combine the seed-specific high oleic acid content resulting from seed-specific antisense inhibition or cosuppression of the microsomal delta-12 desaturase in transgenic seeds with that in the single mutant with 69-77% oleic acid to obtain levels of seed oleic acid that are as high or higher than that in the double mutant without affecting agronomy. The increased levels of oleic acid in the double homozygotes are associated with reduced levels of saturates. The reduction in saturates is also observed with seed-specific inhibition of delta-12 desaturase in soybean seeds.

#### Selection of Hosts, Promoters and Enhancers

A preferred class of heterologous hosts for the expression of the nucleic acid fragments of the invention are eukaryotic hosts, particularly the cells of higher plants. Particularly preferred among the higher plants are the oil-producing species, such as soybean (Glycine max), rapeseed (including Brassica napus, B. campestris), sunflower (Helianthus annus), cotton (Gossypium hirsutum), corn (Zea mays), cocoa (Theobroma cacao), safflower (Carthamus tinctorius), oil palm (Elaeis guineensis), coconut palm (Cocos nucifera), flax (Linum usitatissimum), and peanut (Arachis hypogaea).

Expression in plants will use regulatory sequences functional in such plants. The expression of foreign genes in plants is well-established (De Blaere et al., Meth. Enzymol. (1987) 153:277-291). The source of the promoter chosen to drive the expression of the fragments

009697379 "102600

of the invention is not critical provided it has sufficient transcriptional activity to accomplish the invention by increasing or decreasing, respectively, the level of translatable mRNA for the fatty acid

5 desaturases in the desired host tissue. Preferred promoters include (a) strong constitutive plant promoters, such as those directing the 19S and 35S transcripts in cauliflower mosaic virus (Odell et al., Nature (1985) 313:810-812; Hull et al., Virology (1987)

10 86:482-493), (b) tissue- or developmentally-specific promoters, and (c) other transcriptional promoter systems engineered in plants, such as those using bacteriophage T7 RNA polymerase-promoter sequences to express foreign genes. Examples of tissue-specific

15 promoters are the light-inducible promoter of the small subunit of ribulose 1,5-bis-phosphate carboxylase (if expression is desired in photosynthetic tissues), the maize zein protein promoter (Matzke et al., EMBO J. (1984) 3:1525-1532), and the chlorophyll a/b binding

20 protein promoter (Lampa et al., Nature (1986) 316:750-752).

Particularly preferred promoters are those that allow seed-specific expression. This may be especially useful since seeds are the primary source of vegetable

25 oils and also since seed-specific expression will avoid any potential deleterious effect in non-seed tissues. Examples of seed-specific promoters include, but are not limited to, the promoters of seed storage proteins, which can represent up to 90% of total seed protein in

30 many plants. The seed storage proteins are strictly regulated, being expressed almost exclusively in seeds in a highly tissue-specific and stage-specific manner (Higgins et al., Ann. Rev. Plant Physiol. (1984) 35:191-221; Goldberg et al., Cell (1989) 56:149-160).

Moreover, different seed storage proteins may be expressed at different stages of seed development.

Expression of seed-specific genes has been studied in great detail (See reviews by Goldberg et al., Cell (1989) 56:149-160 and Higgins et al., Ann. Rev. Plant Physiol. (1984) 35:191-221). There are currently numerous examples of seed-specific expression of seed storage protein genes in transgenic dicotyledonous plants. These include genes from dicotyledonous plants for bean b-phaseolin (Sengupta-Gopalan et al., Proc. Natl. Acad. Sci. USA (1985) 82:3320-3324; Hoffman et al., Plant Mol. Biol. (1988) 11:717-729), bean lectin (Voelker et al., EMBO J. (1987) 6:3571-3577), soybean lectin (Okamuro et al., Proc. Natl. Acad. Sci. USA (1986) 83:8240-8244), soybean Kunitz trypsin inhibitor (Perez-Grau et al., Plant Cell (1989) 1:095-1109), soybean b-conglycinin (Beachy et al., EMBO J. (1985) 4:3047-3053; pea vicilin (Higgins et al., Plant Mol. Biol. (1988) 11:683-695), pea convicilin (Newbigin et al., Planta (1990) 180:461-470), pea legumin (Shirsat et al., Mol. Gen. Genetics (1989) 215:326-331); rapeseed napin (Radke et al., Theor. Appl. Genet. (1988) 75:685-694) as well as genes from monocotyledonous plants such as for maize 15 kD zein (Hoffman et al., EMBO J. (1987) 6:3213-3221), maize 18 kD oleosin (Lee et al., Proc. Natl. Acad. Sci. USA (1991) 88:6181-6185), barley b-hordein (Marris et al., Plant Mol. Biol. (1988) 10:359-366) and wheat glutenin (Colot et al., EMBO J. (1987) 6:3559-3564). Moreover, promoters of seed-specific genes operably linked to heterologous coding sequences in chimeric gene constructs also maintain their temporal and spatial expression pattern in transgenic plants. Such examples include use of Arabidopsis thaliana 2S seed storage protein gene promoter to express enkephalin peptides in Arabidopsis



5

10

30

B. napus isocitrate lyase and malate synthase (Comai et al., Plant Cell (1989) 1:293-300), delta-9 desaturase from safflower (Thompson et al. Proc. Natl. Acad. Sci. USA (1991) 88:2578-2582) and castor (Shanklin et al., Proc. Natl. Acad. Sci. USA (1991) 88:2510-2514), acyl carrier protein (ACP) from Arabidopsis (Post-Beittenmiller et al., Nucl. Acids Res. (1989) 17:1777), B. napus (Safford et al., Eur. J. Biochem. (1988) 174:287-295), and B. campestris (Rose et al., Nucl. Acids Res. (1987) 15:7197), b-ketoacyl-ACP synthetase from barley (Siggaard-Andersen et al., Proc. Natl. Acad. Sci. USA (1991) 88:4114-4118), and oleosin from Zea mays (Lee et al., Proc. Natl. Acad. Sci. USA (1991) 88:6181-6185), soybean (Genbank Accession No: X60773) and B. napus (Lee et al., Plant Physiol. (1991) 96:1395-1397) will be of use. If the sequence of the corresponding genes is not disclosed or their promoter region is not identified, one skilled in the art can use the published sequence to isolate the corresponding gene and a fragment thereof containing the promoter. The partial protein sequences for the relatively-abundant enoyl-ACP reductase and acetyl-CoA carboxylase are also published (Slabas et al., Biochim. Biophys. Acta (1987) 877:271-280; Cottingham et al., Biochim. Biophys. Acta (1988) 954:201-207) and one skilled in the art can use these sequences to isolate the corresponding seed genes with their promoters. Similarly, the fragments of the present invention encoding fatty acid desaturases can be used to obtain promoter regions of the corresponding genes for use in expressing chimeric genes.

Attaining the proper level of expression of the nucleic acid fragments of the invention may require the use of different chimeric genes utilizing different promoters. Such chimeric genes can be transferred into

host plants either together in a single expression vector or sequentially using more than one vector.

It is envisioned that the introduction of enhancers or enhancer-like elements into the promoter regions of either the native or chimeric nucleic acid fragments of the invention will result in increased expression to accomplish the invention. This would include viral enhancers such as that found in the 35S promoter (Odell et al., Plant Mol. Biol. (1988) 10:263-272), enhancers from the opine genes (Fromm et al., Plant Cell (1989) 1:977-984), or enhancers from any other source that result in increased transcription when placed into a promoter operably linked to the nucleic acid fragment of the invention.

Of particular importance is the DNA sequence element isolated from the gene for the  $\alpha$ -subunit of b-conglycinin that can confer 40-fold seed-specific enhancement to a constitutive promoter (Chen et al., Dev. Genet. (1989) 10:112-122). One skilled in the art can readily isolate this element and insert it within the promoter region of any gene in order to obtain seed-specific enhanced expression with the promoter in transgenic plants. Insertion of such an element in any seed-specific gene that is expressed at different times than the b-conglycinin gene will result in expression in transgenic plants for a longer period during seed development.

The invention can also be accomplished by a variety of other methods to obtain the desired end. In one form, the invention is based on modifying plants to produce increased levels of fatty acid desaturases by virtue of introducing more than one copy of the foreign gene containing the nucleic acid fragments of the invention. In some cases, the desired level of polyunsaturated fatty acids may require introduction of

foreign genes for more than one kind of fatty acid desaturase.

Any 3' non-coding region capable of providing a polyadenylation signal and other regulatory sequences that may be required for the proper expression of the nucleic acid fragments of the invention can be used to accomplish the invention. This would include 3' ends of the native fatty acid desaturase(s), viral genes such as from the 35S or the 19S cauliflower mosaic virus transcripts, from the opine synthesis genes, ribulose 1,5-bisphosphate carboxylase, or chlorophyll a/b binding protein. There are numerous examples in the art that teach the usefulness of different 3' non-coding regions.

#### Transformation Methods

Various methods of transforming cells of higher plants according to the present invention are available to those skilled in the art (see EPO Pub. 0 295 959 A2 and 0 318 341 A1). Such methods include those based on transformation vectors utilizing the Ti and Ri plasmids of Agrobacterium spp. It is particularly preferred to use the binary type of these vectors. Ti-derived vectors transform a wide variety of higher plants, including monocotyledonous and dicotyledonous plants (Sukhapinda et al., Plant Mol. Biol. (1987) 8:209-216; Potrykus, Mol. Gen. Genet. (1985) 199:183). Other transformation methods are available to those skilled in the art, such as direct uptake of foreign DNA constructs (see EPO Pub. 0 295 959 A2), techniques of electroporation (Fromm et al., Nature (1986) (London) 319:791) or high-velocity ballistic bombardment with metal particles coated with the nucleic acid constructs (Kline et al., Nature (1987) (London) 327:70). Once transformed, the cells can be regenerated by those skilled in the art.

09697379-102600

Of particular relevance are the recently described methods to transform foreign genes into commercially important crops, such as rapeseed (De Block et al., Plant Physiol. (1989) 91:694-701), sunflower (Everett et al., Bio/Technology (1987) 5:1201), and soybean (Christou et al., Proc. Natl. Acad. Sci USA (1989) 86:7500-7504).

#### Application to Molecular Breeding

The 1.6 kb insert obtained from the plasmid pSF2-169K was used as a radiolabelled probe on a Southern blot containing genomic DNA from soybean (Glycine max (cultivar Bonus) and Glycine soja (PI81762)) digested with one of several restriction enzymes. Different patterns of hybridization (polymorphisms) were identified in digests performed with restriction enzymes Hind III and Eco RI. These polymorphisms were used to map two pSF2-169 loci relative to other loci on the soybean genome essentially as described by Helentjaris et al., (Theor. Appl. Genet. (1986) 72:761-769). One mapped to linkage group 11 between 4404.00 and 1503.00 loci (4.5 cM and 7.1 cM from 4404.00 and 1503.00, respectively) and the other to linkage group 19 between 4010.00 and 5302.00 loci (1.9 cM and 2.7 cM from 4010.00 and 5302.00, respectively) [Rafalski, A and Tingey, S. (1993) in Genetic Maps, Ed. O' Brien, S. J.]. The use of restriction fragment length polymorphism (RFLP) markers in plant breeding has been well-documented in the art (Tanksley et al., Bio/Technology (1989) 7:257-264). Thus, the nucleic acid fragments of the invention can be used as RFLP markers for traits linked to expression of fatty acid desaturases. These traits will include altered levels of unsaturated fatty acids. The nucleic acid fragment of the invention can also be used to isolate the fatty acid desaturase gene from variant

009201" 62E26350

(including mutant) plants with altered levels of unsaturated fatty acids. Sequencing of these genes will reveal nucleotide differences from the normal gene that cause the variation. Short oligonucleotides designed  
 5 around these differences may also be used in molecular breeding either as hybridization probes or in DNA-based diagnostics to follow the variation in fatty acids. Oligonucleotides based on differences that are linked to the variation may be used as molecular markers in  
 10 breeding these variant oil traits.

#### EXAMPLES

The present invention is further defined in the following Examples, in which all parts and percentages are by weight and degrees are Celsius, unless otherwise  
 15 stated. It should be understood that these Examples, while indicating preferred embodiments of the invention, are given by way of illustration only. From the above discussion and these Examples, one skilled in the art can ascertain the essential characteristics of this  
 20 invention, and without departing from the spirit and scope thereof, can make various changes and modifications of the invention to adapt it to various usages and conditions. All publications, including patents and non-patent literature, referred to in this  
 25 specification are expressly incorporated by reference herein.

#### EXAMPLE 1

##### ISOLATION OF GENOMIC DNA FLANKING THE T-DNA SITE OF INSERTION IN ARABIDOPSIS THALIANA MUTANT LINE 658

##### 30 Identification of an Arabidopsis thaliana T-DNA Mutant with High Oleic Acid Content

A population of Arabidopsis thaliana (geographic race Wassilewskija) transformants containing the modified T-DNA of Agrobacterium tumefaciens was  
 35 generated by seed transformation as described by

00920T" 62E25950

Feldmann et al., (Mol. Gen. Genetics (1987) 208:1-9).  
 In this population the transformants contain DNA  
 sequences encoding the pBR322 bacterial vector, nopaline  
 synthase, neomycin phosphotransferase (NPTII, confers  
 5 kanamycin resistance), and b-lactamase (confers  
 ampicillin resistance) within the T-DNA border  
 sequences. The integration of the T-DNA into different  
 areas of the chromosomes of individual transformants may  
 cause a disruption of plant gene function at or near the  
 10 site of insertion, and phenotypes associated with this  
 loss of gene function can be analyzed by screening the  
 population for the phenotype.

T3 seed was generated from the wild type seed  
 treated with Agrobacterium tumefaciens by two rounds of  
 15 self-fertilization as described by Feldmann et al.,  
 (Science (1989) 243:1351-1354). These progeny were  
 segregating for the T-DNA insertion, and thus for any  
 mutation resulting from the insertion. Approximately  
 10-12 leaves of each of 1700 lines were combined and the  
 20 fatty acid content of each of the 1700 pooled samples  
 was determined by gas chromatography of the fatty acyl  
 methyl esters essentially as described by Browse et al.,  
 (Anal. Biochem. (1986) 152:141-145) except that 2.5%  
 H<sub>2</sub>SO<sub>4</sub> in methanol was used as the methylation reagent.  
 25 A line designated "658" produced a sample that gave an  
 altered fatty acid profile compared to those of lines  
 657 and 659 sampled at the same time (Table 8).

TABLE 8

Fatty Acid <u>Methyl Ester</u>	657 Leaf <u>Pool</u>	659 Leaf <u>Pool</u>	658 Leaf <u>Pool</u>
16:0	14.4	14.1	13.6
16:1	4.4	4.6	4.5
16:2	2.9	2.2	2.7
16:3	13.9	13.3	13.9

18:0	1.0	1.1	0.9
18:1	2.6	2.5	4.9
18:2	14.0	13.6	12.8
18:3	42.9	46.1	44.4

Analysis of the fatty acid composition of 12 individual T3 seeds of line 658 indicated that the 658 pool was composed of seeds segregating in three classes: "high", "mid-range" and "low" classes with approximately, 37% (12 seeds), 21% (7 seeds), and 14% (3 seeds) oleic acid, respectively (Table 9).

TABLE 9

	<u>"High"</u> <u>Class</u>	<u>"Mid-range"</u> <u>Class</u>	<u>"Low"</u> <u>Class</u>
16:0	8.9	8.7	9.3
16:1c	2.0	1.6	2.6
18:0	4.5	4.3	4.4
18:1	37.0	20.7	14.4
18:2	8.0	24.9	27.7
18:3	10.6	14.3	13.6
20:1	25.5	21.6	20.4

Thus, the high oleic acid mutant phenotype segregates in an approximately Mendelian ratio. To determine the number of independently segregating T-DNA inserts in line 658, 200 T3 seeds were tested for their ability to germinate and grow in the presence of kanamycin [Feldman et al. (1989) Science 243:1351-1354]. In this experiment, only 4 kanamycin-sensitive individual plants were identified. The segregation ratio (approximately 50:1) indicated that line 658 harbored three T-DNA inserts. In this and two other experiments a total of 56 kanamycin-sensitive plants were identified; 53 of these were analyzed for fatty



acid composition and at least seven of these displayed oleic acid levels that were higher than would be expected for wild type seedlings grown under these conditions.

- 5           In order to more rigorously test whether the  
mutation resulting in high oleic acid is the result of  
T-DNA insertion, Applicants identified a derivative line  
that was segregating for the mutant fatty acid phenotype  
as well as a single kanamycin resistance locus. For  
10 this, approximately 100 T3 plants were individually  
grown to maturity and seeds collected. One sample of  
seed from each T3 plant was tested for the ability to  
germinate and grow in the presence of kanamycin. In  
addition, the fatty acid compositions of ten additional  
15 individual seeds from each line were determined. A T3  
plant, designated 658-75, was identified whose progeny  
seeds segregated 28 kanamycin-sensitive to 60 kanamycin-  
resistant and 7 with either low or intermediate oleic  
acid to 2 high oleic acid.
- 20           A total of approximately 400 T4 progeny seeds of  
the derivative line 658-75 were grown and the leaf fatty  
acid composition analyzed. A total of 91 plants were  
identified as being homozygous for the high oleic acid  
trait (18:2/18:1 less than 0.5). The remaining plants  
25 (18:2/18:1 more than 1.2) could not be definitively  
assigned to wild type and heterozygous classes on the  
basis of leaf fatty acid composition and thus could not  
be used to test linkage between the kanamycin marker and  
the fatty acid mutation. Eighty three of the 91  
30 apparently homozygous high oleic acid mutant were tested  
for the presence of nopaline, another T-DNA marker, in  
leaf extracts (Errampalli et al. The Plant Cell  
(1991)3:149-157 and all 83 plants were positive for the  
presence of nopaline. This tight linkage of the mutant  
35 fatty acid phenotype and a T-DNA marker provides

to, the microsomal delta-12 and delta-15 desaturases that act on phosphatidylcholine lipid substrates; the chloroplastic or plastid delta-12 and delta-15 desaturases that act on phosphatidyl glycerol and galactolipids; and other desaturases that act on such fatty acid substrates such as phospholipids, galactolipids, and sulfolipids. "Microsomal desaturase" refers to the cytoplasmic location of the enzyme, while "chloroplast desaturase" and "plastid desaturase" refer to the plastid location of the enzyme. These fatty acid desaturases may be found in a variety of organisms including, but not limited to, higher plants, diatoms, and various eukaryotic and prokaryotic microorganisms such as fungi and photosynthetic bacteria and algae.

The term "homologous fatty acid desaturases" refers to fatty acid desaturases that catalyze the same desaturation on the same lipid substrate. Thus, microsomal delta-15 desaturases, even from different plant species, are homologous fatty acid desaturases.

The term "heterologous fatty acid desaturases" refers to fatty acid desaturases that catalyze desaturations at different positions and/or on different lipid substrates. Thus, for example, microsomal delta-12 and delta-15 desaturases, which act on phosphatidylcholine lipids, are heterologous fatty acid desaturases, even when from the same plant. Similarly, microsomal delta-15 desaturase, which acts on phosphatidylcholine lipids, and chloroplast delta-15 desaturase, which acts on galactolipids, are heterologous fatty acid desaturases, even when from the same plant. It should be noted that these fatty acid desaturases have never been isolated and characterized as proteins.

Accordingly, the terms such as "delta-12 desaturase" and "delta-15 desaturase" are used as a convenience to describe the proteins encoded by nucleic acid fragments

009201" 62E26960

that have been isolated based on the phenotypic effects caused by their disruption. They do not imply any catalytic mechanism. For example, delta-12 desaturase refers to the enzyme that catalyzes the formation of a double bond between carbons 12 and 13 of an 18 carbon fatty acid irrespective of whether it "counts" the carbons from the methyl, carboxyl end, or the first double bond. The term "fatty acid desaturase-related enzyme" refers to enzymes whose catalytic product may not be a carbon-carbon double bond but whose mechanism of action is similar to that of a fatty acid desaturase (that is, catalysis of the displacement of a carbon-hydrogen bond of a fatty acid chain to form a fatty-hydroxyacyl intermediate or end-product). Examples include delta-12 hydroxylase which means a delta-12 fatty acid hydroxylase or the oleate hydroxylase responsible for the synthesis of ricinoleic acid from oleic acid.

The term "nucleic acid" refers to a large molecule which can be single-stranded or double-stranded, composed of monomers (nucleotides) containing a sugar, a phosphate and either a purine or pyrimidine. A "nucleic acid fragment" is a fraction of a given nucleic acid molecule. In higher plants, deoxyribonucleic acid (DNA) is the genetic material while ribonucleic acid (RNA) is involved in the transfer of the information in DNA into proteins. A "genome" is the entire body of genetic material contained in each cell of an organism. The term "nucleotide sequence" refers to the sequence of DNA or RNA polymers, which can be single- or double-stranded, optionally containing synthetic, non-natural or altered nucleotide bases capable of incorporation into DNA or RNA polymers. The term "oligomer" refers to short nucleotide sequences, usually up to 100 bases long. As used herein, the term "homologous to" refers

to the relatedness between the nucleotide sequence of two nucleic acid molecules or between the amino acid sequences of two protein molecules. Estimates of such homology are provided by either DNA-DNA or DNA-RNA hybridization under conditions of stringency as is well understood by those skilled in the art (Hames and Higgins, Eds. (1985) *Nucleic Acid Hybridisation*, IRL Press, Oxford, U.K.); or by the comparison of sequence similarity between two nucleic acids or proteins, such as by the method of Needleman et al. (*J. Mol. Biol.* (1970) 48:443-453). As used herein, "substantially homologous" refers to nucleotide sequences that have more than 90% overall identity at the nucleotide level with the coding region of the claimed sequence, such as genes and pseudo-genes corresponding to the coding regions. The nucleic acid fragments described herein include molecules which comprise possible variations, both man-made and natural, such as but not limited to (a) those that involve base changes that do not cause a change in an encoded amino acid, or (b) which involve base changes that alter an amino acid but do not affect the functional properties of the protein encoded by the DNA sequence, (c) those derived from deletions, rearrangements, amplifications, random or controlled mutagenesis of the nucleic acid fragment, and (d) even occasional nucleotide sequencing errors.

"Gene" refers to a nucleic acid fragment that expresses a specific protein, including regulatory sequences preceding (5' non-coding) and following (3' non-coding) the coding region. "Fatty acid desaturase gene" refers to a nucleic acid fragment that expresses a protein with fatty acid desaturase activity. "Native" gene refers to an isolated gene with its own regulatory sequences as found in nature. "Chimeric gene" refers to a gene that comprises heterogeneous regulatory and

009201" 52E25950

coding sequences not found in nature. "Endogenous" gene refers to the native gene normally found in its natural location in the genome and is not isolated. A "foreign" gene refers to a gene not normally found in the host organism but that is introduced by gene transfer.

"Pseudo-gene" refers to a genomic nucleotide sequence that does not encode a functional enzyme.

"Coding sequence" refers to a DNA sequence that codes for a specific protein and excludes the non-coding sequences. It may constitute an "uninterrupted coding sequence", i.e., lacking an intron or it may include one or more introns bounded by appropriate splice junctions. An "intron" is a nucleotide sequence that is transcribed in the primary transcript but that is removed through cleavage and re-ligation of the RNA within the cell to create the mature mRNA that can be translated into a protein.

"Initiation codon" and "termination codon" refer to a unit of three adjacent nucleotides in a coding sequence that specifies initiation and chain termination, respectively, of protein synthesis (mRNA translation). "Open reading frame" refers to the coding sequence uninterrupted by introns between initiation and termination codons that encodes an amino acid sequence.

"RNA transcript" refers to the product resulting from RNA polymerase-catalyzed transcription of a DNA sequence. When the RNA transcript is a perfect complementary copy of the DNA sequence, it is referred to as the primary transcript or it may be a RNA sequence derived from posttranscriptional processing of the primary transcript and is referred to as the mature RNA. "Messenger RNA (mRNA)" refers to the RNA that is without introns and that can be translated into protein by the cell. "cDNA" refers to a double-stranded DNA that is complementary to and derived from mRNA. "Sense" RNA

009201 6226950

refers to RNA transcript that includes the mRNA.

"Antisense RNA" refers to a RNA transcript that is complementary to all or part of a target primary transcript or mRNA and that blocks the expression of a target gene by interfering with the processing, transport and/or translation of its primary transcript or mRNA. The complementarity of an antisense RNA may be with any part of the specific gene transcript, i.e., at the 5' non-coding sequence, 3' non-coding sequence, introns, or the coding sequence. In addition, as used herein, antisense RNA may contain regions of ribozyme sequences that increase the efficacy of antisense RNA to block gene expression. "Ribozyme" refers to a catalytic RNA and includes sequence-specific endoribonucleases.

As used herein, "suitable regulatory sequences" refer to nucleotide sequences in native or chimeric genes that are located upstream (5'), within, and/or downstream (3') to the nucleic acid fragments of the invention, which control the expression of the nucleic acid fragments of the invention. The term "expression", as used herein, refers to the transcription and stable accumulation of the sense (mRNA) or the antisense RNA derived from the nucleic acid fragment(s) of the invention that, in conjunction with the protein apparatus of the cell, results in altered levels of the fatty acid desaturase(s). Expression or overexpression of the gene involves transcription of the gene and translation of the mRNA into precursor or mature fatty acid desaturase proteins. "Antisense inhibition" refers to the production of antisense RNA transcripts capable of preventing the expression of the target protein. "Overexpression" refers to the production of a gene product in transgenic organisms that exceeds levels of production in normal or non-transformed organisms. "Cosuppression" refers to the expression of a foreign

009207 5252690

gene which has substantial homology to an endogenous gene resulting in the suppression of expression of both the foreign and the endogenous gene. "Altered levels" refers to the production of gene product(s) in

5 transgenic organisms in amounts or proportions that differ from that of normal or non-transformed organisms.

"Promoter" refers to a DNA sequence in a gene, usually upstream (5') to its coding sequence, which controls the expression of the coding sequence by  
 10 providing the recognition for RNA polymerase and other factors required for proper transcription. In artificial DNA constructs promoters can also be used to transcribe antisense RNA. Promoters may also contain DNA sequences that are involved in the binding of  
 15 protein factors which control the effectiveness of transcription initiation in response to physiological or developmental conditions. It may also contain enhancer elements. An "enhancer" is a DNA sequence which can stimulate promoter activity. It may be an innate  
 20 element of the promoter or a heterologous element inserted to enhance the level and/or tissue-specificity of a promoter. "Constitutive promoters" refers to those that direct gene expression in all tissues and at all times. "Tissue-specific" or "development-specific"  
 25 promoters as referred to herein are those that direct gene expression almost exclusively in specific tissues, such as leaves or seeds, or at specific development stages in a tissue, such as in early or late embryogenesis, respectively.

30 The "3' non-coding sequences" refers to the DNA sequence portion of a gene that contains a polyadenylation signal and any other regulatory signal capable of affecting mRNA processing or gene expression. The polyadenylation signal is usually characterized by

009201 6262650

affecting the addition of polyadenylic acid tracts to the 3' end of the mRNA precursor.

"Transformation" herein refers to the transfer of a foreign gene into the genome of a host organism and its genetically stable inheritance. "Restriction fragment length polymorphism" (RFLP) refers to different sized restriction fragment lengths due to altered nucleotide sequences in or around variant forms of genes.

"Molecular breeding" refers to the use of DNA-based diagnostics, such as RFLP, RAPDs, and PCR in breeding. "Fertile" refers to plants that are able to propagate sexually.

"Plants" refer to photosynthetic organisms, both eukaryotic and prokaryotic, whereas the term "Higher plants" refers to eukaryotic plants. "Oil-producing species" herein refers to plant species which produce and store triacylglycerol in specific organs, primarily in seeds. Such species include soybean (Glycine max), rapeseed and canola (including Brassica napus, B. campestris), sunflower (Helianthus annus), cotton (Gossypium hirsutum), corn (Zea mays), cocoa (Theobroma cacao), safflower (Carthamus tinctorius), oil palm (Elaeis guineensis), coconut palm (Cocos nucifera), flax (Linum usitatissimum), castor (Ricinus communis) and peanut (Arachis hypogaea). The group also includes non-agronomic species which are useful in developing appropriate expression vectors such as tobacco, rapid cycling Brassica species, and Arabidopsis thaliana, and wild species which may be a source of unique fatty acids.

"Sequence-dependent protocols" refer to techniques that rely on a nucleotide sequence for their utility. Examples of sequence-dependent protocols include, but are not limited to, the methods of nucleic acid and oligomer hybridization and methods of DNA and RNA

009207 5225950



amplification such as are exemplified in various uses of the polymerase chain reaction (PCR).

Various solutions used in the experimental manipulations are referred to by their common names such as "SSC", "SSPE", "Denhardt's solution", etc. The composition of these solutions may be found by reference to Appendix B of Sambrook, et al. (Molecular Cloning, A Laboratory Manual, 2nd ed. (1989), Cold Spring Harbor Laboratory Press).

10        T-DNA Mutagenesis and Identification of an  
              Arabidopsis Mutant Defective in  
              Microsomal Delta-12 Desaturation

15        In T-DNA mutagenesis (Feldmann, et al., Science (1989) 243:1351-1354), the integration of T-DNA in the genome can interrupt normal expression of the gene at or near the site of the integration. If the resultant mutant phenotype can be detected and shown genetically to be tightly linked to the T-DNA insertion, then the "tagged" mutant locus and its wild type counterpart can be readily isolated by molecular cloning by one skilled in the art.

20        Arabidopsis thaliana seeds were transformed by Agrobacterium tumefaciens C58C1rif strain harboring the avirulent Ti-plasmid pGV3850::pAK1003 that has the T-DNA region between the left and right T-DNA borders replaced by the origin of replication region and ampicillin resistance gene of plasmid pBR322, a bacterial kanamycin resistance gene, and a plant kanamycin resistance gene (Feldmann, et al., Mol. Gen. Genetics (1987) 208:1-9).

25        Plants from the treated seeds were self-fertilized and the resultant progeny seeds, germinated in the presence of kanamycin, were self-fertilized to give rise to a population, designated T3, that was segregating for T-DNA insertions. T3 seeds from approximately 1700 T2

30        plants were germinated and grown under controlled

00920T"62E26950

environment. One leaf from each of ten T3 plants of each line were pooled and analyzed for fatty acid composition. One line, designated 658, showed an increased level of oleic acid (18:1). Analysis of twelve individual T3 seeds of line 658 identified two seeds that contained greater than 36% oleic acid while the remaining seeds contained 12-22% oleic acid. The mutant phenotype of increased level of oleic acid in leaf and seed tissues of line 658 and its segregation in individual T3 seeds suggested that line 658 harbors a mutation that affects desaturation of oleic acid to linoleic acid in both leaf and seed tissues. When approximately 200 T3 seeds of line 658 were tested for their ability to germinate in the presence of kanamycin, four kanamycin-sensitive seeds were identified, suggesting multiple, possibly three, T-DNA inserts in the original T2 line. When progeny seeds of 100 individual T3 plants were analyzed for fatty acid composition and their ability to germinate on kanamycin, one plant, designated 658-75, was identified whose progeny segregated 7 wild type:2 mutant for the increased oleic acid and 28 sensitive:60 resistant for kanamycin resistance. Approximately 400 T4 progeny seeds of derivative line 658-75 were grown and their leaves analyzed for fatty acid composition. Ninety one of these seedlings were identified as homozygous for the mutant (high oleic acid) phenotype. Eighty-three of these homozygous plants were tested for the presence of nopaline, another marker for T-DNA, and all of them were nopaline positive. On the basis of these genetic studies it was concluded that the mutation in microsomal delta-12 desaturation was linked to the T-DNA.

009201" 5252550

Isolation of Arabidopsis 658-75 Genomic DNA  
Containing the Disrupted Gene Controlling  
Microsomal Delta-12 Desaturation

In order to isolate the gene controlling microsomal  
5 delta-12 desaturation from wild-type Arabidopsis, a  
T-DNA-plant DNA "junction" fragment containing a T-DNA  
border integrated into the host plant DNA was isolated  
from the homozygous mutant plants of the 658-75 line of  
Arabidopsis. For this, genomic DNA from the mutant  
10 plant was isolated and completely digested by either Bam  
HI or Sal I restriction enzymes. In each case, one of  
the resultant fragments was expected to contain the  
origin of replication and ampicillin-resistance gene of  
pBR322 as well as the left T-DNA-plant DNA junction  
15 fragment. Such fragments were rescued as plasmids by  
ligating the digested genomic DNA fragments at a dilute  
concentration to facilitate self-ligation and then using  
the ligated fragments to transform E. coli cells. While  
no ampicillin-resistant colony was obtained from the  
20 plasmid rescue of Sal I-digested plant genomic DNA, a  
single ampicillin-resistant colony was obtained from the  
plasmid rescue of Bam HI-digested plant genomic DNA.  
The plasmid obtained from this transformant was  
designated p658-1. Restriction analysis of plasmid  
25 p658-1 with Bam HI, Sal I and Eco RI restriction enzymes  
strongly suggested that it contained the expected  
14.2 kb portion of the T-DNA (containing pBR322  
sequences) and a putative plant DNA/left T-DNA border  
fragment in a 1.6 kb Eco RI-Bam HI fragment. The 1.6 kb  
30 Eco RI-Bam HI fragment was subcloned into pBluescript SK  
[Stratagene] by standard cloning procedures described in  
Sambrook et al., (Molecular Cloning, A Laboratory  
Manual, 2nd ed. (1989), Cold Spring Harbor Laboratory  
Press) and the resultant plasmid, designated pS1658.

009207" 62626950

Isolation of Microsomal Delta-12 Desaturase cDNA  
and Gene from Wild type Arabidopsis

The 1.6 kb Eco RI-Bam HI fragment, which contained the putative plant DNA flanking T-DNA, in plasmid p658-1 was isolated and used as a radiolabeled hybridization probe to screen a cDNA library made to polyA<sup>+</sup> mRNA from the above-ground parts of Arabidopsis thaliana plants, which varied in size from those that had just opened their primary leaves to plants which had bolted and were flowering [Elledge et al. (1991) Proc. Natl. Acad Sci. USA 88:1731-1735]. The cDNA inserts in the library were made into an Xho I site flanked by Eco RI sites in lambda Yes vector [Elledge et al. (1991) Proc. Natl. Acad Sci. USA 88:1731-1735]. Of the several positively-hybridizing plaques, four were subjected to plaque purification. Plasmids were excised from the purified phages by site-specific recombination using the cre-lox recombination system in E. coli strain BNN132 [Elledge et al. (1991) Proc. Natl. Acad Sci. USA 88:1731-1735]. The four excised plasmids were digested by Eco RI restriction enzyme and shown to contain cDNA inserts ranging in size between 1 kb and 1.5 kb. Partial nucleotide sequence determination and restriction enzyme mapping of all four cDNAs revealed their common identity.

The partial nucleotide sequences of two cDNAs, designated pSF2b and p92103, containing inserts of ca. 1.2 kb and ca. 1.4 kb, respectively, were determined. The composite sequence derived from these plasmids is shown as SEQ ID NO:1 and is expected to be contained completely in plasmid p92103. SEQ ID NO:1 shows the 5' to 3' nucleotide sequence of 1372 base pairs of the Arabidopsis cDNA which encodes microsomal delta-12 fatty acid desaturase. Nucleotides 93-95 are the putative initiation codon of the open reading frame (nucleotides

009201 522555

93-1244), (identified by comparison of other plant delta-12 desaturases in this application). Nucleotides 1242-1244 are the termination codon. Nucleotides 1 to 92 and 1245-1372 are the 5' and 3' untranslated  
 5 nucleotides, respectively. The 383 amino acid protein sequence in SEQ ID NO:2 is that deduced from the open reading frame and has an estimated molecular weight of 44 kD.

The gene corresponding to SEQ ID NO:1 was isolated  
 10 by screening an Arabidopsis genomic DNA library using radiolabeled pSF2b cDNA insert, purifying the positively-hybridizing plaque, and subcloning a 6 kB Hind III insert fragment from the phage DNA in pBluescript vector. The sequence of 2973 nucleotides of  
 15 the gene is shown in SEQ ID NO:15. Comparison of the sequences of the gene (SEQ ID NO:15) and the cDNA (SEQ ID NO:1) revealed the presence of a single intron of 1134 bp at a position between nucleotide positions 88 and 89 of the cDNA, which is 4 nucleotides 5' to the  
 20 initiation codon.

The 1.6 kB Eco RI-Bam HI genomic border fragment insert in pS1658 was also partially sequenced from the Bam HI and Eco RI ends. Comparison of the nucleotide sequences of the gene (SEQ ID NO:15), the cDNA (SEQ ID  
 25 NO:1), the border fragment, and the published sequence of the left end of T-DNA (Yadav et al., (1982) Proc. Natl. Acad. Sci. 79:6322-6326) revealed that a) the sequence of the first 451 nucleotides of the border fragment from the Bam HI end is collinear with that of  
 30 nucleotides 539 (Bam HI site) to 89 of the cDNA, b) from the Eco RI end, the border fragment is collinear from nucleotides 1 to 61 with that of the left end of T-DNA (except for a deletion of 9 contiguous nucleotides at position 42 in the border fragment), and is collinear  
 35 from nucleotides 57 to 104 with that of nucleotides

009201" 52E26950

41-88 of the cDNA, and c) the sequence divergences between the border fragment and the cDNA are due to the presence of the intron in the border fragment. These results show that the T-DNA disrupted the microsomal delta-12 desaturase gene in the transcribed region between the promoter and the coding region and 5' to the intron in the untranslated sequence.

A phage DNA containing Arabidopsis microsomal delta-12 desaturase gene was used as a RFLP marker on a Southern blot containing genomic DNA from several progeny of Arabidopsis thaliana (ecotype Wassileskija and marker line W100 ecotype Landesberg background) digested with Hind III. This mapped the microsomal delta-12 desaturase gene 13.6 cM proximal to locus c3838, 9.2 cM distal to locus 1At228, and 4.9 cM proximal to Fad D locus on chromosome 3 [Koorneef, M. et al., (1993) in Genetic Maps, Ed. O'Brien, S. J.; Yadav et al. (1993) Plant Physiology 103:467-476]. This position corresponds closely to previously suggested locus for microsomal delta-12 desaturation (*Fad 2*) [Hugly, S. et al., (1991) Heredity 82:4321].

The open reading frames in SEQ ID NO:1 and in sequences encoding Arabidopsis microsomal delta-15 desaturase [WO 9311245], Arabidopsis plastid delta-15 desaturase [WO 9311245], and cyanobacterial desaturase, *des A*, [Wada, et al., Nature (1990) 347:200-203; Genbank ID:CSDESA; GenBank Accession No:X53508] as well as their deduced amino acid sequences were compared by the method of Needleman et al. [J. Mol. Biol. (1970) 48:443-453] using gap weight and gap length weight values of 5.0 and 0.3, respectively, for the nucleotide sequences and 3.0 and 0.1, respectively, for protein sequences. The overall identities are summarized in Table 2.

TABLE 2

Percent Identity Between Different Fatty Acid  
Desaturases at the Nucleotide and Amino Acid Levels

		<u>a3</u>	<u>ad</u>	<u>des A</u>
a2	nucleotide	48(8 gaps)	46(6 gaps)	43(10 gaps)
	amino acid	39(9 gaps)	34(8 gaps)	24(10 gaps)
a3	nucleotide	-	65(1 gap)	43(9 gaps)
	amino acid	-	65(2 gaps)	26(11 gaps)
ad	nucleotide	-	-	43(9 gaps)
	amino acid	-	-	26(11 gaps)

a2, a3, ad, and des A refer, respectively, to SEQ ID NO:1/2, Arabidopsis microsomal delta-15 desaturase, Arabidopsis plastid delta-15 desaturase, and cyanobacterial desaturase, des A. The percent identities in each comparison are shown at both the nucleotide and amino acid levels; the number of gaps imposed by the comparisons are shown in brackets following the percent identities. As expected on the basis of unsuccessful attempts in using delta-15 fatty acid nucleotide sequences as hybridization probes to isolate nucleotide sequences encoding microsomal delta-12 fatty acid desaturase, the overall homology at the nucleotide level between microsomal delta-12 fatty acid desaturase (SEQ ID NO:1) and the nucleotide sequences encoding the other three desaturases is poor (ranging between 43% and 48%). At the amino acid level too, the microsomal delta-12 fatty acid desaturase (SEQ ID NO:2) is poorly related to cyanobacterial des A (less than 24% identity) and the plant delta-15 desaturases (less than 39% identity).

While the overall relatedness between the deduced amino acid sequence of the said invention and the published fatty acid desaturases is limited, more significant identities are observed in shorter stretches

009201" 6/26/95

of amino acid sequences in the above comparisons. These results confirmed that the T-DNA in line 658-75 had interrupted the normal expression of a fatty acid desaturase gene. Based on the fatty acid phenotype of homozygous mutant line 658-75, Applicants concluded that SEQ ID NO:1 encoded the delta-12 desaturase. Further, Applicants concluded that it was the microsomal delta-12 desaturase, and not the chloroplastic delta-12 desaturase, since: a) the mutant phenotype was expressed strongly in the seed but expressed poorly, if at all, in the leaf of line 658-75, and b) the delta-12 desaturase polypeptide, by comparison to the microsomal and plastid delta-15 desaturase polypeptides [WO 9311245], did not have an N-terminal extension of a transit peptide expected for a nuclear-encoded plastid desaturase.

Plasmid p92103 was deposited on October 16, 1992 with the American Type Culture Collection of Rockville, Maryland under the provisions of the Budapest Treaty and bears accession number ATCC 69095.

Expression Of Microsomal Delta-12 Fatty Acid Desaturase In Arabidopsis Fad2-1 Mutant To Complement Its Mutation In Delta-12 Fatty Acid Desaturation

To confirm the identity of SEQ ID NO:1 (Arabidopsis microsomal delta-12 fatty acid desaturase cDNA) a chimeric gene comprising of SEQ ID NO:1 was transformed into an Arabidopsis mutant affected in microsomal delta-12 fatty acid desaturation. For this, the ca. 1.4 kb Eco RI fragment containing the cDNA (SEQ ID NO:1) was isolated from plasmid p92103 and sub-cloned in pGA748 vector [An et. al. (1988) Binary Vectors. In: Plant Molecular Biology Manual. Eds Gelvin, S. B. et al. Kluwer Academic Press], which was previously linearized with Eco RI restriction enzyme. In one of the resultant binary plasmid, designated pGA-Fad2, the cDNA was placed

009201"62E26950



in the sense orientation behind the CaMV 35S promotor of the vector to provide constitutive expression.

Binary vector pGA-Fad2 was transformed by the freeze/thaw method [Holsters et al. (1978) Mol. Gen. Genet. 163:181-187] into Agrobacterium tumefaciens strain R1000, carrying the Ri plasmid pRiA4b from Agrobacterium rhizogenes [Moore et al., (1979) Plasmid 2:617-626] to result in transformants R1000/pGA-Fad2.

Agrobacterium strains R1000 and R1000/pGA-Fad2 were used to transform Arabidopsis mutant fad2-1 [Miquel, M. & Browse, J. (1992) Journal of Biological Chemistry 267:1502-1509] and strain R1000 was used to transform wild type Arabidopsis. Young bolts of plants were sterilized and cut so that a single node was present in each explant. Explants were inoculated by Agrobacteria and incubated at 25°C in the dark on drug-free MS minimal organics medium with 30 g/L sucrose (Gibco). After four days, the explants were transferred to fresh MS medium containing 500 mg/L cefotaxime and 250 mg/ml carbenicillin for the counterselection of Agrobacterium. After 5 days, hairy roots derived from R1000/pGA-Fad2 transformation were excised and transferred to the same medium containing 50 mg/ml kanamycin. Fatty acid methyl esters were prepared from 5-10 mm of the roots essentially as described by Browse et al., (Anal. Biochem. (1986) 152:141-145) except that 2.5% H<sub>2</sub>SO<sub>4</sub> in methanol was used as the methylation reagent and samples were heated for 1.5 h at 80°C to effect the methanolysis of the seed triglycerides. The results are shown in Table 3. Root samples 41 to 46, 48 to 51, 58, and 59 are derived from transformation of fad2-1 plants with R1000/pGA Fad2; root samples 52, 53, and 57 were derived from transformation of fad2-1 plants with R1000 and serve as controls; root sample 60 is derived from

transformation of wild type Arabidopsis with R1000 and also serves as a control.

TABLE 3

Fatty acid Composition in Transgenic  
Arabidopsis fad2-1 Hairy Roots Transformed  
with Agrobacterium R1000/pGA-fad2

<u>Sample</u>	<u>16:0</u>	<u>16:1</u>	<u>18:0</u>	<u>18:1</u>	<u>18:2</u>	<u>18:3</u>
41	24.4	1.8	1.7	5.0	29.4	33.8
42	25.6	3.7	1.3	20.0	22.0	27.5
43	23.6	-	1.6	7.2	27.6	36.1
44	24.4	1.3	4.6	16.0	18.1	33.6
45	20.7	-	8.1	44.7	11.8	14.8
46	20.1	-	1.8	7.5	33.7	36.0
48	26.1	2.9	2.1	9.5	17.6	33.4
49	30.8	1.0	2.4	8.7	18.7	31.1
50	19.8	1.9	3.3	27.7	21.8	24.4
51	20.9	1.1	5.0	13.7	25.0	32.1
58	23.5	0.3	1.4	3.6	22.1	45.9
59	22.6	0.6	1.4	2.8	29.9	40.4
52, cont.	12.3	-	2.6	64.2	4.6	16.4
53, cont.	20.3	9.1	2.2	55.2	1.7	9.2
57, cont.	10.4	2.4	0.7	65.9	3.8	12.7
60, WT	23.0	1.7	0.8	6.0	35.0	31.8

- 5        These results show that expression of Arabidopsis  
microsomal delta-12 desaturase in a mutant Arabidopsis  
lacking delta-12 desaturation can result in partial to  
complete complementation of the mutant. The decrease in  
oleic acid levels in transgenic roots is accompanied by  
10    increases in the levels of both 18:2 and 18:3. Thus,  
overexpression of this gene in other oil crops, especially  
canola, which is a close relative of Arabidopsis and which  
naturally has high levels of 18:1 in seeds, is also expected  
to result in higher levels of 18:2, which in conjunction with

the overexpression of the microsomal delta-15 fatty acid desaturase will result in very high levels of 18:3.

Using Arabidopsis Microsomal Delta-12 Desaturase  
cDNA as a Hybridization Probe to Isolate Microsomal  
 5 Delta-12 Desaturase cDNAs from Other Plant Species

Evidence for conservation of the delta-12 desaturase sequences amongst species was provided by using the Arabidopsis cDNA insert from pSF2b as a hybridization probe to clone related sequences from  
 10 Brassica napus, and soybean. Furthermore, corn and castor bean microsomal delta-12 fatty acid desaturase were isolated by PCR using primers made to conserved regions of microsomal delta-12 desaturases.

Cloning of a Brassica napus Seed  
 15 cDNA Encoding Seed Microsomal Delta-12  
Fatty Acid Desaturase

For the purpose of cloning the Brassica napus seed cDNA encoding a delta-12 fatty acid desaturase, the cDNA insert from pSF2b was isolated by digestion of pSF2b  
 20 with EcoR I followed by purification of the 1.2 kb insert by gel electrophoresis. The 1.2 kb fragment was radiolabeled and used as a hybridization probe to screen a lambda phage cDNA library made with poly A<sup>+</sup> mRNA from developing Brassica napus seeds 20-21 days after  
 25 pollination. Approximately 600,000 plaques were screened under low stringency hybridization conditions (50 mM Tris pH 7.6, 6X SSC, 5X Denhardt's, 0.5% SDS, 100 ug denatured calf thymus DNA and 50°C) and washes (two washes with 2X SSC, 0.5% SDS at room temperature  
 30 for 15 min each, then twice with 0.2X SSC, 0.5% SDS at room temperature for 15 min each, and then twice with 0.2X SSC, 0.5% SDS at 50°C for 15 min each). Ten strongly-hybridizing phage were plaque-purified and the size of the cDNA inserts was determined by PCR  
 35 amplification of the insert using phage as template and

009207" 6262660

T3/T7 oligomers for primers. Two of these phages, 165D and 165F, had PCR amplified inserts of 1.6 kb and 1.2 kb respectively and these phages were also used to excise the phagemids as described above. The phagemid derived from phage 165D, designated pCF2-165D, contained a 1.5 kb insert which was sequenced completely on one strand. SEQ ID NO:3 shows the 5' to 3' nucleotide sequence of 1394 base pairs of the *Brassica napus* cDNA which encodes delta-12 desaturase in plasmid pCF2-165d. Nucleotides 99 to 101 and nucleotides 1248 to 1250 are, respectively, the putative initiation codon and the termination codon of the open reading frame (nucleotides 99 to 1250). Nucleotides 1 to 98 and 1251 to 1394 are, respectively, the 5' and 3' untranslated nucleotides. The 383 amino acid protein sequence deduced from the open reading frame in SEQ ID NO:3 is shown in SEQ ID NO:4. While the other strand can easily be sequenced for confirmation, comparisons of SEQ ID NOS:1 and 3 and of SEQ ID NOS:2 and 4, even admitting of possible sequencing errors, showed an overall homology of approximately 84% at both the nucleotide and amino acid levels, which confirmed that pCF2-165D is a *Brassica napus* seed cDNA that encoded delta-12 desaturase. Plasmid pCF2-165D has been deposited on October 16, 1992 with the American Type Culture Collection of Rockville, Maryland under the provisions of the Budapest Treaty and bears accession number ATCC 69094.

Cloning of a Soybean (*Glycine max*)  
cDNA Encoding Seed Microsomal Delta-12  
Fatty Acid Desaturase

A cDNA library was made to poly A<sup>+</sup> mRNA isolated from developing soybean seeds, and screened as described above.. Radiolabelled probe prepared from pSF2b as described above was added, and allowed to hybridize for 18 h at 50°C. The probes were washed as described

above. Autoradiography of the filters indicated that there were 14 strongly hybridizing plaques, and numerous weakly hybridizing plaques. Six of the 14 strongly hybridizing plaques were plaque purified as described above and the cDNA insert size was determined by PCR amplification of the insert using phage as template and T3/T7 oligomers for primers. One of these phages, 169K, had an insert sizes of 1.5 kb and this phage was also used to excise the phagemid as described above. The phagemid derived from phage 169K, designated pSF2-169K, contained a 1.5 kb insert which was sequenced completely on both strands. SEQ ID NO:5 shows the 5' to 3' nucleotide sequence of 1473 base pairs of soybean (Glycine max) cDNA which encodes delta-12 desaturase in plasmid pSF2-169K. Nucleotides 108 to 110 and nucleotides 1245 to 1247 are, respectively, the putative initiation codon and the termination codon of the open reading frame (nucleotides 108 to 1247). Nucleotides 1 to 107 and 1248 to 1462 are, respectively, the 5' and 3' untranslated nucleotides. The 380 amino acid protein sequence deduced from the open reading frame in SEQ ID NO:5 is shown in SEQ ID NO:6. Comparisons of SEQ ID NOS:1 and 5 and of SEQ ID NOS:2 and 6, even admitting of possible sequencing errors, showed an overall homology of approximately 65% at the nucleotide level and approximately 70% at the amino acid level, which confirmed that pSF2-169K is a soybean seed cDNA that encoded delta-12 desaturase. A further description of this clone can be obtained by comparison of the SEQ ID NO:1, SEQ ID NO:3, and SEQ ID NO:5 and by analyzing the phenotype of transgenic plants produced by using chimeric genes incorporating the insert of pSF2-169K, in sense or antisense orientation, with suitable regulatory sequences. Plasmid pSF2-169K was deposited on October 16, 1992 with the American Type Culture

009201" 52E25950

Collection of Rockville, Maryland under the provisions of the Budapest Treaty and bears accession number ATCC 69092.

Cloning of a Corn (Zea mays)

5                   cDNA Encoding Seed Microsomal Delta-12  
                     Fatty Acid Desaturase

Corn microsomal delta-12 desaturase cDNA was isolated using a PCR approach. For this, a cDNA library was made to poly A<sup>+</sup> RNA from developing corn embryos in  
 10 Lambda Zap II vector. This library was used as template for PCR using sets of degenerate oligomers NS3 (SEQ ID NO:13) and RB5A/B (SEQ ID NOS:16 and 17) as sense and antisense primers, respectively. NS3 and RB5A/B correspond to stretches of amino acids 101-109 and  
 15 318-326, respectively, of SEQ ID NO:2, which are conserved in most microsomal delta-12 desaturases (for example, SEQ ID NOS:2, 4, 6, 8). PCR was carried out using a PCR kit (Perkin-Elmer) by 40 cycles of 94°C 1', 45°C, 1 min, and 55°C, 2 min. Analyses of the PCR  
 20 products on an agarose gel showed the presence of a product of the expected size (720 bp), which was absent in control reactions containing either the sense or antisense primers alone. The fragment was gel purified and then used as a probe for screening the corn cDNA  
 25 library at 60°C as described above. One positively-hybridizing plaque was purified and partial sequence determination of its cDNA showed it to be a nucleotide sequence encoding microsomal delta-12 desaturase but truncated at the 3' end. The cDNA insert encoding the  
 30 partial desaturase was gel isolated and used to probe the corn cDNA library again. Several positive plaques were recovered and characterized. DNA sequence analysis revealed that all of these clones seem to represent the same sequence with the different length of 5' or 3'  
 35 ends. The clone containing the longest insert,

009201"52E26950

designated pFad2#1, was sequenced completely. The total length of the cDNA is 1790 bp (SEQ ID NO:7) comprising of an open reading frame from nucleotide 165 to 1328 bp that encoded a polypeptide of 388 amino acids. The deduced amino acid sequence of the polypeptide (SEQ ID NO:8) shared overall identities of 71%, 40%, and 38% to Arabidopsis microsomal delta-12 desaturase, Arabidopsis microsomal delta-15 desaturase, and Arabidopsis plastid delta-15 desaturase, respectively. Furthermore, it lacked an N-terminal amino acid extension that would indicate it is a plastid enzyme. Based on these considerations, it is concluded that it encodes a microsomal delta-12 desaturase.

#### Isolation of cDNAs Encoding

##### Delta-12 Microsomal Fatty Acid Desaturases and Desaturase-Related Enzymes from Castor Bean Seed

Polysomal mRNA was isolated from castor beans of stages I-II (5-10 DAP) and also from castor beans of stages IV-V (20-25 DAP). Ten ng of each mRNA was used for separate RT-PCR reactions, using the Perkin-Elmer RT-PCR kit. The reverse transcriptase reaction was primed with random hexamers and the PCR reaction with degenerate delta-12 desaturase primers NS3 and NS9 (SEQ ID NOS:13 and 14). The annealing-extension temperature of the PCR reaction was 50°C. A DNA fragment of approx. 700 bp was amplified from both stage I-II and stage IV-V mRNA. The amplified DNA fragment from one of the reactions was gel purified and cloned into a pGEM-T vector using the Promega pGEM-T PCR cloning kit to create the plasmid pRF2-1C. The 700 bp insert in pRF2-1C was sequenced, as described above, and the resulting DNA sequence is shown in SEQ ID NO:9. The DNA sequence in SEQ ID NO:9 contains an open-reading frame encoding 219 amino acids (SEQ ID NO:10) which has 81% identity (90% similarity) with amino acids 135 to 353 of

the Arabidopsis microsomal delta-12 desaturase described in SEQ ID NO:2. The cDNA insert in pRF2-1C is therefore a 676 bp fragment of a full-length cDNA encoding a castor bean seed microsomal delta-12 desaturase. The full length castor bean seed microsomal delta-12 desaturase cDNA may be isolated by screening a castor seed cDNA library, at 60°C, with the labeled insert of pRF2-1C as described in the example above. The insert in pRF2-1C may also be used to screen castor bean libraries at lower temperatures to isolate delta-12 desaturase-related sequences, such as the delta-12 hydroxylase.

A cDNA library made to poly A<sup>+</sup> mRNA isolated from developing castor beans (stages IV-V, 20-25 DAP) was screened as described above. Radiolabeled probe prepared from pSF2b or pRF2-1C, as described above, were added, and allowed to hybridize for 18 h at 50°C. The filters were washed as described above. Autoradiography of the filters indicated that there were numerous hybridizing plaques, which appeared either strongly-hybridising or weakly-hybridising. Three of the strongly hybridising plaques (190A-41, 190A-42 and 190A-44) and three of the weakly hybridising plaques, (190B-41, 190b-43 and 197c-42), were plaque purified using the methods described above. The cDNA insert size of the purified phages were determined by PCR amplification of the insert using phage as template and lambda-gt11 oligomers (Clontech lambda-gt11 Amplimers) for primers. The PCR-amplified inserts of the amplified phages were subcloned into pBluescript (Pharmacia) which had been cut with Eco RI and filled in with Klenow (Sambrook et al. (Molecular Cloning, A Laboratory Approach, 2nd. ed. (1989) Cold Spring Harbor Laboratory Press). The resulting plasmids were called pRF190a-41, pRF190a-42, pRF190a-44, pRF190b-41, pRF190b-43 and



pRF197c-42. All of the inserts were about 1.1 kb with the exception of pRF197c-42 which was approx. 1.5 kb. The inserts in the plasmids were sequenced as described above. The insert in pRF190b-43 did not contain any open reading frame and was not identified. The inserts in pRF190a-41, pRF190a-42, pRF190a-44 and pRF190b-41 were identical. The insert in pRF197c-42 contained all of the nucleotides of the inserts in pRF190a-41, pRF190a-42, pRF190a-44 and pRF190b-41 plus an additional approx. 400 bp. It was deduced therefore that the insert in pRF197c-42 was a longer version of the inserts in pRF190a-41, pRF190a-42, pRF190a-44 and pRF190b-41 and all were derived from the same full-length mRNA. The complete cDNA sequence of the insert in plasmid pRF197c-42 is shown in SEQ ID NO:11. The deduced amino acid sequence of SEQ ID NO:11, shown in SEQ ID NO:12, is 78.5% identical (90% similarity) to the castor microsomal delta-12 desaturase described above (SEQ ID NO:10) and 66% identical (80% similarity) to the Arabidopsis delta-12 desaturase amino acid sequence in SEQ ID NO:2. These similarities confirm that pRF197c-42 is a castor bean seed cDNA that encodes a microsomal delta-12 desaturase or a microsomal delta-12 desaturase-related enzyme, such as a delta-12 hydroxylase. Specific PCR primers for pRF2-1C and pRF197c-42 were made. For pRF2-1c the upstream primer was bases 180 to 197 of the cDNA sequence in SEQ ID NO:9. For pRF197c-42 the upstream primer was bases 717 to 743 of the cDNA sequence in SEQ ID NO:11. A common downstream primer was made corresponding to the exact complement of the nucleotides 463 to 478 of the sequence described in SEQ ID NO:9. Using RT-PCR with random hexamers and the above primers it was observed that the cDNA contained in pRF2-1C is expressed in both Stage I-II and Stage IV-V castor bean seeds whereas the cDNA contained in plasmid

pRF197c-42 is expressed only in Stage IV-V castor bean seeds, i.e., it is only expressed in tissue actively synthesizing ricinoleic acid. Thus, it is possible that this cDNA encodes a delta-12 hydroxylase.

5        There is enough deduced amino acid sequence from the two castor sequences described in SEQ ID NOS:10 and 12 to compare the highly conserved region corresponding to amino acids 311 to 353 of SEQ ID NO:2. When SEQ ID NOS:2, 4, 6, 8, and 10 are aligned by the Hein method  
10        described above the consensus sequence corresponds exactly to the amino acids 311 to 353 of SEQ ID NO:2. All of the seed microsomal delta-12 desaturases described above have a high degree of identity with the consensus over this region, namely Arabidopsis (100%  
15        identity), soybean (90% identity), corn (95% identity), canola (93% identity) and one (pRF2-1c) of the castor bean sequences (100% identity). The other castor bean seed delta-12 desaturase or desaturase-related sequence (pRF197c-42) however has less identity with the  
20        consensus, namely 81% for the deduced amino acid sequence of the insert in pRF197c-42 (described in SEQ ID NO:12). Thus while it remains possible that the insert in pRF197c-42 encodes a microsomal delta-12 desaturase, this observation supports the hypothesis  
25        that it encodes a desaturase-related sequence, namely the delta-12 hydroxylase.

      An additional approach to cloning a castor bean seed delta-12 hydroxylase is the screening of a differential population of cDNAs. A lambda-Zap  
30        (Stratagene) cDNA library made to polysomal mRNA isolated from developing castor bean endosperm (stages IV-V, 20-25 DAP) was screened with <sup>32</sup>P-labeled cDNA made from polysomal mRNA isolated from developing castor bean endosperm (stage I-II, 5-10 DAP) and with <sup>32</sup>P-labeled  
35        cDNA made from polysomal mRNA isolated from developing

009201" 52E26950

castor bean endosperm (stages IV-V, 20-25 DAP). The library was screened at a density of 2000 plaques per 137 mm plate so that individual plaques were isolated. About 60,000 plaques were screened and plaques which  
 5 hybridised with late (stage IV/V) cDNA but not early (stage I/II) cDNA, which corresponded to about 1 in every 200 plaques, were pooled.

The library of differentially expressed cDNAs may be screened with the castor delta-12 desaturase cDNA  
 10 described above and/or with degenerate oligonucleotides based on sequences of amino conserved among delta-12 desaturases to isolate related castor cDNAs, including the cDNA encoding the delta-12 oleate hydroxylase enzyme. These regions of amino acid conservation may  
 15 include, but are not limited to amino acids 101 to 109, 137 to 145, and 318 to 327 of the amino acid sequence described in SEQ ID NO:2 or any of the sequences described in Table 7 below. Examples of such oligomers  
 20 are SEQ ID NOS:13, 14, 16, and 17. The insert in plasmid pCF2-197c may be cut with Eco RI to remove vector sequences, purified by gel electrophoresis and labeled as described above. Degenerate oligomers based on the above conserved amino acid sequences may be  
 25 labeled with  $^{32}\text{P}$  as described above. The cDNAs cloned from the developing endosperm difference library which do not hybridize with early mRNA probe but do hybridize with late mRNA probe and hybridize with either castor delta-12 desaturase cDNA or with an oligomer based on delta-12 desaturase sequences are likely to be the  
 30 castor delta-12 hydroxylase. The pBluescript vector containing the putative hydroxylase cDNA can be excised and the inserts directly sequenced, as described above.

Clones such as pRF2-1C and pRF197c-42, and other clones from the differential screening, which, based on  
 35 their DNA sequence, are less related to castor bean seed

009920T 62626959

microsomal delta-12 desaturases and are not any of the known fatty-acid desaturases described above or in WO 9311245, may be expressed, for example, in soybean embryos or another suitable plant tissue, or in a  
 5 microorganism, such as yeast, which does not normally contain ricinoleic acid, using suitable expression vectors and transformation protocols. The presence of novel ricinoleic acid in the transformed tissue(s) expressing the castor cDNA would confirm the identity of  
 10 the castor cDNA as DNA encoding for an oleate hydroxylase.

Sequence Comparisons Among Seed Microsomal  
Delta-12 Desaturases

The percent overall identities between coding  
 15 regions of the full-length nucleotide sequences encoding microsomal delta-12 desaturases was determined by their alignment by the method of Needleman et al. (J. Mol. Biol. (1970) 48:443-453) using gap weight and gap length weight values of 5.0 and 0.3 (Table 4). Here, a2, c2,  
 20 s2, z2 and des A refer, respectively, to the nucleotide sequences encoding microsomal delta-12 fatty acid desaturases from Arabidopsis (SEQ ID NO:1), Brassica napus (SEQ ID NO:3), soybean (SEQ ID NO:5), corn (SEQ ID NO:7), and cyanobacterial des A, whereas r2 refers to  
 25 the microsomal delta-12 desaturase or desaturase-related enzyme from castor bean (SEQ ID NO:12).

TABLE 4

Percent Identity Between the Coding Regions of  
Nucleotide Sequences Encoding Different Microsomal  
Delta-12 Fatty Acid Desaturases

	<u>c2</u>	<u>s2</u>	<u>z2</u>	<u>des A</u>
a2	84	66	64	43
c2	-	65	62	42
s2	-	-	62	42

The overall relatedness between the deduced amino acid sequences of microsomal delta-12 desaturases or desaturase-related enzymes of the invention (i.e., SEQ ID NOS:2, 4, 6, 8, and 12) determined by their alignment by the method of Needleman et al. (J. Mol. Biol. (1970) 48:443-453) using gap weight and gap length weight values of 3.0 and 0.1, respectively, is shown in Table 5. Here a2, c2, s2, z2 and des A refer, respectively, to microsomal delta-12 fatty acid desaturases from Arabidopsis (SEQ ID NO:2), Brassica napus (SEQ ID NO:4), soybean (SEQ ID NO:6), corn (SEQ ID NO:8), and cyanobacterial des A, whereas r2 refers to the microsomal desaturase or desaturase-related enzyme from castor bean (SEQ ID NO:12). The relatedness between the enzymes is shown as percent overall identity/percent overall similarity.

TABLE 5

Relatedness Between Different Microsomal  
Delta-12 Fatty Acid Desaturases

	<u>c2</u>	<u>s2</u>	<u>r2</u>	<u>z2</u>	<u>des A</u>
a2	84/89	70/85	66/80	71/83	24/50
c2	-	67/80	63/76	69/79	24/51
s2	-	-	67/83	66/82	23/49
r2	-	-	-	61/78	24/51
z2	-	-	-	-	25/49

The high degree of overall identity (60% or greater) at the amino acid levels between the Brassica napus, soybean, castor and corn enzymes with that of Arabidopsis microsomal delta-12 desaturase and their lack of an N-terminal extension of a transit peptide expected for a nuclear-encoded chloroplast desaturase leads Applicants to conclude that SEQ ID NOS:4, 6, 8, 10, and 12 encode the microsomal delta-12

009207" 62E/6950

desaturases or desaturase-related enzymes. Further confirmation of this identity will come from biological function, that is, by analyzing the phenotype of transgenic plants or other organisms produced by using chimeric genes incorporating the above-mentioned sequences in sense or antisense orientation, with suitable regulatory sequences. Thus, one can isolate cDNAs and genes for homologous fatty acid desaturases from the same or different higher plant species, especially from the oil-producing species. Furthermore, based on these comparisons, the Applicants expect all higher plant microsomal delta-12 desaturases from all higher plants to show an overall identity of 60% or more and to be able to readily isolate homologous fatty acid desaturase sequences using SEQ ID NOS:1, 3, 5, 7, 9, and 11 by sequence-dependent protocols.

The overall percent identity at the amino acid level, using the above alignment method, between selected plant desaturases is illustrated in Table 6.

**TABLE 6**  
Percent Identity Between Selected Plant Fatty Acid Desaturases at the Amino Acid Level

	<u>a3</u>	<u>aD</u>	<u>c3</u>	<u>cD</u>	<u>S3</u>
a2	38	33	38	35	34
a3	-	65	93	66	67
aD	-	-	66	87	65
c3	-	-	-	67	67
cD	-	-	-	-	65

In Table 6, a2, a3, aD, c3, cD, and S3 refer, respectively, to SEQ ID NO:2, Arabidopsis microsomal delta-15 desaturase, Arabidopsis plastid delta-15 desaturase, canola microsomal delta-15 desaturase, canola plastid delta-15 desaturase, and soybean microsomal delta-15 desaturase. Based on these

comparisons, the delta-15 desaturases, of both  
 microsomal and plastid types, have overall identities of  
 65% or more at the amino acid level, even when from the  
 same plant species. Based on the above the Applicants  
 5 expect microsomal delta-12 desaturases from all higher  
 plants to show similar levels of identity (that is, 60%  
 or more identity at the amino acid level) and that SEQ  
 ID NOS:1, 3, 5, 7, and 9 can also be used as  
 hybridization probe to isolate homologous delta-12  
 10 desaturase sequences, and possibly for sequences for  
 fatty acid desaturase-related enzymes, such as oleate  
 hydroxylase, that have an overall amino acid homology of  
 50% or more.

Similar alignments of protein sequences of plant  
 15 microsomal fatty acid delta-12 desaturases [SEQ ID  
 NOS:2, 4, 6, and 8] and plant delta-15 desaturases  
 [microsomal and plastid delta-15 desaturases from  
Arabidopsis and Brassica napus, WO 9311245] allows  
 identification of amino acid sequences conserved between  
 20 the different desaturases (Table 7).

**TABLE 7**  
 Amino Acid Sequences Conserved Between  
 Plant Microsomal Delta-12 Desaturases and Microsomal and  
Plastid Delta-15 Desaturases

Region	Conserved AA Positions in SEQ ID NO:2	Consensus Conserved AA Sequence in $\Delta^{12}$ Desaturases	Consensus Conserved AA Sequence in $\Delta^{15}$ Desaturases	Consensus AA Sequence
A	39-44	<u>AIPPHC</u>	<u>AIPKHC</u>	AIP(P/K)HC
B	86-90	<u>WP(L/I)YW</u>	<u>WPLYW</u>	WP(L/I)YW
C	104-109	<u>AHECGH</u>	<u>GHDCGH</u>	(A/G)H(D/E)CGH
D	130-134	<u>LLVPY</u>	<u>ILVPY</u>	(L/I)LVPY
E	137-142	<u>WKYSHR</u>	<u>WRISHR</u>	W(K/R)(Y/I)SHR
F	140-145	<u>SHRRHH</u>	<u>SHRTHH</u>	SHR(R/T)HH
G	269-274	<u>ITYLQ</u>	<u>VTYLH</u>	(I/V)TYL(Q/H)
H	279-282	<u>LPHY</u>	<u>LPWY</u>	LP(H/W)Y

I	289-294	<u>WL(R/K)GAL</u>	<u>YLRGGL</u>	(W/Y)L(R/K)G(A/G)L
J	296-302	<u>TVDRDYG</u>	<u>TLDRDYG</u>	T(V/L)DRDYG
K	314-321	<u>THVAHHLF</u>	<u>THVIHHLF</u>	THV(A/I)HHLF
L	318-327	<u>HHLFSTMPHY</u>	<u>HHLFQIPHY</u>	HHFL(S/P) (T/Q)(I/M)PHY

Table 7 shows twelve regions of conserved amino acid sequences, designated A-L (column 1), whose positions in SEQ ID NO:2 are shown in column 2. The consensus sequences for these regions in plant delta-12 fatty acid desaturases and plant delta-15 fatty acid desaturases are shown in columns 3 and 4, respectively; amino acids are shown by standard abbreviations, the underlined amino acids are conserved between the delta-12 and the delta-15 desaturases, and amino acids in brackets represent substitutions found at that position. The consensus sequence of these regions are shown in column 5. These short conserved amino acids and their relative positions further confirm that the isolated isolated cDNAs encode a fatty acid desaturase.

Isolation of Nucleotide Sequences Encoding  
Homologous and Heterologous Fatty acid Desaturases  
and Desaturase-like Enzymes

Fragments of the instant invention may be used to isolate cDNAs and genes of homologous and heterologous fatty acid desaturases from the same species as the fragments of the invention or from different species. Isolation of homologous genes using sequence-dependent protocols is well-known in the art and Applicants have demonstrated that Arabidopsis microsomal delta-12 desaturase cDNA sequence can be used to isolate cDNA for related fatty acid desaturases from Brassica napus, soybean, corn and castor bean.

More importantly, one can use the fragments containing SEQ ID NOS:1, 3, 5, 7, and 9 or their



smaller, more conserved regions to isolate novel fatty acid desaturases and fatty acid desaturase-related enzymes.

5 In a particular embodiment of the present invention, regions of the nucleic acid fragments of the invention that are conserved between different desaturases may be used by one skilled in the art to design a mixture of degenerate oligomers for use in sequence-dependent protocols aimed at isolating nucleic acid fragments encoding homologous or heterologous fatty acid desaturase cDNA's or genes. For example, in the polymerase chain reaction (Innis, et al., Eds, (1990) PCR Protocols: A Guide to Methods and Applications, Academic Press, San Diego), two short pieces of the present fragment of the invention can be used to amplify a longer fatty acid desaturase DNA fragment from DNA or RNA. The polymerase chain reaction may also be performed on a library of cloned nucleotide sequences with one primer based on the fragment of the invention and the other on either the poly A<sup>+</sup> tail or a vector sequence. These oligomers may be unique sequences or degenerate sequences derived from the nucleic acid fragments of the invention. The longer piece of homologous fatty acid desaturase DNA generated by this method could then be used as a probe for isolating related fatty acid desaturase genes or cDNAs from Arabidopsis or other species and subsequently identified by differential screening with known desaturase sequences and by nucleotide sequence determination. The design of oligomers, including long oligomers using deoxyinosine, and "guessmers" for hybridization or for the polymerase chain reaction are known to one skilled in the art and discussed in Sambrook et al., (Molecular Cloning, A Laboratory Manual, 2nd ed. (1989), Cold Spring Harbor Laboratory Press). Short stretches of

10  
15  
20  
25  
30  
35

006673-10260

amino acid sequences that are conserved between cyanobacterial delta-12 desaturase (Wada et al., Nature (1990) 347:200-203) and plant delta-15 desaturases [WO 9311245] were previously used to make oligo-

5 nucleotides that were degenerate and/or used deoxyinosine/s. One set of these oligomers made to a stretch of 12 amino acids conserved between cyanobacterial delta-12 desaturase and higher plant delta-15 desaturases was successful in cloning the

10 plastid delta-12 desaturase cDNAs; these plant desaturases have more than 50% identity to the cyanobacterial delta-12 desaturase. Some of these oligonucleotides were also used as primers to make polymerase chain reaction (PCR) products using poly A<sup>+</sup>

15 RNA from plants. However, none of the oligonucleotides and the PCR products were successful as radiolabeled hybridization probes in isolating nucleotide sequences encoding microsomal delta-12 fatty acid desaturases. Thus, as expected, none of the stretches of four or more

20 amino acids conserved between Arabidopsis delta-12 and Arabidopsis delta-15 desaturases are identical in the cyanobacterial desaturase, just like none of the stretches of four or more amino acids conserved between Arabidopsis delta-15 and the cyanobacterial desaturase

25 are identical in SEQ ID NO:2. Stretches of conserved amino acids between the present invention and delta-15 desaturases now allow for the design of oligomers to be used to isolate sequences encoding other desaturases and desaturase-related enzymes. For example, conserved

30 stretches of amino acids between delta-12 desaturase and delta-15 desaturase, shown in Table 7, are useful in designing long oligomers for hybridization as well as shorter ones for use as primers in the polymerase chain reaction. In this regard, sequences conserved between

35 delta-12 and delta-15 desaturases (shown in Table 7)

009201 5426950

evidence that the high oleic acid trait in mutant 658 is the result of T-DNA insertion.

#### Plasmid Rescue and Analysis

One-half and one microgram of genomic DNA from the  
5 homozygous mutant plants of the 658-75 line, prepared  
from leaf tissue as described [Rogers, S. O. and A. J.  
Bendich (1985) Plant Molecular Biology 5:69-76], was  
digested with 20 units of either Bam HI or Sal I  
restriction enzyme (Bethesda Research Laboratory) in a  
10 50 µL reaction volume according to the manufacturer's  
specifications. After digestion the DNA was extracted  
with buffer-saturated phenol (Bethesda Research  
Laboratory) followed by precipitation in ethanol.  
One-half to one microgram of Bam HI or Sal I digested  
15 genomic DNA was resuspended in 200 uL or 400 uL of  
ligation buffer containing 50 mM Tris-Cl, pH 8.0, 10 mM  
MgCl<sub>2</sub>, 10 mM dithiothreitol, 1 mM ATP, and 4 units of T4  
DNA ligase (Bethesda Research Laboratory). The dilute  
DNA concentration of approximate 2.5 ug/mL in the  
20 ligation reaction was chosen to facilitate  
circularization, as opposed to intermolecular joining.  
The reaction was incubated for 16 h at 16°C. Competent  
DH10B cells (Bethesda Research Laboratory) were  
transfected with 10 ng of ligated DNA per 100 µL of  
25 competent cells according to the manufacturer's  
specifications. Transformants from Sal I or Bam HI  
digests were selected on LB plates (10 g Bacto-tryptone,  
5 g Bacto-yeast extract, 5 g NaCl, 15 g agar per liter,  
pH 7.4) containing 100 µg/mL ampicillin. After  
30 overnight incubation at 37°C the plates were scored for  
ampicillin-resistant colonies.

A single ampicillin-resistant transformant derived  
from Bam HI-digested plant DNA was used to start a  
culture in 35 mL LB medium (10 g Bacto-tryptone, 5 g  
35 yeast-extract, 5 g NaCl per liter) containing 25 mg/L

09697379-102600

ampicillin. The culture was incubated with shaking overnight at 37°C and the cells were then collected by centrifugation at 1000xg for 10 min. Plasmid DNA, designated p658-1, was isolated from the cells by the alkaline lysis method of Birnboim et al. [Nucleic Acid Research (1979) 7:1513-1523], as described in Sambrook et al., (Molecular Cloning, A Laboratory Manual, 2nd ed (1989) Cold Spring Harbor Laboratory Press). Plasmid p658-1 DNA was digested by restriction enzymes Bam HI, Eco RI and Sal I (Bethesda Research Laboratory) and electrophoresed through a 1% agarose gel in 1xTBE buffer (0.089M tris-borate, 0.002M EDTA). The restriction pattern indicated the presence in this plasmid of the expected 14.2 kB T-DNA fragment and a 1.6 kB putative plant DNA/T-DNA border fragment.

#### EXAMPLE 2

#### CLONING OF ARABIDOPSIS THALIANA MICROSOMAL DELTA-12 DESATURASE cDNA USING GENOMIC DNA FLANKING THE T-DNA SITE OF INSERTION IN ARABIDOPSIS THALIANA MUTANT LINE 658-75 AS A HYBRIDIZATION PROBE

Two hundred nanograms of the 1.6 kB Eco RI-Bam HI fragment from plasmid p658-1, following digestion of the plasmid with Eco RI and Bam HI and purification by electrophoresis in agarose, was radiolabelled with alpha[32P]-dCTP using a Random Priming Labeling Kit (Bethesda Research Laboratory) under conditions recommended by the manufacturer.

The radiolabeled DNA was used as a probe to screen an Arabidopsis cDNA library made from RNA isolated from above ground portions of various growth stages (Elledge et al., (1991) Proc. Nat. Acad. Sci., 88:1731-1735) essentially as described in Sambrook et al., (Molecular Cloning, A Laboratory Manual, 2nd ed. (1989), Cold Spring Harbor Laboratory Press). For this, approximately 17,000 plaque-forming units were plated on

0969739.102600

seven 90mm petri plates containing a lawn of LE392 E. coli cells on NZY agar media (5 g NaCl, 2 g MgSO<sub>4</sub>-7 H<sub>2</sub>O, 5 g yeast extract, 10 g casein acid hydrolysate, 13 g agar per liter). Replica filters of the phage

5     plaques were prepared by adsorbing the plaques onto nitrocellulose filters (BA85, Schleicher and Schuell) then soaking successively for five min each in 0.5 M NaOH/1 M NaCl, 0.5 M Tris(pH 7.4)/1.5 M NaCl and 2xSSPE (0.36 M NaCl, 20 mM NaH<sub>2</sub>PO<sub>4</sub>(pH 7.4), 20 mM EDTA

10    (pH 7.4)). The filters were then air dried and baked for 2 h at 80°C. After baking the filters were wetted in 2X SSPE, and then incubated at 42°C in prehybridization buffer (50% Formamide, 5X SSPE, 1% SDS, 5X Denhardt's Reagent, and 100 ug/mL denatured salmon

15    sperm DNA) for 2 h. The filters were removed from the prehybridization buffer, and then transferred to hybridization buffer (50% Formamide, 5X SSPE, 1% SDS, 1X Denhardt's Reagent, and 100 ug/mL denatured salmon sperm DNA) containing the denatured radiolabeled probe (see

20    above) and incubated for 40 h at 42°C. The filters were washed three times in 2X SSPE/0.2% SDS at 42°C (15 min each) and twice in 0.2X SSPE/0.2% SDS at 55°C (30 min each), followed by autoradiography on Kodak XAR-5 film with an intensifying screen at -80°C, overnight.

25    Fifteen plaques were identified as positively-hybridizing on replica filters. Five of these were subjected to plaque purification essentially as described in Sambrook et al., (Molecular Cloning, A Laboratory Manual, 2nd ed. (1989), Cold Spring Harbor

30    Laboratory Press). The lambda YES-R cDNA clones were converted to plasmid by propagating the phage in the E. coli BNN-132 cells, which expresses Cre protein that excises the cDNA insert as a double-stranded plasmid by cre-mediated in vivo site-specific recombination at a

35    'lox' sites present in the phage. Ampicillin-resistant

plasmid clones containing cDNA inserts were grown in liquid culture, and plasmid DNA was prepared using the alkaline lysis method as previously described. The sizes of the resulting plasmids were analyzed by electrophoresis in agarose gels. The agarose gels were treated with 0.5 M NaOH/1 M NaCl, and 0.5 M Tris(pH 7.4), 1.5 M NaCl for 15 min each, and the gel was then dried completely on a gel drier at 65°C. The gel was hydrated in 2X SSPE and incubated overnight, at 42°C, in hybridization buffer containing the denatured radiolabeled probe, followed by washing as described above. After autoradiography, the inserts of four of the purified cDNA clones were found to have hybridized to the probe. Plasmid DNA from the hybridizing clones was purified by equilibration in a CsCl/ethidium bromide gradient (see above). The four cDNA clones were sequenced using Sequenase T7 DNA polymerase (US Biochemical Corp.) and the manufacturer's instructions, beginning with primers homologous to vector sequences that flank the cDNA insert. After comparing the partial sequences of the inserts obtained from the four clones, it was apparent that they each contained sequences in common. One cDNA clone, p92103, containing ca. 1.4 kB cDNA insert, was sequenced. The longest three clones were subcloned into the plasmid vector pBluescript (Stratagene). One of these clones, designated pSF2b, containing ca 1.2 kB cDNA insert was also sequenced serially with primers designed from the newly acquired sequences as the sequencing experiment progressed. The composite sequence derived from pSF2b and p92103 is shown in SEQ ID NO:1.

03697379.102600

EXAMPLE 3CLONING OF PLANT FATTY ACIDDESATURASE cDNAs USING THE ARABIDOPSIS THALIANAMICROSOMAL DELTA-12 DESATURASE cDNA CLONE AS A5 HYBRIDIZATION PROBE

An approximately 1.2 kb fragment containing the Arabidopsis delta-12 desaturase coding sequence of SEQ ID NO:1 was obtained from plasmid pSF2b. This plasmid was digested with EcoR I and the 1.2 kb delta-12  
 10 desaturase cDNA fragment was purified from the vector sequence by agarose gel electrophoresis. The fragment was radiolabelled with <sup>32</sup>P as previously described.

Cloning of a Brassica napus SeedcDNA Encoding Microsomal Delta-12 Fatty Acid15 Desaturase

The radiolabelled probe was used to screen a Brassica napus seed cDNA library. In order to construct the library, Brassica napus seeds were harvested 20-21 days after pollination, placed in liquid nitrogen, and  
 20 polysomal RNA was isolated following the procedure of Kamalay et al., (Cell (1980) 19:935-946). The polyadenylated mRNA fraction was obtained by affinity chromatography on oligo-dT cellulose (Aviv et al., Proc. Natl. Acad. Sci. USA (1972) 69:1408-1411). Four  
 25 micrograms of this mRNA were used to construct a seed cDNA library in lambda phage (Uni-ZAP™ XR vector) using the protocol described in the ZAP-cDNA™ Synthesis Kit (1991 Stratagene Catalog, Item #200400). Approximately 600,000 clones were screened for positively hybridizing  
 30 plaques using the radiolabelled EcoR I fragment from pSF2b as a probe essentially as described in Sambrook et al., (Molecular Cloning: A Laboratory Manual, 2nd ed. (1989) Cold Spring Harbor Laboratory Press) except that low stringency hybridization conditions (50 mM Tris, pH  
 35 7.6, 6X SSC, 5X Denhardt's, 0.5% SDS, 100 µg denatured

009207 5225950

calf thymus DNA and 50°C) were used and post-hybridization washes were performed twice with 2X SSC, 0.5% SDS at room temperature for 15 min, then twice with 0.2X SSC, 0.5% SDS at room temperature for 15 min, and then twice with 0.2X SSC, 0.5% SDS at 50°C for 15 min. Ten positive plaques showing strong hybridization were picked, plated out, and the screening procedure was repeated. From the secondary screen nine pure phage plaques were isolated. Plasmid clones containing the cDNA inserts were obtained through the use of a helper phage according to the in vivo excision protocol provided by Stratagene. Double-stranded DNA was prepared using the alkaline lysis method as previously described, and the resulting plasmids were size-analyzed by electrophoresis in agarose gels. The largest one of the nine clones, designated pCF2-165D, contained an approximately 1.5 kb insert which was sequenced as described above. The sequence of 1394 bases of the cDNA insert of pCF2-165D is shown in SEQ ID NO:3. Contained in the insert but not shown in SEQ ID NO:3 are approximately 40 bases of the extreme 5' end of the 5' non-translated region and a poly A tail of about 38 bases at the extreme 3' end of the insert.

#### Cloning of a Soybean Seed

#### 25 cDNA Encoding Microsomal Delta-12

#### Fatty Acid Desaturase

A cDNA library was made as follows: Soybean embryos (ca. 50 mg fresh weight each) were removed from the pods and frozen in liquid nitrogen. The frozen embryos were ground to a fine powder in the presence of liquid nitrogen and then extracted by Polytron homogenization and fractionated to enrich for total RNA by the method of Chirgwin et al. (Biochemistry (1979) 18:5294-5299). The nucleic acid fraction was enriched for poly A<sup>+</sup>RNA by passing total RNA through an oligo-dT



cellulose column and eluting the poly A<sup>+</sup>RNA with salt as described by Goodman et al. (Meth. Enzymol. (1979) 68:75-90). cDNA was synthesized from the purified poly A<sup>+</sup>RNA using cDNA Synthesis System (Bethesda Research Laboratory) and the manufacturer's instructions. The resultant double-stranded DNA was methylated by Eco RI DNA methylase (Promega) prior to filling-in its ends with T4 DNA polymerase (Bethesda Research Laboratory) and blunt-end ligation to phosphorylated Eco RI linkers using T4 DNA ligase (Pharmacia). The double-stranded DNA was digested with Eco RI enzyme, separated from excess linkers by passage through a gel filtration column (Sephadex CL-4B), and ligated to lambda ZAP vector (Stratagene) according to manufacturer's instructions. Ligated DNA was packaged into phage using the Gigapack packaging extract (Stratagene) according to manufacturer's instructions. The resultant cDNA library was amplified as per Stratagene's instructions and stored at -80°C.

Following the instructions in the Lambda ZAP Cloning Kit Manual (Stratagene), the cDNA phage library was used to infect *E. coli* BB4 cells and approximately 600,000 plaque forming units were plated onto 150 mm diameter petri plates. Duplicate lifts of the plates were made onto nitrocellulose filters (Schleicher & Schuell). The filters were prehybridized in 25 mL of hybridization buffer consisting of 6X SSPE, 5X Denhardt's solution, 0.5% SDS, 5% dextran sulfate and 0.1 mg/mL denatured salmon sperm DNA (Sigma Chemical Co.) at 50°C for 2 h. Radiolabelled probe prepared from pSF2b as described above was added, and allowed to hybridize for 18 h at 50°C. The filters were washed exactly as described above. Autoradiography of the filters indicated that there were 14 strongly hybridizing plaques. The 14 plaques were subjected to a

second round of screening as before. Numerous, strongly hybridizing plaques were observed on 6 of the 14 filters, and one, well-isolated from other phage, was picked from each of the six plates for further analysis.

5       Following the Lambda ZAP Cloning Kit Instruction Manual (Stratagene), sequences of the pBluescript vector, including the cDNA inserts, from the purified phages were excised in the presence of a helper phage and the resultant phagemids were used to infect *E. coli* 10 XL-1 Blue cells. DNA from the plasmids was made by the Promega "Magic Miniprep" according to the manufacturers instructions. Restriction analysis indicated that the plasmids contained inserts ranging in size from 1 kb to 2.5 kb. The alkali-denatured double-stranded DNA from 15 one of these, designated pSF2-169K contained an insert of 1.6 kb, was sequenced as described above. The nucleotide sequence of the cDNA insert in plasmid pSF2-169K shown in SEQ ID NO:5.

Cloning of a Corn (*Zea mays*)

20       cDNA Encoding Seed Microsomal Delta-12  
           Fatty Acid Desaturase

Corn microsomal delta-12 desaturase cDNA was isolated using a PCR approach. For this, a cDNA library was made to poly A<sup>+</sup> RNA from developing corn embryos in 25 Lambda ZAP II vector (Stratagene). 5-10 ul of this library was used as a template for PCR using 100 pmol each of two sets of degenerate oligomers NS3 (SEQ ID NO:13) and equimolar amounts of RB5a/b (that is, equimolar amounts of SEQ ID NOS:16/17) as sense and 30 antisense primers, respectively. NS3 and RB5a/b correspond to stretches of amino acids 101-109 and 318-326, respectively, of SEQ ID NO:2, which are conserved in most microsomal delta-12 desaturases (SEQ ID NOS:2, 4, 6, 8). PCR was carried out using the PCR 35 kit (Perkin-Elmer) using 40 cycles of 94°C 1 min, 45°C,

009201" 57E76960

1 min, and 55°C, 2 min. Analyses of the PCR products on an agarose gel showed the presence of a product of the expected size (720 bp), which was absent in control reactions containing either the sense or antisense primers alone. The PCR product fragment was gel purified and then used as a probe for screening the same corn cDNA library at 60°C as described above. One positively-hybridizing plaque was purified and partial sequence determination of its cDNA showed it to be a nucleotide sequence encoding microsomal delta-12 desaturase but truncated at the 3' end. The cDNA insert encoding the partial desaturase was gel isolated and used to probe the corn cDNA library again. Several positive plaques were recovered and characterized. DNA sequence analysis revealed that all of these clones seem to represent the same sequence with the different length of 5' or 3' ends. The clone containing the longest insert, designated pFad2#1, was sequenced completely. SEQ ID NO:7 shows the 5' to 3' nucleotide sequence of 1790 base pairs of corn (*Zea mays*) cDNA which encodes microsomal delta-12 desaturase in plasmid pFad2#1. Nucleotides 165 to 167 and nucleotides 1326 to 1328 are, respectively, the putative initiation codon and the termination codon of the open reading frame (nucleotides 164 to 1328). SEQ ID NO:8 is the 387 amino acid protein sequence deduced from the open reading frame (nucleotides 164 to 1328) in SEQ ID NO:7. The deduced amino acid sequence of the polypeptide shared overall identities of 71%, 40%, and 38% to *Arabidopsis* microsomal delta-12 desaturase, *Arabidopsis* microsomal delta-15 desaturase, and *Arabidopsis* plastid delta-15 desaturase, respectively. Furthermore, it lacked an N-terminal amino acid extension that would indicate it is a plastid enzyme. Based on these considerations, it

096973-10260

is concluded that it encodes a microsomal delta-12 desaturase.

Cloning of a cDNA Encoding A Microsomal Delta-12 Desaturase and of cDNAs Encoding Microsomal Delta-12

5 Desaturase-Related Enzymes from Castor Bean Seed

Castor microsomal delta-12 desaturase cDNA was isolated using a RT-PCR approach. Polysomal mRNA was isolated from castor beans of stages I-II (5-10 DAP) and also from castor beans of stages IV-V (20-25 DAP).

10 Ten ng of each mRNA was used for separate RT-PCR reactions, using the Perkin-Elmer RT-PCR kit with the reagent concentration as recommended by the kit protocol. The reverse transcriptase reaction was primed with random hexamers and the PCR reaction with 100 pmol

15 each of the degenerate delta-12 desaturase primers NS3 and NS9 (SEQ ID NOS:13 and 14, respectively). The reverse transcriptase reaction was incubated at 25°C for 10 min, 42°C for 15 min, 99°C for 5 min and 5°C for 5 min. The PCR reaction was incubated at 95°C for 2 min

20 followed by 35 cycles of 95°C for 1 min/50°C for 1 min. A final incubation at 60°C for 7 min completed the reaction. A DNA fragment of 720 bp was amplified from both stage I-II and stage IV-V mRNA. The amplified DNA fragment from one of the reactions was gel purified and

25 cloned into a pGEM-T vector using the Promega pGEM-T PCR cloning kit to create the plasmid pRF2-1C. The 720 bp insert in pRF2-1C was sequenced, as described above, and the resulting DNA sequence is shown in SEQ ID NO:9. The DNA sequence in SEQ ID NO:9 contains an open-reading

30 frame encoding 219 amino acids (SEQ ID NO:10), which has 81% identity (90% similarity) with amino acids 135 to 353 of the *Arabidopsis* microsomal delta-12 desaturase described in SEQ ID NO:2. The cDNA insert in pRF2-1C is therefore a 673 bp fragment of a full-length cDNA

35 encoding a castor bean seed microsomal delta-12

009201 526960

5 The insert in pRF2-1C may also be used to screen castor  
bean libraries at lower temperatures to isolate delta-12  
desaturase related sequences, such as the delta-12  
hydroxylase.

A cDNA library made to poly A<sup>+</sup> mRNA isolated from developing castor beans (stages IV-V, 20-25 DAP) was screened as described above. Radiolabeled probe prepared from pSF2b or pRF2-1C, as described above, were added, and allowed to hybridize for 18 h at 50°C. The filters were washed as described above. Autoradiography of the filters indicated that there were numerous hybridizing plaques, which appeared either strongly hybridising or weakly hybridising. Three of the strongly hybridising plaques (190A-41, 190A-42 and 190A-44) and three of the weakly hybridising plaques, (190B-41, 190b-43 and 197c-42), were plaque purified using the methods described above. The cDNA insert size of the purified phages were determined by PCR amplification of the insert using phage as template and lambda-gt11 oligomers (Clontech lambda-gt11 Amplimers) for primers. The PCR-amplified inserts of the amplified phages were subcloned into pBluescript (Pharmacia) which had been cut with Eco RI and filled in with Klenow (Sambrook et al. (Molecular Cloning, A Laboratory Approach, 2nd. ed. (1989) Cold Spring Harbor Laboratory Press). The resulting plasmids were called pRF190a-41, pRF190a-42, pRF190a-44, pRF190b-41, pRF190b-43 and pRF197c-42. All of the inserts were about 1.1 kb with the exception of pRF197c-42 which was approx. 1.5 kb. The inserts in the plasmids were sequenced as described above. The insert in pRF190b-43 did not contain any

open reading frame and was not identified. The inserts in pRF190a-41, pRF190a-42, pRF190a-44 and pRF190b-41 were identical. The insert in pRF197c-42 contained all of the nucleotides of the inserts in pRF190a-41,

5 pRF190a-42, pRF190a-44 and pRF190b-41 plus an additional approx. 400 bp. It was deduced therefore that the insert in pRF197c-42 was a longer version of the inserts in pRF190a-41, pRF190a-42, pRF190a-44 and pRF190b-41 and all were derived from the same full-length mRNA. The

10 complete cDNA sequence of the insert in plasmid pRF197c-42 is shown in SEQ ID NO:11. The deduced amino acid sequence of SEQ ID NO:11, shown in SEQ ID NO:12, is 78.5% identical (90% similarity) to the castor microsomal delta-12 desaturase described above (SEQ ID

15 NO:10) and 66% identical (80% similarity) to the Arabidopsis delta-12 desaturase amino acid sequence in SEQ ID NO:2. These similarities confirm that pRF197c-42 is a castor bean seed cDNA that encodes a microsomal delta-12 desaturase or a microsomal delta-12 desaturase-related enzyme, such as a delta-12 hydroxylase.

20 Specific PCR primers for pRF2-1C and pRF197c-42 were made. For pRF2-1c the upstream primer was bases 180 to 197 of the cDNA sequence in SEQ ID NO:9. For pRF197c-42 the upstream primer was bases 717 to 743 of the cDNA

25 sequence in SEQ ID NO:11. A common downstream primer was made corresponding to the exact complement of the nucleotides 463 to 478 of the sequence described in SEQ ID NO:9. Using RT-PCR with random hexamers and the above primers, and the incubation temperatures described

30 above, it was observed that mRNA which gave rise to the cDNA contained in pRF2-1C is present in both Stage I-II and Stage IV-V castor bean seeds whereas mRNA which gave rise to the cDNA contained in plasmid pRF197c-42 is present only in Stage IV-V castor bean seeds, i.e., it

35 is only expressed in tissue actively synthesizing

009201" 52E25950

ricinoleic acid. Thus it is possible that this cDNA encodes a delta-12 hydroxylase.

Clones such as pRF2-1C and pRF197c-42, and other clones from the differential screening, which, based on their DNA sequence, are less related to castor bean seed microsomal delta-12 desaturases and are not any of the known fatty-acid desaturases described above or in WO 9311245, may be expressed, for example, in soybean embryos or another suitable plant tissue, or in a microorganism, such as yeast, which does not normally contain ricinoleic acid, using suitable expression vectors and transformation protocols. The presence of novel ricinoleic acid in the transformed tissue(s) expressing the castor cDNA would confirm the identity of the castor cDNA as DNA encoding for an oleate hydroxylase.

#### EXAMPLE 4

#### USE OF THE ARABIDOPSIS THALIANA DELTA-12 DESATURASE GENOMIC CLONE AS A RESTRICTION FRAGMENT LENGTH POLYMORPHISM (RFLP) MARKER TO MAP THE DELTA-12 DESATURASE LOCUS IN ARABIDOPSIS

The gene encoding Arabidopsis microsomal delta-12 desaturase was used to map the genetic locus encoding the microsomal delta-12 desaturase of Arabidopsis thaliana. pSF2b cDNA insert encoding Arabidopsis microsomal delta-12 desaturase DNA was radiolabeled and used to screen an Arabidopsis genomic DNA library. DNA from several pure strongly-hybridizing phages was isolated. Southern blot analysis of the DNA from different phages using radiolabeled pSF2b cDNA insert as the probe identified a 6 kb Hind III insert fragment to contain the coding region of the gene. This fragment was subcloned in pBluescript vector to result in plasmid pAGF2-6 and used for partial sequence determination. This sequence (SEQ ID NO:15) confirmed that it is the

009221 62E/65950

microsomal delta-12 desaturase gene. DNA from two phages was isolated and labelled with  $^{32}\text{P}$  using a random priming kit from Pharmacia under conditions recommended by the manufacturer. The radioactive DNA was used to

5 probe a Southern blot containing genomic DNA from Arabidopsis thaliana (ecotype Wassileskija and marker line W100 ecotype Landesberg background) digested with one of several restriction endonucleases. Following hybridization and washes under standard conditions

10 (Sambrook et al., Molecular Cloning: A Laboratory Manual, 2nd ed. (1989) Cold Spring Harbor Laboratory Press), autoradiograms were obtained. A different pattern of hybridization (polymorphism) was identified in Hind III-digested genomic DNAs using one of the phage

15 DNAs. This polymorphism was located to a 7 kb Hind III fragment in the phage DNA that revealed the polymorphism. The 7 kb fragment was subcloned in pBluescript vector to result in plasmid pAGF2-7. Plasmid pAGF2-7 was restricted with Hind III enzyme and

20 used as a radiolabelled probe to map the polymorphism essentially as described by Helentjaris et al., (Theor. Appl. Genet. (1986) 72:761-769). The radiolabelled DNA fragment was applied as described above to Southern blots of Hind III-digested genomic DNA isolated from 117

25 recombinant inbred progeny (derived from single-seed descent lines to the  $F_6$  generation) resulting from a cross between Arabidopsis thaliana marker line W100 and ecotype Wassileskija (Burr et al., Genetics (1988) 118:519-526). The bands on the autoradiograms were

30 interpreted as resulting from inheritance of either paternal (ecotype Wassileskija) or maternal (marker line W100) DNA or both (a heterozygote). The resulting segregation data were subjected to genetic analysis using the computer program Mapmaker (Lander et al.,

35 Genomics (1987) 1:174-181). In conjunction with

09697379 102660



previously obtained segregation data for 63 anonymous RFLP markers and 9 morphological markers in Arabidopsis thaliana (Chang et al., Proc. Natl. Acad. Sci. USA (1988) 85:6856-6860; Nam et al., Plant Cell (1989) 1:699-705), a single genetic locus was positioned corresponding to the microsomal delta-12 desaturase gene. The location of the microsomal delta-12 desaturase gene was thus determined to be 13.6 cM proximal to locus c3838, 9.2 cM distal to locus 1At228, and 4.9 cM proximal to FadD locus on chromosome 3 [Koorneef, M. et. al. (1993) in Genetic Maps, Ed. O'Brien, S. J.; Yadav et al. (1993) Plant Physiology 103:467-476.]

#### EXAMPLE 5

#### 15     USE OF SOYBEAN MICROSOMAL DELTA-12 DESATURASE cDNA           SEQUENCE AS A RESTRICTION FRAGMENT           LENGTH POLYMORPHISM (RFLP) MARKER

The 1.6 kb insert obtained from the plasmid pSF2-169K as previously described was radiolabelled with <sup>32</sup>P using a Random Priming Kit from Bethesda Research Laboratories under conditions recommended by the manufacturer. The resulting radioactive probe was used to probe a Southern blot (Sambrook et al., Molecular Cloning: A Laboratory Manual, 2nd Ed. (1989) Cold Spring Harbor Laboratory Press) containing genomic DNA from soybean (Glycine max (cultivar Bonus) and Glycine soja (PI81762)) digested with one of several restriction enzymes. After hybridization and washes under low stringency conditions (50 mM Tris, pH 7.5, 6X SSPE, 10% dextran sulfate, 1% SDS at 56°C for the hybridization and initial washes, changing to 2X SSPE and 0.1% SDS for the final wash), autoradiograms were obtained, and different patterns of hybridization (polymorphisms) were identified in digests performed with restriction enzymes Hind III and Eco RI. These polymorphisms were used to

map two pSF2-169k loci relative to other loci on the soybean genome essentially as described by Helentjaris et al., (Theor. Appl. Genet. (1986) 72:761-769). The map positions of the polymorphisms were determined to be

5 in linkage group 11 between 4404.00 and 1503.00 loci (4.5 cM and 7.1 cM from 4404.00 and 1503.00, respectively) and linkage group 19 between 4010.00 and 5302.00 loci (1.9 cM and 2.7 cM from 4010.00 and 5302.00, respectively) [Rafalski, A. and Tingey, S.

10 (1993) in Genetic Maps, Ed. O' Brien, S. J.].

#### EXAMPLE 6

#### EXPRESSION OF MICROSOMAL DELTA-12 DESATURASE IN SOYBEANS

##### Construction of Vectors for Transformation of Glycine max for Reduced Expression of Microsomal Delta-12 Desaturases in Developing Soybean Seeds

Plasmids containing the antisense *G. max* microsomal delta-12 desaturase cDNA sequence under control of the soybean Kunitz Trypsin Inhibitor 3 (KTI3) promoter .

20 (Jofuku and Goldberg, Plant Cell (1989) 1:1079-1093) , the *Phaseolus vulgaris* 7S seed storage protein (phaseolin) promoter (Sengupta-Gopalan et al., Proc. Natl. Acad. Sci. USA (1985) 82:3320-3324; Hoffman et al., Plant Mol. Biol. (1988) 11:717-729) and soybean

25 beta-conglycinin promoter (Beachy et al., EMBO J. (1985) 4:3047-3053), were constructed. The construction of vectors expressing the soybean delta-12 desaturase antisense cDNA under the control of these promoters was facilitated by the use of the following plasmids:

30 pML70, pCW108 and pCW109A.

The pML70 vector contains the KTI3 promoter and the KTI3 3' untranslated region and was derived from the commercially available vector pTZ18R (Pharmacia) via the intermediate plasmids pML51, pML55, pML64 and pML65. A

35 2.4 kb Bst BI/Eco RI fragment of the complete soybean

KTi3 gene (Jofuku and Goldberg (1989) Plant Cell 1:1079-1093), which contains all 2039 nucleotides of the 5' untranslated region and 390 bases of the coding sequence of the KTi3 gene ending at the Eco RI site corresponding to bases 755 to 761 of the sequence described in Jofuku et al (1989) Plant Cell 1:427-435, was ligated into the Acc I/Eco RI sites of pTZ18R to create the plasmid pML51. The plasmid pML51 was cut with Nco I, filled in using Klenow, and religated, to destroy an Nco I site in the middle of the 5' untranslated region of the KTi3 insert, resulting in the plasmid pML55. The plasmid pML55 was partially digested with Xmn I/Eco RI to release a 0.42 kb fragment, corresponding to bases 732 to 755 of the above cited sequence, which was discarded. A synthetic Xmn I/Eco RI linker containing an Nco I site, was constructed by making a dimer of complementary synthetic oligonucleotides consisting of the coding sequence for an Xmn I site (5'-TCTTCC-3') and an Nco I site (5'-CCATGGG-3') followed directly by part of an Eco RI site (5'-GAAGG-3'). The Xmn I and Nco I/Eco RI sites were linked by a short intervening sequence (5'-ATAGCCCCCAA-3'). This synthetic linker was ligated into the Xmn I/Eco RI sites of the 4.94 kb fragment to create the plasmid pML64. The 3' untranslated region of the KTi3 gene was amplified from the sequence described in Jofuku et al (Ibid.) by standard PCR protocols (Perkin Elmer Cetus, GeneAmp PCR kit) using the primers ML51 and ML52. Primer ML51 contained the 20 nucleotides corresponding to bases 1072 to 1091 of the above cited sequence with the addition of nucleotides corresponding to Eco RV (5'-GATATC-3'), Nco I (5'-CCATGG-3'), Xba I (5'-TCTAGA-3'), Sma I (5'-CCCGGG-3') and Kpn I (5'-GGTACC-3') sites at the 5' end of the primer. Primer ML52 contained to the exact compliment of the

009201 5225960

nucleotides corresponding to bases 1242 to 1259 of the above cited sequence with the addition of nucleotides corresponding to Sma I (5'-CCCGGG-3'), Eco RI (5'-GAATTC-3'), Bam HI (5'-GGATCC-3') and Sal I (5'-GTCGAC-3') sites at the 5' end of the primer. The PCR-amplified 3' end of the KTi3 gene was ligated into the Nco I/Eco RI sites of pML64 to create the plasmid pML65. A synthetic multiple cloning site linker was constructed by making a dimer of complementary synthetic oligonucleotides consisting of the coding sequence for Pst I (5'-CTGCA-3'), Sal I (5'-GTCGAC-3'), Bam HI (5'-GGATCC-3') and Pst I (5'-CTGCA-3') sites. The linker was ligated into the Pst I site (directly 5' to the KTi3 promoter region) of pML65 to create the plasmid pML70.

The 1.46 kb Sma I/Kpn I fragment from pSF2-169K (soybean delta-12 desaturase cDNA described above) was ligated into the corresponding sites in pML70 resulting in the plasmid pBS10. The desaturase cDNA fragment was in the reverse (antisense) orientation with respect to the KTi3 promoter in pBS10. The plasmid pBS10 was digested with Bam HI and a 3.47 kb fragment, representing the KTi3 promoter/antisense desaturase cDNA/KTi3-3' end transcriptional unit was isolated by agarose gel electrophoresis. The vector pML18 consists of the non-tissue specific and constitutive cauliflower mosaic virus (35S) promoter (Odell et al., Nature (1985) 313:810-812; Hull et al., Virology (1987) 86:482-493), driving expression of the neomycin phosphotransferase gene described in (Beck et al. (1982) Gene 19:327-336) followed by the 3' end of the nopaline synthase gene including nucleotides 848 to 1550 described by (Depicker et al., (1982) J. Appl. Genet. 1:561-574). This transcriptional unit was inserted into the commercial cloning vector pGEM9Z (Gibco-BRL) and is flanked at the

009201" 52E26960

5' end of the 35S promoter by the restriction sites Sal I, Xba I, Bam HI and Sma I in that order. An additional Sal I site is present at the 3' end of the NOS 3' sequence and the Xba I, Bam HI and Sal I sites are unique. The 3.47 kb transcriptional unit released from pBS10 was ligated into the Bam HI site of the vector pML18. When the resulting plasmids were double digested with Sma I and Kpn I, plasmids containing inserts in the desired orientation yielded 3 fragments of 5.74, 2.69 and 1.46 kb. A plasmid with the transcriptional unit in the correct orientation was selected and was designated pBS13.

The pCW108 vector contains the bean phaseolin promoter and 3' untranslated region and was derived from the commercially available pUC18 plasmid (Gibco-BRL) via plasmids AS3 and pCW104. Plasmid AS3 contains 495 base pairs of the bean (*Phaseolus vulgaris*) phaseolin (7S seed storage protein) promoter starting with 5'-TGGTCTTTTGGT-3' followed by the entire 1175 base pairs of the 3' untranslated region of the same gene (see sequence descriptions in Doyle et al., (1986) J. Biol. Chem. 261:9228-9238 and Slightom et al., (1983) Proc. Natl. Acad. Sci. USA, 80:1897-1901. Further sequence description may be found in WO 9113993) cloned into the Hind III site of pUC18. The additional cloning sites of the pUC18 multiple cloning region (Eco RI, Sph I, Pst I and Sal I) were removed by digesting with Eco RI and Sal I, filling in the ends with Klenow and religating to yield the plasmid pCW104. A new multiple cloning site was created between the 495bp of the 5' phaseolin and the 1175bp of the 3' phaseolin by inserting a dimer of complementary synthetic oligonucleotides consisting of the coding sequence for a Nco I site (5'-CCATGG-3') followed by three filler bases (5'-TAG-3'), the coding sequence for a Sma I site

009201" 52625350

(5'-CCCGGG-3'), the last three bases of a Kpn I site (5'-TAC-3'), a cytosine and the coding sequence for an Xba I site (5'-TCTAGA-3') to create the plasmid pCW108. This plasmid contains unique Nco I, Sma I, Kpn I and Xba I sites directly behind the phaseolin promoter. The 1.4 kb Eco RV/Sma I fragment from pSF2-169K was ligated into the Sma I site of the commercially available phagemid pBC SK+ (Stratagene). A phagemid with the cDNA in the desired orientation was selected by digesting with Pfl MI/Xho I to yield fragments of approx. 1 kb and 4 kb and designated pM1-SF2. The 1.4 kb Xmn I/Xba I fragment from pM1-SF2 was inserted into the Sma I/Xba I sites of pCW108 to yield the plasmid pBS11, which has the soybean delta-12 desaturase cDNA in the reverse (3'-5') orientation behind the phaseolin promoter. The plasmid pBS11 was digested with Bam HI and a 3.07 kb fragment, representing the phaseolin promoter/antisense desaturase cDNA/phaseolin 3' end transcriptional unit was isolated by agarose gel electrophoresis and ligated into the Hind III site of pML18 (described above). When the resulting plasmids were digested with Xba I, plasmids containing inserts in the desired orientation yielded 2 fragments of 8.01 and 1.18 kb. A plasmid with the transcriptional unit in the correct orientation was selected and was designated pBS14.

The vector pCW109A contains the soybean b-conglycinin promoter sequence and the phaseolin 3' untranslated region and is a modified version of vector pCW109 which was derived from the commercially available plasmid pUC18 (Gibco-BRL). The vector pCW109 was made by inserting into the Hind III site of the cloning vector pUC18 a 555 bp 5' non-coding region (containing the promoter region) of the b-conglycinin gene followed by the multiple cloning sequence containing the restriction endonuclease sites for Nco I, Sma I, Kpn I

and Xba I, as described for pCW108 above, then 1174 bp of the common bean phaseolin 3' untranslated region into the Hind III site (described above). The b-conglycinin promoter region used is an allele of the published b-conglycinin gene (Doyle et al., J. Biol. Chem. (1986) 261:9228-9238) due to differences at 27 nucleotide positions. Further sequence description of this gene may be found in Slightom (W0 9113993). To facilitate use in antisense constructions, the Nco I site and potential translation start site in the plasmid pCW109 was destroyed by digestion with Nco I, mung bean exonuclease digestion and re-ligation of the blunt site to give the modified plasmid pCW109A. The plasmid pCW109A was digested with Hind III and the resulting 1.84 kb fragment, which contained the b-conglycinin/antisense delta-12 desaturase cDNA/phaseolin 3' untranslated region, was gel isolated. The plasmid pML18 (described above) was digested with Xba I, filled in using Klenow and religated, in order to remove the Xba I site. The resulting plasmid was designated pBS16. The 1.84 kb fragment of plasmid pCW109A (described above) was ligated into the Hind III site of pBS16. A plasmid containing the insert in the desired orientation yielded a 3.53 kb and 4.41 kb fragment when digested with Kpn I and this plasmid was designated pCST2. The Xmn I/Xba I fragment of pML1-SF2 (described above) was ligated into the Sma I/Xba I sites of pCST2 to yield the vector pST11.

#### 30      Transformation Of Somatic Soybean Embryo Cultures                     and Regeneration Of Soybean Plants

Soybean embryogenic suspension cultures were maintained in 35 mL liquid media (SB55 or SBP6) on a rotary shaker, 150 rpm, at 28°C with mixed florescent and incandescent lights on a 16:8 h day/night schedule. Cultures were subcultured every four weeks by

inoculating approximately 35 mg of tissue into 35 mL of liquid medium.

Soybean embryogenic suspension cultures were transformed with pCS3FdST1R by the method of particle gun bombardment (see Kline et al. (1987) Nature (London) 327:70). A DuPont Biolistic PDS1000/HE instrument (helium retrofit) was used for these transformations.

To 50 mL of a 60 mg/mL 1 mm gold particle suspension was added (in order); 5 uL DNA (1 ug/uL), 20 uL spermidine (0.1 M), and 50 uL CaCl<sub>2</sub> (2.5 M). The particle preparation was agitated for 3 min, spun in a microfuge for 10 sec and the supernatant removed. The DNA-coated particles were then washed once in 400 uL 70% ethanol and re suspended in 40 uL of anhydrous ethanol. The DNA/particle suspension was sonicated three times for 1 sec each. Five uL of the DNA-coated gold particles were then loaded on each macro carrier disk.

Approximately 300-400 mg of a four week old suspension culture was placed in an empty 60x15 mm petri dish and the residual liquid removed from the tissue with a pipette. For each transformation experiment, approximately 5-10 plates of tissue were normally bombarded. Membrane rupture pressure was set at 1000 psi and the chamber was evacuated to a vacuum of 28 inches of mercury. The tissue was placed approximately 3.5 inches away from the retaining screen and bombarded three times. Following bombardment, the tissue was placed back into liquid and cultured as described above.

Eleven days post bombardment, the liquid media was exchanged with fresh SB55 containing 50 mg/mL hygromycin. The selective media was refreshed weekly. Seven weeks post bombardment, green, transformed tissue was observed growing from untransformed, necrotic embryogenic clusters. Isolated green tissue was removed



and inoculated into individual flasks to generate new, clonally propagated, transformed embryogenic suspension cultures. Thus each new line was treated as independent transformation event. These suspensions can then be  
5 maintained as suspensions of embryos clustered in an immature developmental stage through subculture or regenerated into whole plants by maturation and germination of individual somatic embryos.

Transformed embryogenic clusters were removed from  
10 liquid culture and placed on a solid agar media (SB103) containing no hormones or antibiotics. Embryos were cultured for eight weeks at 26°C with mixed florescent and incandescent lights on a 16:8 h day/night schedule. During this period, individual embryos were removed from  
15 the clusters and analyzed at various stages of embryo development After eight weeks somatic embryos become suitable for germination. For germination, eight week old embryos were removed from the maturation medium and dried in empty petri dishes for 1 to 5 days. The dried  
20 embryos were then planted in SB71-1 medium where they were allowed to germinate under the same lighting and germination conditions described above. Germinated embryos were transferred to sterile soil and grown to maturity for seed collection.

TABLE 10

Media:		B5 Vitamin Stock
SB55 and SBP6 Stock		10 g m-inositol
Solutions		100 mg nicotinic acid
(g/L):		100 mg pyridoxine HCl
MS Sulfate 100X Stock		1 g thiamine
MgSO <sub>4</sub> 7H <sub>2</sub> O	37.0	SB55 (per Liter)
MnSO <sub>4</sub> H <sub>2</sub> O	1.69	10 mL each MS stocks
ZnSO <sub>4</sub> 7H <sub>2</sub> O	0.86	1 mL B5 Vitamin stock
CuSO <sub>4</sub> 5H <sub>2</sub> O	0.0025	0.8 g NH <sub>4</sub> NO <sub>3</sub>
MS Halides 100X Stock		3.033 g KNO <sub>3</sub>
CaCl <sub>2</sub> 2H <sub>2</sub> O	44.0	1 mL 2,4-D (10mg/mL stock)
KI	0.083	60 g sucrose
CoCl <sub>2</sub> 6H <sub>2</sub> O	0.00125	0.667 g asparagine
KH <sub>2</sub> PO <sub>4</sub>	17.0	pH 5.7
H <sub>3</sub> BO <sub>3</sub>	0.62	For SBP6- substitute 0.5 mL
Na <sub>2</sub> MoO <sub>4</sub> 2H <sub>2</sub> O	0.025	2,4-D
MS FeEDTA 100X Stock		SB103 (per Liter)
Na <sub>2</sub> EDTA	3.724	MS Salts
FeSO <sub>4</sub> 7H <sub>2</sub> O	2.784	6% maltose
		750 mg MgCl <sub>2</sub>
		0.2% Gelrite
		pH 5.7
		SB71-1 (per liter)
		B5 salts
		1ml B5 vitamin stock
		3% sucrose
		750mg MgCl <sub>2</sub>
		0.2% gelrite
		pH 5.7

009201" 62E26950

Analysis Of Transgenic Glycine Max Embryos and  
Seeds Containing An Antisense Delta-15 Desaturase:  
Demonstration That The Phenotype Of Transgenic Soybean  
Somatic Embryos Is Predictive Of The Phenotype Of Seeds  
5 Derived From Plants Regenerated From Those Embryos

While in the globular embryo state in liquid culture as described above, somatic soybean embryos contain very low amounts of triacylglycerol or storage proteins typical of maturing, zygotic soybean embryos.

10 At this developmental stage, the ratio of total triacylglyceride to total polar lipid (phospholipids and glycolipid) is about 1:4, as is typical of zygotic soybean embryos at the developmental stage from which the somatic embryo culture was initiated. At the

15 globular stage as well, the mRNAs for the prominent seed proteins (alpha' subunit of beta-conglycinin, Kunitz Trypsin Inhibitor 3 and Soybean Seed Lectin) are essentially absent. Upon transfer to hormone free media to allow differentiation to the maturing somatic embryo

20 state as described above, triacylglycerol becomes the most abundant lipid class. As well, mRNAs for alpha'-subunit of beta-conglycinin, Kunitz Trypsin Inhibitor 3 and Soybean Seed Lectin become very abundant messages in the total mRNA population. In these respects the

25 somatic soybean embryo system behaves very similarly to maturing zygotic soybean embryos *in vivo*, and is therefore a good and rapid model system for analyzing the phenotypic effects of modifying the expression of genes in the fatty acid biosynthesis pathway.

30 Furthermore, the model system is predictive of the fatty acid composition of seeds from plants derived from transgenic embryos. Liquid culture globular embryos transformed with a vector containing a soybean microsomal delta-15 desaturase, in a reverse orientation

35 and under the control of soybean conglycinin promoter

000001-6262650

(pCS3FdST 1R), gave rise to mature embryos with a reduced 18:3 content (WO 9311245). A number of embryos from line A2872 (control tissue transformed with pCST) and from lines 299/1/3, 299/15/1, 303/7/1, 306/3/1, 306/4/3, 306/4/5 (line 2872 transformed with plasmid pCS3FdST1R) were analyzed for fatty acid content. Fatty acid analysis was performed as described in WO 9311245 using single embryos as the tissue source. Mature, somatic embryos from each of these lines were also regenerated into soybean plants by transfer to regeneration medium as described above. A number of seeds taken from plants regenerated from these embryo lines were analyzed for fatty acid content. The relative fatty-acid composition of embryos taken from tissue transformed with pCS3FdST1R was compared with relative fatty-acid composition of seeds taken from plants derived from embryos transformed with pCS3FdST1R. Also, relative fatty acid compositions of embryos and seeds transformed with pCS3FdST1R were compared with control tissue, transformed with pCST. In all cases where a reduced 18:3 content was seen in a transgenic embryo line, compared with the control, a reduced 18:3 content was also observed in segregating seeds of plants derived from that line, when compared with the control seed (Table 11).

TABLE 11

Antisense Delta-15 Desaturase:  
Relative 18:3 Content Of Embryos And Seeds Of Control  
(A2172) And Transgenic (299-, 303-, 306-) Soybean Lines

Soybean Line	Embryo	Embryo	Seed	Seed
	av. %18:3	lowest %18:3	av. %18:3*	lowest %18:3
A2872 (control)	12.1 (2.6)	8.5	8.9 (0.8)	8.0
299/1/3	5.6 (1.2)	4.5	4.3 (1.6)	2.5
299/15/1	8.9 (2.2)	5.2	2.5 (1.8)	1.4

303/7/1	7.3 (1.1)	5.9	4.9 (1.9)	2.8
306/3/1	7.0 (1.9)	5.3	2.4 (1.7)	1.3
306/4/3	8.5 (1.9)	6.4	4.5 (2.2)	2.7
306/4/5	7.6 (1.6)	5.6	4.6 (1.6)	2.7

\*Seeds which were segregating with wild-type phenotype and without a copy of the transgene are not included in these averages. The number in brackets is S.D., n=10.

Thus the Applicants conclude that an altered polyunsaturated fatty acid phenotype observed in a transgenic, mature somatic embryo line is predictive of an altered fatty acid composition of seeds of plants derived from that line.

Analysis Of Transgenic Glycine Max Embryos Containing An Antisense Microsomal Delta-12 Desaturase Construct

The vectors pBS13, pBS14 and pST11 contain the soybean microsomal delta-12 desaturase cDNA, in the antisense orientation, under the control of the soybean Kunitz Trypsin Inhibitor 3 (KTI3), *Phaseolus* phaseolin, and soybean beta-conglycinin promoters as described above. Liquid culture globular embryos transformed with vectors pBS13, pBS14 and pST11, gave rise to mature embryo lines as described above. Fatty acid analysis was performed as described in WO 9311245 using single, mature embryos as the tissue source. A number of embryos from line A2872 (control tissue transformed with pCST) and from line A2872 transformed with vectors pBS13, pBS14 and pST11 were analyzed for fatty acid content. About 30% of the transformed lines showed an increased 18:1 content when compared with control lines transformed with pCST described above, demonstrating that the delta-12 desaturase had been inhibited in these lines. The remaining transformed lines showed relative fatty acid compositions similar to those of the control line. The relative 18:1 content of the lines showing an increased 18:1 content was as high as 50% compared with

a maximum of 12.5% in the control embryo lines. The average 18:1 content of embryo lines which showed an increased 18:1 content was about 35% (Table 12). In all the lines showing an increased 18:1 content there was a proportional decrease in the relative 18:2 content (Table 13).

TABLE 12

Summary Of Experiment In Which Soybean Embryos Were Transformed With Plasmids Containing A Soybean Antisense Microsomal Delta-12 Desaturase cDNA

	# of <u>Vector Lines</u>	<u># of lines</u>		
		with high <u>18:1</u>	highest <u>18:1</u>	av. (%) <u>18:1</u>
pCST (control)	---	---	12.5	10.5
pBS13	11	4	53.5	35.9
pBS14	11	2	48.7	32.6
pST11	11	3	50.1	35.9

In Table 12 the average 18:1 of transgenics is the average of all embryos transformed with a particular vector whose relative 18:1 content is greater than two standard deviations from the highest control value (12.5). The control average is the average of ten A2872 embryos (standard deviation = 1.2). The data in Table 12 are derived from Table 13 below.

005201" 62E/5950

TABLE 13

Relative Fatty Acid Contents Of Embryo Lines  
Transformed With Plasmids Containing A  
Soybean Antisense Delta-12 Desaturase cDNA

Embryo Line	Relative % Fatty-Acid Content				
A2872 (control)					
#	16:0	18:0	18:1	18:2	18:3
1	11.7	3.2	11.7	52.7	16.1
2	16.4	4.0	10.8	47.1	19.3
3	17.1	3.4	8.3	48.3	20.6
4	15.3	2.7	9.4	51.1	19.0
5	15.2	3.6	10.8	51.0	17.5
6	18.6	3.9	10.9	45.8	18.1
7	14.6	3.4	12.5	52.3	16.4
8	14.2	3.5	11.2	53.9	16.7
9	15.2	3.2	9.8	49.5	16.1
10	19.0	3.8	9.6	47.4	19.0
G335/4/197 (pBS13)					
#	16:0	18:0	18:1	18:2	18:3
1	12.2	3.3	42.0	23.0	17.4
2	12.4	2.7	22.4	39.0	21.9
3	12.0	3.2	42.0	23.2	18.4
G335/4/221 (pBS13)					
#	16:0	18:0	18:1	18:2	18:3
1	12.2	2.7	30.4	36.0	17.9
2	11.5	2.4	14.3	53.4	17.6
3	13.0	2.6	15.2	47.4	19.9
4	12.0	2.6	27.4	37.9	19.1
5	11.7	2.7	25.1	42.3	15.6
6	11.7	3.4	21.6	44.3	17.8
7	12.0	2.5	11.3	53.6	20.0
8	12.0	2.5	20.8	44.1	19.5
9	11.7	2.6	25.3	39.6	18.3

009201" 62E25950

## G335/8/174 (pBS13)

#	16:0	18:0	18:1	18:2	18:3
1	14.1	2.1	30.3	32.1	20.3
2	14.7	2.5	5.9	40.6	34.8
3	14.3	2.4	7.3	45.2	29.8

## G335/8/202 (pBS13)

#	16:0	18:0	18:1	18:2	18:3
1	11.7	1.5	30.1	32.4	23.3
2	11.4	2.3	48.5	20.6	16.1
3	12.9	2.3	46.6	17.1	19.5
4	12.7	2.6	32.0	31.1	20.5
5	12.9	1.9	41.7	23.5	18.9
6	12.3	2.6	40.1	25.6	17.9
7	11.3	2.4	53.5	16.6	14.5
8	11.4	2.5	15.5	21.7	17.8
9	10.2	2.0	45.4	23.2	18.5
10	12.8	2.2	43.2	23.5	16.9

## G335/6/42 (pBS14)

#	16:0	18:0	18:1	18:2	18:3
1	13.7	2.4	38.6	28.2	15.6
2	12.6	2.3	37.6	28.8	17.2
3	11.7	3.0	48.7	21.1	14.6

## G335/6/104 (pBS14)

#	16:0	18:0	18:1	18:2	18:3
1	13.8	2.5	30.5	35.4	16.0
2	12.3	2.3	14.6	53.2	16.4
3	12.7	2.6	27.1	36.6	20.0
4	12.6	2.2	32.1	34.9	17.4
5	12.7	2.6	23.2	41.2	19.3
6	12.6	2.2	11.7	52.5	20.1
7	13.3	2.1	23.3	41.2	18.4

## G335/1/25 (pST11)

#	16:0	18:0	18:1	18:2	18:3
1	13.7	2.8	50.7	17.5	12.1
2	14.5	3.0	41.8	23.5	15.0



3	13.9	2.9	49.1	16.8	13.6
4	12.3	2.8	47.5	19.3	14.8
G335/2/7/1 (pST11)					
#	16:0	18:0	18:1	18:2	18:3
1	15.5	4.3	21.8	38.0	17.5
2	17.8	4.1	22.0	39.5	14.0
3	15.2	3.0	20.5	42.2	16.5
G335/2/118 (pST11)					
#	16:0	18:0	18:1	18:2	18:3
1	14.1	2.7	44.7	22.6	14.0
2	15.8	2.8	37.7	26.9	14.8
3	17.3	3.4	23.3	37.9	16.0

N.B. All other transformed embryos (24 lines) had fatty acid profiles similar to those of the control.

One of these embryo lines, G335/1/25, had an average 18:2 content of less than 20% and an average 18:1 content greater than 45% (and as high as 53.5%). The Applicants expect, based on the data in Table 13, that seeds derived from plants regenerated from such lines will have an equivalent or greater increase in 18:1 content and an equivalent or greater increase decrease in 18:2 content.

Analysis Of Transgenic Glycine Max Seeds Containing An Antisense Delta-12 Desaturase Construct

The vector pST11 contains the soybean microsomal delta-12 desaturase cDNA, in the antisense orientation, under the control of the soybean beta-conglycinin promoter as described above. Liquid culture globular embryos transformed with vector pST11 gave rise to mature embryo lines as described above. These embryo were germinated, as described above, without analysis. Plants regenerated from these embryos were self-fertilized and between 15 and 30 seeds were taken from each plant. These seeds were analyzed for fatty acid

composition as described in WO 9311245. The relative fatty acid content, expressed as a percentage of the five major fatty acids are shown below in Table 13-A:

TABLE 13-A:

Relative Fatty Acids Contents Of Seeds Containing A  
Soybean Antisense Delta-12 Desaturase cDNA

LINE #	SEED #	16 0	18 0	18 1	18 2	18 3
336/3/4/1	F1	11.4	2.6	19.7	55.5	10.3
	F2	14.9	3.5	14.1	54.9	12.0
	F3	10.7	2.6	22.4	53.0	10.7
	F4	12.0	3.1	21.7	53.8	8.9
	F5	10.7	2.4	18.8	55.6	12.2
	F6	10.9	2.8	22.3	53.9	9.6
	F7	11.6	2.6	19.1	55.9	10.3
	F8	12.2	2.6	21.8	52.5	10.5
	F9	10.6	2.7	23.8	51.1	11.5
	F10	12.3	2.5	18.3	54.2	12.5
	F11	12.9	3.6	12.6	58.4	12.3
	F12	12.4	3.4	16.5	54.9	12.6
	F13	13.4	3.4	10.5	55.9	16.8
	F14	11.4	3.2	26.3	51.0	7.7
	F15	11.8	2.9	20.3	53.7	11.0
336/7/1/1	F16	13.2	2.6	25.8	43.1	14.8
	F17	9.6	3.2	65.4	11.9	9.5
	F18	10.1	2.6	41.8	37.0	8.0
	F19	8.6	2.4	60.3	18.7	9.4
	F20	11.4	3.2	41.5	30.6	12.9
	F21	8.7	2.9	63.2	16.9	8.2
	F22	10.8	3.0	39.4	35.2	11.6
	F23	7.8	2.5	79.9	3.4	6.8
	F24	9.0	2.4	63.0	17.5	8.1
	F25	9.5	2.6	60.4	19.6	8.0
	F26	8.8	3.5	67.2	11.5	8.7
	F27	10.4	3.1	53.7	22.5	10.2

009201" 62E25950

	F28	11.1	3.0	47.0	27.8	11.1
	F29	10.6	2.8	47.3	30.6	8.4
	F30	9.7	2.7	59.0	18.2	10.4
	F136	11.6	3.2	27.4	49.3	8.4
	F137	10.7	2.8	24.2	53.6	8.3
	F138	9.8	2.6	45.6	32.2	9.4
	F139	12.1	3.2	47.0	27.2	10.3
	F140	9.7	2.6	50.8	28.2	8.1
	F141	8.4	3.5	66.3	12.0	9.4
	F142	10.3	2.5	30.9	48.5	7.3
	F143	9.1	3.6	72.1	7.0	7.5
	F144	9.8	3.2	58.6	18.2	9.7
	F145	8.3	2.8	71.8	9.5	7.0
	F146	11.1	2.8	25.9	51.5	8.6
	F147	9.5	3.0	43.1	34.5	9.4
	F148	10.6	2.5	26.7	50.6	9.1
	F149	11.7	3.3	20.2	55.4	9.0
336/7/3/3	F31	11.5	2.7	20.0	55.0	10.7
	F32	12.6	3.0	24.1	52.0	8.1
	F33	11.6	2.9	20.5	56.1	8.5
	F34	13.0	3.0	20.2	53.9	10.0
	F35	11.8	2.8	20.8	53.2	11.2
	F36	10.9	3.0	29.5	48.2	8.2
	F37	11.2	3.0	26.4	49.2	10.0
	F38	12.7	3.4	15.3	56.9	11.6
	F39	11.5	3.2	19.0	54.6	11.7
	F40	10.8	3.1	23.0	52.9	10.2
	F41	11.0	3.3	32.1	45.0	8.4
	F42	13.6	3.8	13.2	55.7	13.7
	F43	10.6	2.7	25.7	50.9	10.1
	F44	9.8	2.5	25.8	49.8	12.1
	F45	12.4	3.0	19.7	53.8	11.1
336/7/4/11	F46	11.4	2.9	36.4	39.4	9.8
	F47	12.8	3.2	23.4	50.6	10.1
	F48	11.0	2.8	26.0	49.6	10.4

009207" 52E26950

	F49	11.7	2.9	22.9	52.9	9.2
	F50	10.8	2.9	32.2	46.6	7.2
	F51	11.4	3.4	20.0	54.0	11.1
	F52	11.6	3.0	26.6	49.6	9.0
	F53	12.6	2.7	31.3	43.4	9.6
	F54	10.8	3.4	23.0	51.3	11.0
	F55	11.5	2.8	22.4	54.1	8.8
	F56	11.1	2.7	25.8	50.6	9.5
	F57	11.8	3.4	20.1	54.3	10.3
	F58	13.0	3.3	21.1	51.2	11.4
	F59	12.4	3.2	19.5	53.7	10.8
	F60	9.7	2.9	41.7	37.4	8.1
336/7/5/7	F61	11.0	3.5	43.8	31.2	10.1
	F62	12.1	3.2	18.4	56.8	9.5
	F63	10.9	3.9	48.6	26.0	10.4
	F64	12.6	3.7	14.3	58.8	10.4
	F65	9.1	2.9	61.7	16.6	9.1
	F66	9.9	3.3	59.4	17.6	9.5
	F67	9.5	2.9	59.3	16.5	11.2
	F68	10.3	4.0	46.0	28.2	11.2
	F69	11.6	2.5	30.6	46.4	8.6
	F70	10.7	3.6	54.6	22.1	8.6
	F71	12.5	4.0	16.0	56.4	10.9
	F72	11.1	2.8	42.2	34.9	8.6
	F73	11.6	3.4	23.0	53.4	8.2
	F74	12.1	3.5	15.8	58.0	10.4
	F75	10.8	3.4	51.4	24.2	9.7
336/7/7/19	F76	11.4	2.7	41.7	32.8	11.5
	F77	8.6	2.5	76.6	5.1	6.8
	F78	9.5	3.2	63.2	14.6	9.1
	F79	8.2	2.8	77.7	3.8	6.8
	F80	9.6	3.4	69.8	7.9	9.2
	F81	8.3	2.8	72.5	8.6	7.0
	F82	11.1	2.8	41.5	31.9	12.3
	F83	10.2	3.0	65.7	11.2	9.4

009201" 52E25950

	F84	8.8	2.7	69.7	10.5	7.6
	F85	8.5	2.9	75.2	5.5	7.2
	F86	9.5	2.6	71.3	7.8	8.3
	F87	10.1	2.9	60.5	17.1	9.0
	F88	8.5	2.5	76.6	4.9	6.8
	F89	9.6	2.4	68.7	10.5	8.1
	F90	10.7	3.5	55.1	20.2	10.4
	F150	9.1	2.9	66.9	12.5	8.0
	F151	7.8	2.6	77.1	6.1	6.1
	F152	9.5	2.9	29.7	48.8	8.6
	F153	7.6	2.8	74.9	6.5	7.6
	F154	12.4	3.1	18.6	54.7	10.7
	F155	8.4	2.9	79.0	3.0	5.8
	F156	10.2	3.3	51.7	25.4	9.1
	F157	10.5	2.5	62.5	15.8	8.3
	F158	9.6	3.3	71.1	8.5	6.9
	F159	7.3	2.9	77.4	4.5	7.4
	F160	12.4	3.3	20.4	54.4	9.2
	F161	8.9	2.9	75.4	5.8	6.4
	F162	12.0	3.7	38.4	35.2	10.4
	F163	11.1	3.6	37.9	36.6	10.7
	F164	8.3	3.7	77.5	3.5	6.7
	F165	11.0	2.8	53.6	20.8	11.1
336/7/5/5	F91	12.8	3.7	17.3	55.3	10.6
	F92	9.1	2.8	66.3	12.8	8.5
	F93	8.9	2.2	70.5	10.4	7.4
	F94	10.5	2.5	28.1	50.9	7.6
	F95	10.9	3.2	42.8	31.9	10.8
	F96	11.4	3.4	44.9	30.5	9.5
	F97	11.7	2.7	35.8	39.5	9.8
	F98	9.7	2.5	49.0	27.7	10.9
	F99	12.3	3.5	15.9	57.1	10.8
	F100	9.4	3.0	56.7	21.2	9.3
	F101	9.8	2.6	64.4	13.4	9.6
	F102	12.3	2.8	19.6	54.4	10.7

009207 52E/5950

	F103	10.0	2.9	57.3	19.0	10.8
	F104	8.9	2.6	68.7	12.2	7.3
	F105	10.2	2.1	56.3	23.3	7.7
336/7/7/1	F106	12.3	3.2	15.7	57.9	10.6
	F107	11.2	2.7	18.8	57.4	9.7
	F108	10.9	3.3	24.3	50.1	11.2
	F109	12.2	2.5	21.0	52.5	11.5
	F110	10.8	2.6	44.5	30.4	11.5
	F111	9.8	2.9	33.4	43.9	9.8
	F112	12.5	2.7	18.0	55.7	10.7
	F113	12.4	2.8	18.4	55.2	11.2
	F114	11.0	2.4	29.3	49.4	7.5
	F115	13.0	2.9	15.2	54.8	13.9
	F116	10.6	2.6	41.1	35.4	9.8
	F117	11.4	3.1	21.4	52.7	11.1
	F118	11.2	3.2	24.7	51.3	9.4
	F119	12.0	2.5	24.3	51.4	9.3
	F120	12.2	2.8	20.5	51.8	12.3
336/7/7/3	F121	11.3	3.0	23.9	53.0	8.4
	F122	6.7	2.5	77.7	4.4	8.0
	F123	7.8	3.2	70.1	10.2	8.4
	F124	8.4	3.4	65.1	12.9	9.7
	F125	11.5	3.2	16.3	57.6	11.0
	F126	11.5	2.7	20.1	51.5	14.2
	F127	11.6	2.6	24.8	50.8	9.7
	F128	9.2	2.6	41.5	38.1	8.7
	F129	7.7	2.6	78.4	3.6	7.0
	F130	10.4	2.6	52.6	24.9	9.4
	F131	12.2	2.9	20.0	54.8	9.8
	F132	8.0	3.0	77.0	4.6	7.0
	F133	10.2	3.3	22.0	52.9	11.6
	F134	11.8	3.0	16.3	56.0	12.9
	F135	8.7	2.6	53.3	25.2	9.7
336/3/3/1	F166	11.8	3.3	25.9	49.4	9.6
	F167	13.1	3.5	15.5	54.8	13.0

009207" 5/2E/5950

	F168	12.9	2.7	22.8	51.6	9.9
	F169	12.8	3.0	17.7	55.6	10.8
	F170	13.7	3.2	15.4	55.9	11.8
	F171	12.6	3.1	17.6	54.2	12.4
	F172	12.7	3.3	17.6	54.2	12.2
	F173	11.7	3.2	15.6	56.8	12.5
	F174	12.5	3.5	17.2	56.7	9.9
	F175	12.6	3.1	22.2	51.2	10.8
	F176	12.7	3.2	17.6	54.9	11.5
	F177	12.5	3.0	22.0	52.1	10.3
	F178	12.3	2.6	32.2	43.9	9.0
	F179	12.0	3.0	17.8	56.0	11.2
	F180	12.6	3.4	18.8	55.0	10.1
	F181	11.5	2.5	27.6	48.1	10.3
	F182	14.9	3.4	14.7	49.8	17.1
	F183	10.7	2.5	24.7	53.9	8.2
	F184	14.1	3.2	13.2	54.6	14.8
	F185	13.3	3.6	16.3	54.8	11.9
336/3/4/7	F186	11.9	4.0	12.7	58.7	12.8
	F187	11.2	2.8	21.2	56.1	8.6
	F188	12.3	3.5	12.3	58.6	13.2
	F189	12.5	3.8	12.7	58.2	12.7
	F190	12.4	3.1	13.5	55.2	15.8
	F191	13.5	3.7	12.2	55.9	14.6
	F192	12.1	2.9	16.4	56.7	11.8
	F193	11.8	3.5	14.9	56.8	13.0
	F194	13.0	3.9	12.5	56.5	14.0
	F195	11.8	2.8	19.4	53.9	12.0
	F196	12.7	3.8	11.1	58.1	14.2
	F197	12.4	3.5	15.1	56.1	12.8
	F198	12.2	3.6	16.2	57.2	10.8
	F199	13.1	3.1	13.3	57.4	13.0
	F200	11.8	4.0	13.5	58.8	11.9
	F201	11.9	4.1	12.4	59.6	11.9
	F202	13.0	4.0	12.1	56.8	14.1

336/5/7/3	F203	13.0	3.6	15.0	55.6	12.8
	F204	11.8	4.0	17.9	55.3	11.0
	F205	13.5	3.9	10.8	57.0	14.7
	F206	10.2	2.5	21.3	54.3	11.6
	F207	11.6	3.5	14.4	56.6	14.0
	F208	12.4	3.4	16.6	55.1	12.4
	F209	12.0	3.0	23.9	51.4	9.6
	F210	12.8	3.0	18.0	55.3	10.8
	F211	11.2	2.7	25.6	50.3	10.0
	F212	12.8	3.2	12.1	57.6	14.3
	F213	11.2	2.8	18.2	55.4	12.3
	F214	13.8	2.9	18.6	51.2	13.5
	F215	11.8	2.2	21.0	46.4	18.6
	F216	10.9	2.5	34.2	43.7	8.6
	F217	10.0	2.6	36.9	42.8	7.7
	F218	12.0	2.9	18.1	54.5	12.4
	F219	11.8	3.2	19.8	54.2	11.0
	F220	10.3	2.5	26.7	49.9	10.6
	F221	12.8	3.3	17.7	53.2	13.0
	F222	11.7	2.7	25.6	49.6	10.4
336/7/3/9	F223	12.2	3.2	15.1	55.6	13.8
	F224	12.2	3.3	18.8	55.5	10.2
	F225	11.4	2.4	27.4	49.0	9.7
	F226	11.7	3.2	21.1	52.0	11.9
	F227	12.0	3.2	14.3	55.6	14.8
	F228	12.0	3.0	21.9	52.9	10.2
	F229	11.6	2.9	18.0	54.8	12.6
	F230	14.0	3.1	15.6	55.6	11.6
	F231	11.1	2.5	23.9	51.8	10.7
	F232	13.0	2.8	14.0	57.6	12.6
	F233	12.6	3.1	17.5	54.8	12.0
	F234	13.6	3.6	17.8	51.5	13.4
	F235	13.0	3.7	16.7	54.2	12.4
	F236	12.8	3.4	19.0	53.2	11.6
	F237	10.5	2.5	30.5	46.6	9.9

009201" 52E25950



	F238	11.5	2.8	21.5	54.5	9.5
	F239	13.8	3.0	17.1	54.6	11.6
	F240	14.3	3.9	15.3	53.8	12.7
	F241	14.2	3.4	9.9	57.9	14.5
	F242	12.5	2.9	22.8	52.1	9.7
	F243	11.9	3.3	22.4	52.4	10.0
	F244	14.5	2.5	18.4	45.0	19.6
	F245	11.3	3.2	26.5	48.4	10.6
336/7/3/13	F246	13.0	2.8	25.8	46.3	12.1
	F247	12.5	3.3	16.7	56.5	10.9
	F248	11.6	2.7	31.0	44.1	10.5
	F249	13.6	4.0	14.2	53.4	14.8
	F250	10.8	3.0	36.8	41.2	8.2
	F251	13.5	2.6	16.4	50.1	17.4
	F252	14.2	2.9	18.1	53.4	11.3
	F253	13.7	3.7	18.9	49.2	14.5
	F254	11.4	2.7	28.8	47.8	9.2
	F255	12.9	3.6	18.4	52.9	12.2
	F256	11.2	3.0	23.9	50.2	11.6
	F257	13.2	3.2	18.7	53.2	11.8
	F258	10.8	2.9	31.0	45.9	9.3
	F259	13.0	3.8	17.5	53.7	12.0
	F260	13.2	3.6	18.5	50.7	14.0
	F261	12.6	3.8	24.2	49.7	9.6
	F262	14.7	3.2	14.1	54.0	14.0
	F263	14.7	4.5	11.3	55.3	14.1
	F264	12.8	3.2	19.4	54.4	10.1
	F265	12.2	3.7	14.6	56.2	13.3
336/7/4/2	F266	12.5	3.9	46.6	28.2	8.8
	F267	9.6	2.5	61.8	18.5	7.1
	F268	10.1	2.9	62.6	13.8	9.8
	F269	12.4	3.2	25.5	49.5	9.1
	F270	8.2	3.0	72.1	9.2	6.8
	F271	14.3	3.8	14.8	57.2	9.8
	F272	12.2	3.3	17.6	49.0	17.9

009204" 62E26950

336/7/4/9	F273	11.7	3.2	41.7	34.3	8.8
	F274	11.6	3.5	44.1	30.1	10.5
	F275	13.5	3.5	22.4	48.7	11.8
	F276	12.5	3.4	23.1	48.6	12.4
	F277	13.0	3.8	22.6	49.0	11.6
	F278	12.3	3.7	27.1	46.5	10.4
	F279	10.6	3.0	58.2	19.5	8.6
	F280	12.8	4.0	13.0	59.7	10.4
	F281	12.8	3.4	17.4	56.6	9.8
	F282	10.6	4.4	46.3	29.5	9.1
	F283	13.4	4.4	23.7	47.2	11.3
	F284	11.2	3.0	29.2	46.7	9.4
	F285	11.9	3.5	36.8	38.5	8.8
	F286	9.9	2.6	39.6	37.8	10.1
	F287	8.9	2.7	57.3	23.6	7.5
	F288	10.4	2.6	45.2	33.2	8.0
	F289	10.0	2.9	30.3	46.3	10.5
	F290	10.7	2.5	34.9	42.4	9.3
	F291	9.9	2.9	52.8	25.2	8.7
	F292	10.5	2.4	39.0	38.1	9.6
	F293	9.2	2.7	33.8	43.9	10.2
	F294	11.6	2.7	29.0	46.6	9.9
	F295	12.2	2.8	18.0	56.6	10.2
	F296	10.7	2.7	17.4	56.6	12.5
	F297	10.4	2.4	27.1	41.9	7.7
	F298	10.4	2.8	19.8	58.1	8.7
	F299	10.9	2.6	18.0	57.9	10.3
	F300	9.4	2.4	40.4	39.4	7.8
	F301	9.5	2.7	45.8	33.5	8.0
	F302	10.0	2.5	40.1	38.0	9.4
	F303	9.4	2.6	45.1	33.2	9.1
	F304	10.6	2.8	22.9	51.9	11.8
	F305	10.1	2.3	34.5	45.1	7.8
336/7/6/10	F306	8.8	3.4	71.4	9.1	7.3
	F307	13.0	3.0	20.5	56.0	7.4

009204" 62E26950

336/7/7/9

F308	7.8	3.0	76.3	5.6	6.8
F309	11.1	3.1	24.4	54.2	7.1
F310	9.3	2.5	63.5	11.2	13.5
F311	9.3	3.4	70.0	9.1	8.1
F312	8.0	3.8	61.1	16.4	10.5
F313	8.6	3.3	64.1	14.5	9.5
F314	8.7	3.1	72.5	9.7	5.7
F315	9.0	3.6	58.0	20.8	8.5
F316	12.3	3.8	14.5	59.8	9.4
F317	10.1	3.9	13.6	63.6	8.8
F318	10.9	3.1	16.6	57.3	12.1
F319	8.7	2.7	74.1	7.5	6.8
F320	10.4	3.1	63.2	13.2	9.8
F321	9.6	3.2	50.4	28.2	8.6
F322	7.6	3.4	57.4	23.4	8.0
F323	9.1	2.8	66.0	14.6	7.4
F324	10.3	3.3	60.0	15.6	10.7
F325	10.1	3.5	53.7	21.8	10.7
F326	12.3	3.9	16.5	57.8	9.4
F327	8.4	3.1	72.8	7.6	7.4
F328	11.2	2.9	34.6	43.4	7.8
F329	9.5	3.3	60.8	16.6	9.8
F330	11.7	3.2	31.5	44.7	8.7
F331	12.2	2.9	22.8	53.3	8.8
F332	9.7	2.6	52.9	26.6	8.2
F333	12.2	3.3	28.7	47.6	8.2
F334	10.4	3.6	54.3	21.8	9.8
F335	10.0	3.3	64.9	11.8	10.0
F336	9.0	3.0	73.2	6.0	8.7
F337	9.2	2.7	68.8	10.1	9.2
F338	9.9	3.0	62.7	14.7	9.6
F339	8.5	3.2	72.2	6.6	9.4
F340	12.8	3.7	15.2	55.2	13.0
F341	13.2	2.8	22.5	49.8	11.7
F342	11.2	4.2	42.1	30.8	11.6

009207" 52E/6950

	F343	12.6	3.9	26.8	44.9	11.7
	F344	9.1	3.3	66.6	10.8	9.9
	F345	13.8	4.3	17.9	43.0	21.0
336/7/7/17	F346	9.9	3.2	58.2	17.5	11.2
	F347	9.1	2.8	75.0	4.8	7.8
	F348	8.8	2.8	64.1	14.0	10.2
	F349	9.7	2.9	56.4	20.0	11.0
	F350	13.9	3.9	12.8	51.1	18.2
	F351	9.2	3.3	70.0	7.8	9.6
	F352	11.3	2.5	44.4	32.9	8.9
	F353	12.2	3.0	20.6	53.8	10.4
	F354	12.0	2.8	22.1	52.8	10.4
	F355	14.1	3.5	15.2	54.2	13.1
	F356	12.7	3.0	19.7	54.2	10.4
	F357	8.0	2.4	71.4	11.2	7.0
	F358	13.0	3.4	14.3	56.1	13.1
	F359	8.3	3.0	69.4	10.3	8.8
	F360	8.7	2.6	75.8	4.7	7.8
	F361	13.1	3.3	25.4	44.8	13.3
	F362	10.6	3.6	56.0	17.2	12.6
	F363	13.9	4.0	12.6	55.5	14.0
	F364	12.9	3.8	20.7	51.0	11.4
	F365	12.7	2.6	29.0	42.7	12.8

The 18:1 content of 95 seeds taken from non-transformed A2872 soybean seeds ranged from 9.9% to 32.2% with a mean 18:1 content of 18.4%. Segregating seeds of plants derived from embryos transformed with pST11 ranged from 12.6% to 79.9% with a mean of 44.6% (Table 13-A). Seeds which had an elevated 18:1 content relative to untransformed plants also had a reduced total saturate content. For example, the total saturate content of untransformed A2872 seeds is around 15% (11% 16:0, 4% 18:0). The total saturate content of seed F122 (line 336/7/7/3, 18:1 content 77.7%) was 9.2% (6.7% 16:0, 2.5% 18:0).

EXAMPLE 7EXPRESSION OF MICROSOMAL DELTA-12 DESATURASE IN CANOLAConstruction Of Vectors For Transformation ofBrassica Napus For Reduced Expression of5       Microsomal Delta-12 Desaturasesin Developing Canola Seeds

An extended poly A tail was removed from the canola delta-12 desaturase sequence contained in plasmid pCF2-165D and additional restriction sites for cloning were introduced as follows. A PCR primer was synthesized corresponding to bases 354 through 371 of SEQ ID NO:3. The second PCR primer was synthesized as the complement to bases 1253 through 1231 with 15 additional bases (GCAGATATCGCGGCC) added to the 5' end. The additional bases encode both an EcoRV site and a NotI site. pCF2-165D was used as the template for PCR amplification using these primers. The 914 base pair product of PCR amplification was digested with EcoRV and PflMI to give an 812 base pair product corresponding to bases 450 through 1253 of pCF2-165D with the added NotI site.

pCF2-165D was digested with PstI, the PstI overhang was blunted with Klenow fragment and then digested with PflMI. The 3.5 kB fragment corresponding to pBluescript along with the 5' 450 bases of the canola Fad2 cDNA was gel purified and ligated to the above described 812 base pair fragment. The ligation product was amplified by transformation of E. coli and plasmid DNA isolation. The EcoRI site remaining at the cloning junction between pBluescript and the canola Fad2 cDNA was destroyed by digestion, blunting and religation. The recovered plasmid was called pM2CFd2.

pM2CFd2 was digested with EcoRV and SmaI to remove the Fad2 insert as a blunt ended fragment. The fragment was gel purified and cloned into the SmaI site of pBC

009201" 52E46950

(Stratagene, La Jolla, CA). A plasmid with the NotI site introduced by PCR oriented away from the existing NotI site in pBC was identified by NotI digestion and gel fractionation of the digests. The resulting  
 5 construct then had NotI sites at both ends of the canola Fad2 cDNA fragment and was called pM3CFd2.

Vectors for transformation of the antisense cytoplasmic delta-12 desaturase constructions under control of the  $\beta$ -conglycinin, Kunitz trypsin inhibitor  
 10 III, napin and phaseolin promoters into plants using Agrobacterium tumefaciens were produced by constructing a binary Ti plasmid vector system (Bevan, (1984) Nucl. Acids Res. 12:8711-8720). One starting vector for the system, (pZS199) is based on a vector which contains:  
 15 (1) the chimeric gene nopaline synthase/neomycin phosphotransferase as a selectable marker for transformed plant cells (Brevan et al. (1984) Nature 304: 184-186), (2) the left and right borders of the T-DNA of the Ti plasmid (Brevan et al. (1984) Nucl.  
 20 Acids Res. 12:8711-8720), (3) the E. coli lacZ a-complementing segment (Vieria and Messing (1982) Gene 19:259-267) with unique restriction endonuclease sites for Eco RI, Kpn I, Bam HI, and Sal I, (4) the bacterial replication origin from the Pseudomonas plasmid pVS1  
 25 (Itoh et al. (1984) Plasmid 11:206-220), and (5) the bacterial neomycin phosphotransferase gene from Tn5 (Berg et al. (1975) Proc. Natnl. Acad. Sci. U.S.A. 72:3628-3632) as a selectable marker for transformed A. tumefaciens. The nopaline synthase promoter in the  
 30 plant selectable marker was replaced by the 35S promoter (Odell et al. (1985) Nature, 313:810-813) by a standard restriction endonuclease digestion and ligation strategy. The 35S promoter is required for efficient Brassica napus transformation as described below. A  
 35 second vector (pZS212) was constructed by reversing the

order of restriction sites in the unique site cloning region of pZS199

Canola napin promoter expression cassettes were constructed as follows: Ten oligonucleotide primers were synthesized based upon the nucleotide sequence of napin lambda clone CGN1-2 published in European Patent Application EP 255378). The oligonucleotide sequences were:

- BR42 and BR43 corresponding to bases 1132 to 1156 (BR42) and the complement of bases 2248 to 2271 (BR43) of the sequence listed in Figure 2 of EP 255378.
- BR45 and BR46 corresponding to bases 1150 to 1170 (BR46) and the complement of bases 2120 to 2155 (BR45) of the sequence listed in Figure 2 of EP 255378. In addition BR46 had bases corresponding to a Sal I site (5'-GTCGAC-3') and a few additional bases (5'-TCAGGCCT-3') at its 5' end and BR45 had bases corresponding to a Bgl II site (5'-AGATCT-3') and two (5'-CT-3') additional bases at the 5' end of the primer,
- BR47 and BR48 corresponding to bases 2705 to 2723 (BR47) and bases 2643 to 2666 (BR48) of the sequence listed in Figure 2 of EP 255378. In addition BR47 had two (5'-CT-3') additional bases at the 5' end of the primer followed by bases corresponding to a Bgl II site (5'-AGATCT-3') followed by a few additional bases (5'-TCAGGCCT-3'),
- BR49 and BR50 corresponding to the complement of bases 3877 to 3897 (BR49) and the complement of bases 3985 to 3919 (BR50) of the sequence listed in Figure 2 of EP 255378. In addition BR49 had bases corresponding to a Sal I site (5'-GTCGAC-3') and a few additional bases (5'-TCAGGCCT-3') at its 5' end,
- BR57 and BR58 corresponding to the complement of bases 3875 to 3888 (BR57) and bases 2700 to 2714 (BR58) of

009201" 62E25950

the sequence listed in Figure 2 of EP 255378. In addition the 5' end of BR57 had some extra bases (5'-CCATGG-3') followed by bases corresponding to a Sac I site (5'-GAGCTC-3') followed by more additional bases (5'-GTCGACGAGG-3'). The 5' end of BR58 had additional bases (5'-GAGCTC-3') followed by bases corresponding to a Nco I site (5'-CCATGG-3') followed by additional bases (5'-AGATCTGGTACC-3').

- BR61 and BR62 corresponding to bases 1846 to 1865 (BR61) and bases 2094 to 2114 (BR62) of the sequence listed in Figure 2 of EP 255378. In addition the 5' end of BR 62 had additional bases (5'-GACA-3') followed by bases corresponding to a Bgl II site (5'-AGATCT-3') followed by a few additional bases (5'-GCGGCCGC-3').

Genomic DNA from the canola variety 'Hyola401' (Zeneca Seeds) was used as a template for PCR amplification of the napin promoter and napin terminator regions. The promoter was first amplified using primers BR42 and BR43, and reamplified using primers BR45 and BR46. Plasmid pIMC01 was derived by digestion of the 1.0 kb promoter PCR product with SalI/BglII and ligation into SalI/BamHI digested pBluescript SK<sup>+</sup> (Stratagene). The napin terminator region was amplified using primers BR48 and BR50, and reamplified using primers BR47 and BR49. Plasmid pIMC06 was derived by digestion of the 1.2 kb terminator PCR product with SalI/BglII and ligation into SalI/BglII digested pSP72 (Promega). Using pIMC06 as a template, the terminator region was reamplified by PCR using primer BR57 and primer BR58. Plasmid pIMC101 containing both the napin promoter and terminator was generated by digestion of the PCR product with SacI/NcoI and ligation into SacI/NcoI digested pIMC01. Plasmid pIMC101 contains a 2.2 kb napin expression cassette including complete napin 5' and 3'

009207 526550



non-translated sequences and an introduced NcoI site at the translation start ATG. Primer BR61 and primer BR62 were used to PCR amplify an ~270 bp fragment from the 3' end of the napin promoter. Plasmid pIMC401 was obtained  
 5 by digestion of the resultant PCR product with EcoRI/BglII and ligation into EcoRI/BglII digested pIMC101. Plasmid pIMC401 contains a 2.2 kb napin expression cassette lacking the napin 5' non-translated sequence and includes a NotI site at the transcription  
 10 start.

To construct the antisense expression vector, pM3CFd2 was digested with NotI as was pIMC401. The delta-12 desaturase containing insert from the digest of pM3CFd2 was gel isolated and ligated into the NotI  
 15 digested and phosphatase treated pIMC401. An isolate in which the delta-12 desaturase was oriented antisense to the napin promoter was selected by digestion with XhoI and PflMI to give plasmid pNCFd2R. pNCFd2R was digested with SalI, phosphatase treated and ligated into pZS212  
 20 which had been opened by the same treatment. A plasmid with desired orientation of the introduced napin:delta-12 desaturase antisense transcription unit relative to the selectable marker was chosen by digestion with PvuI and the resulting binary vector was  
 25 given the name pZNCf2R.

Plasmid pML70 (described in Example 6 above) was digested with NcoI, blunted then digested with KpnI. Plasmid pM2CFd was digested with KpnI and SmaI and the isolated fragment ligated into the opened pML70 to give  
 30 the antisense expression cassette pMKCFd2R. The promotor:delta-12 desaturase:terminator sequence was removed from pMKCFd2R by BamHI digestion and ligated into pZS199 which had been BamHI digested and phosphatase treated. The desired orientation relative  
 35 to the selectable marker was determined by digestion

with XhoI and PflMI to give the expression vector pZKCFd2R.

The expression vector containing the  $\beta$ -conglitin promoter was constructed by SmaI and EcoRV digestion of pM2CFd2 and ligation into SmaI cut pML109A. An isolate with the antisense orientation was identified by digestion with XhoI and PflMI, and the transcription unit was isolated by SalI and EcoRI digestion. The isolated SalI-EcoRI fragment was ligated into EcoRI-SalI digested pZS199 to give pCCFd2R.

The expression vector containing the phaseolin promoter was obtained using the same procedure with pCW108 as the starting, promoter containing vector and pZS212 as the binary portion of the vector to give pZPhCFd2R.

#### Agrobacterium-Mediated Transformation Of Brassica Napus

The binary vectors pZNCf2R, pZCCf2R, pZPhCFd2R, and pZNCf2R were transferred by a freeze/thaw method (Holsters et al. (1978) Mol Gen Genet 163:181-187) to the Agrobacterium strain LBA4404/pAL4404 (Hockema et al. (1983), Nature 303:179-180).

Brassica napus cultivar "Westar" was transformed by co-cultivation of seedling pieces with disarmed Agrobacterium tumefaciens strain LBA4404 carrying the the appropriate binary vector.

B. napus seeds were sterilized by stirring in 10% Chlorox, 0.1% SDS for thirty min, and then rinsed thoroughly with sterile distilled water. The seeds were germinated on sterile medium containing 30 mM CaCl<sub>2</sub> and 1.5% agar, and grown for six days in the dark at 24°C.

Liquid cultures of Agrobacterium for plant transformation were grown overnight at 28°C in Minimal A medium containing 100 mg/L kanamycin. The bacterial cells were pelleted by centrifugation and resuspended at

a concentration of  $10^8$  cells/mL in liquid Murashige and Skoog Minimal Organic medium containing 100  $\mu$ M acetosyringone.

5 B. napus seedling hypocotyls were cut into 5 mm segments which were immediately placed into the bacterial suspension. After 30 min, the hypocotyl pieces were removed from the bacterial suspension and placed onto BC-35 callus medium containing 100  $\mu$ M acetosyringone. The plant tissue and Agrobacteria were  
10 co-cultivated for three days at 24°C in dim light.

The co-cultivation was terminated by transferring the hypocotyl pieces to BC-35 callus medium containing 200 mg/L carbenicillin to kill the Agrobacteria, and 25 mg/L kanamycin to select for transformed plant cell  
15 growth. The seedling pieces were incubated on this medium for three weeks at 28°C under continuous light.

After four weeks, the segments were transferred to BS-48 regeneration medium containing 200 mg/L carbenicillin and 25 mg/L kanamycin. Plant tissue was  
20 subcultured every two weeks onto fresh selective regeneration medium, under the same culture conditions described for the callus medium. Putatively transformed calli grew rapidly on regeneration medium; as calli reached a diameter of about 2 mm, they were removed from  
25 the hypocotyl pieces and placed on the same medium lacking kanamycin.

Shoots began to appear within several weeks after transfer to BS-48 regeneration medium. As soon as the shoots formed discernable stems, they were excised from  
30 the calli, transferred to MSV-1A elongation medium, and moved to a 16:8 h photoperiod at 24°C.

Once shoots had elongated several internodes, they were cut above the agar surface and the cut ends were dipped in Rootone. Treated shoots were planted directly  
35 into wet Metro-Mix 350 soilless potting medium. The pots

were covered with plastic bags which were removed when the plants were clearly growing -- after about ten days.

Plants were grown under a 16:8 h photoperiod, with a daytime temperature of 23°C and a nighttime temperature of 17°C. When the primary flowering stem began to elongate, it was covered with a mesh pollen-containment bag to prevent outcrossing. Self-pollination was facilitated by shaking the plants several times each day. Fifty-one plants have thus far been obtained from transformations using both pZCCFd2R and pZPhCFd2R, 40 plants have been obtained from pZKCFd2R and 26 from pZNCFd2R.

#### Minimal A Bacterial Growth Medium

Dissolve in distilled water:

15      10.5 grams potassium phosphate, dibasic  
          4.5 grams potassium phosphate, monobasic  
          1.0 gram ammonium sulfate  
          0.5 gram sodium citrate, dihydrate  
          Make up to 979 mL with distilled water

20      Autoclave  
          Add 20 mL filter-sterilized 10% sucrose  
          Add 1 mL filter-sterilized 1 M MgSO<sub>4</sub>

#### Brassica Callus Medium BC-35

Per liter:

25      Murashige and Skoog Minimal Organic Medium (MS salts, 100 mg/L i-inositol, 0.4 mg/L thiamine; GIBCO #510-3118)  
          30 grams sucrose  
          18 grams mannitol

30      0.5 mg/L 2,4-D  
          0.3 mg/L kinetin  
          0.6% agarose  
          pH 5.8

#### Brassica Regeneration Medium BS-48

35      Murashige and Skoog Minimal Organic Medium

009201" 52E25950

Gamborg B5 Vitamins (SIGMA #1019)

10 grams glucose

250 mg xylose

600 mg MES

5 0.4% agarose

pH 5.7

Filter-sterilize and add after autoclaving:

2.0 mg/L zeatin

0.1 mg/L IAA

10 Brassica Shoot Elongation Medium MSV-1A

Murashige and Skoog Minimal Organic Medium

Gamborg B5 Vitamins

10 grams sucrose

0.6% agarose

15 pH 5.8

Analysis Of Transgenic Brassica Napus Seeds Containing  
An Antisense Microsomal Delta-12 Desaturase Construct

Fifty-one plants were obtained from transformation with both pZPhCFd2R and pZCCFd2R, 40 were obtained from pZKCFd2R, and 26 from pZNCFd2R. The relative levels of oleate (18:1), linoleate (18:2) and linolate (18:3) change during development so that reliable determination of seed fatty acid phenotype is best obtained from seed which has undergone normal maturation and drydown.

25 Relatively few transformed plants have gone through to maturity, however seeds were sampled from plants which had been transferred to pots for at least 80 days and which had pods that had yellowed and contained seeds with seed coats which had black pigmentation. Plants were chosen for early analysis based on promoter type, presence and copy number of the inserted delta-12 desaturase antisense gene and fertility of the plant.

35 Fatty acid analysis was done on either individual seeds from transformed and control plants, or on 40 mg of bulk seed from individual plants as described in

009207" 62E26960

Example 6. Southern analysis for detection of the presence of canola delta-12 desaturase antisense genes was done on DNA obtained from leaves of transformed plants. DNA was digested either to release the  
 5 promotor:delta-12 desaturase fragment from the transformation vector or to cut outside the coding region of the delta-12 desaturase antisense gene, but within the left and right T-DNA borders of the vector.

TABLE 14

Relative Fatty Acid Profiles of Microsomal Delta-12 Desaturase  
Antisense Transformed and Control Brassica Napus Seeds

PLANT #	PROMOTER	COPY#	AGE*	% of TOTAL FATTY ACIDS				
				16:0	18:0	18:1	18:2	18:3
Westar	control	none	82	4.6	1.2	64.6	20.9	6.6
151-22	phaseolin	>8	82	4.4	1.0	76.6	10.0	6.2
158-8	napin	1	83	3.5	1.5	81.3	6.3	4.6
westar	control	none	106	4.1	1.7	64.4	19.9	7.1
151-22	phaseolin	>8	106	4.2	1.9	74.4	9.9	6.3
151-127	phaseolin	0	106	4.1	2.3	68.4	16.9	5.2
151-268	phaseolin	1	106	4.2	2.7	73.3	12.0	4.2
153-83	conglycinin	2	106	4.1	1.6	68.5	16.7	6.3

\*Seed sampling date in days after the plant was transferred to soil

10

The expected fatty acid phenotype for antisense suppression of the delta-12 desaturase is decreased relative content of 18:2 with a corresponding increase in 18:1. Plant numbers 151-22 and 158-8 both show a  
 15 substantial decrease in 18:2 content of bulk seed when compared to the westar control at 83 days after planting. Plant 151-22 also shows this difference at maturity in comparison to either the westar control or plant 151-127, which was transformed with the selectable  
 20 marker gene but not the delta-12 desaturase antisense gene.

Since the fatty acid analysis was done on seeds from the primary transformant, individual seed should be segregating for the presense of the transgene copy or copies. The segregating phenotypes serve as an internal control for the effect of the delta-12 desaturase antisense gene. The relative fatty acid phenotypes for 10 individual westar seeds, 10 individual 151-22 seeds and 12 individual 158-8 seeds are given in Table 15 below.

10

TABLE 15

Relative Fatty Acid Profiles for Individual Seeds of Control and Genetically Segregating Delta-12 Desaturase Transformed Brassica Napus Seeds

<u>westar control</u>				
<u>16:0</u>	<u>18:0</u>	<u>18:1</u>	<u>18:2</u>	<u>18:3</u>
4.65	1.05	63.45	21.31	7.29
4.65	1.37	65.41	20.72	6.18
3.86	1.31	62.19	22.50	8.18
4.46	1.41	66.81	19.40	5.63
4.76	1.30	61.90	22.39	7.65
4.59	1.10	64.77	20.62	6.56
4.61	1.16	68.66	18.20	5.07
4.71	1.26	67.28	19.32	5.18
4.67	0.98	61.96	22.93	7.61
4.73	1.33	63.85	21.65	6.23
<u>151-22</u>				
<u>16:0</u>	<u>18:0</u>	<u>18:1</u>	<u>18:2</u>	<u>18:3</u>
4.56	1.08	73.40	12.40	7.60
4.25	1.20	77.90	10.00	5.40
4.40	1.00	76.90	10.10	5.90
4.40	0.94	77.40	9.40	6.10
4.50	1.00	73.60	11.30	7.90
4.60	0.98	75.40	10.50	6.50
4.49	0.96	76.70	9.90	6.00
4.20	1.10	77.20	9.70	5.50

00920T" 6226950

4.20	1.00	80.00	7.90	4.90
4.50	1.00	78.00	8.80	5.80
<u>158-8</u>				
<u>16:0</u>	<u>18:0</u>	<u>18:1</u>	<u>18:2</u>	<u>18:3</u>
3.62	1.67	84.45	3.60	3.73
3.46	1.64	85.56	3.02	3.36
3.48	1.61	83.64	4.43	4.21
3.53	1.40	83.80	4.41	4.36
3.48	1.39	83.66	4.35	4.44
3.80	1.50	68.17	16.57	7.56
3.41	1.40	83.76	4.38	4.40
3.49	1.29	82.77	5.16	4.60
3.77	1.39	69.47	16.40	6.54
3.44	1.36	83.86	4.49	4.27
3.48	1.38	83.15	4.91	4.53
3.55	1.92	83.69	4.20	3.70

The westar control shows comparatively little seed  
 to seed variation in content of 18:1 or 18:2. Further  
 the ratio of 18:3/18:2 remains very constant between  
 5 seeds at about 0.35. Plant #158-8 should show a  
 segregation ratio of either 1:2:1 or 1:3 since by  
 Southern analysis it contains a single transgene. The  
 1:2:1 ratio would indicate a semi-dominant, copy number  
 effect while the 1:3 ratio would indicate complete  
 10 dominance. Two wild type 158-8 segregants are clear in  
 Table 15, while the remaining seeds may either be the  
 same, or the two seeds at greater than 84% 18:1 may  
 represent the homozygous transgeneic. In either case  
 the fatty acid phenotypes of the seeds are as expected  
 15 for effective delta-12 desaturase suppression in this  
 generation. The fatty acid phenotypes of the seeds of  
 plant 151-22 show variation in their 18:1 and 18:2  
 content, with 18:1 higher than the control average and  
 18:2 lower. The segregation is apparently quite



complex, as would be expected of a multi-copy transgenic plant.

#### EXAMPLE 8

##### COMBINATION OF HIGH OLEIC TRAITS IN

##### TRANSFORMANT 158-8-1 AND RAPESEED MUTANT IMC 129

*Brassica napus* primary transformant 158-8 that was transformed with a chimeric napin:antisense rapeseed delta-12 desaturase gene (see Example 7 herein) was selfed to give rise to seeds, designated T2 generation, that were segregating for high oleic acid phenotypes. Several high oleic T2 seeds were germinated and leaf DNA of the T2 plants was used in quantitative Southern analysis to identify individuals homozygous for the transformed chimeric gene. One homozygous T2 plant, designated 158-8-1, was selfed and the fatty acid composition of the T3 progeny determined by single seed analysis (Table 16). The data confirmed that 158-8-1 was homozygous for the high oleate trait.

TABLE 16

Fatty Acid Composition of Single Seeds of a Selfed T2 Transgenic Rapeseed Line 158-8-1

Progeny	Relative Fatty Acid Content (%)				
	16:0	18:0	18:1	18:2	18:3
1	3.91	1.59	82.36	5.75	5.78
2	3.81	1.53	83.34	5.55	5.16
3	3.82	1.55	85.29	4.47	4.27
4	3.67	1.75	83.21	5.05	5.72
5	3.74	1.79	82.64	5.29	5.86
6	3.92	1.63	83.43	5.51	4.91
7	3.82	1.47	84.10	5.10	4.89
8	3.83	1.67	82.63	5.97	5.34
Average	3.82	1.62	83.38	5.34	5.24

T2 plant 158-8-1 was also used as the male parent in a cross with either rapeseed cultivar Westar (control) or IMC 129. Three plants were grown from the F1 hybrid seed from each cross and selfed. Single, segregating F2 seeds from each cross were analyzed for fatty acid composition by imbibing the seeds in water overnight, removing the seed coat and cutting off approximately one-half of the outer cotyledon. Fatty acids in the partial cotyledon were analyzed by direct transesterification in methanol and gas chromatography as described in Example 7.

The combined analysis from 272 F2 seeds from the three 158-8-1 X IMC 129 crosses and of 60 F2 seeds from two 158-8-1 X Westar crosses are shown in Table 17 as % individuals in each class of 18:1 content.

TABLE 17

Frequency Distribution of F2 seeds of 158-8-1 X IMC 129  
Cross on the Basis of Oleic Acid Content

% 18:1 class	% seeds from 158-8-1 X Westar cross	% of seeds from 158-8-1 X IMC129 cross
61-63	0	0.4
64-66	16.7	1.1
67-69	5	1.1
70-72	3.3	5.1
73-75	0	8.8
76-78	13.3	7.0
79-81	45	7.7
82-84	15	22.1
85-87	1.7	36.8
88-90	0	9.9

The data shows that the average 18:1 content in F2 progeny of the cross involving IMC 129 was significantly higher than that involving Westar.

Thirty F2 half-seeds selected for their very high oleic acid content were germinated and their leaf DNA subjected to molecular analysis (Southern analysis and polymerase chain reaction) with gene specific probes to identify 6 F2 plants that were homozygous for both the mutant delta-12 desaturase gene from IMC 129 and the transgene from 158-8-1. The fatty acid composition of the F2 seeds from which these 6 F2 plants were derived is shown in Table 18 and their average fatty acid compositions is compared to those of 158-8-1 T3 seeds (Table 16) and to those of Westar control seeds (Table 19).

TABLE 18

Fatty Acid Composition of Six F2 Seeds Homozygous for the Transgene and High Oleic Acid Mutation in IMC 129

Progeny	Relative Fatty Acid Content (%)				
	16:0	18:0	18:1	18:2	18:3
1	2.7	1.9	89.0	2.1	3.5
2	3.0	1.4	89.2	2.4	3.6
3	3.0	1.3	88.2	3.0	3.4
4	2.6	1.3	90.9	2.4	2.9
5	3.3	1.1	84.4	4.6	6.3
6	3.2	1.3	88.4	2.65	4.4
Average	2.97	1.38	88.4	2.86	4.0

TABLE 19

Seed Fatty Acid Composition of Westar, 158-8-1 and Six Seeds Homozygous for the Transgene from 158-8-1 and High Oleic Acid Mutation from IMC 129.

Progeny	Relative Fatty Acid Content (%)				
	16:0	18:0	18:1	18:2	18:3
Westar	4.20	2.20	63.00	19.90	10.60
158-8-1	3.33	1.63	83.38	5.34	5.24
158-8-1 x IMC129	2.97	1.38	88.40	2.86	4.02

The data in Table 19 shows that the the seed oleic acid content in the double homozygotes resulting from the 158-8-1 X IMC 129 cross are higher than either parent and that this increase is associated with reduced  
5 saturates (16:0+18:0). Furthermore, the double homozygote plants appear normal in their growth and development.

009207" 62E26960

SEQUENCE LISTING

## (1) GENERAL INFORMATION:

- (i) APPLICANT: JONATHAN EDWARD LIGHTNER  
JOHN JOSEPH OKULEY
- (ii) TITLE OF INVENTION: GENES FOR MICROSOMAL  
FATTY ACID DELTA-12  
DESATURASES AND RELATED  
ENZYMES FROM PLANTS
- (iii) NUMBER OF SEQUENCES: 17
- (iv) CORRESPONDENCE ADDRESS:
  - (A) ADDRESSEE: E. I. DU PONT DE NEMOURS  
AND COMPANY
  - (B) STREET: 1007 MARKET STREET
  - (C) CITY: WILMINGTON
  - (D) STATE: DELAWARE
  - (E) COUNTRY: U.S.A.
  - (F) ZIP: 19898
- (v) COMPUTER READABLE FORM:
  - (A) MEDIUM TYPE: Floppy disk
  - (B) COMPUTER: MacIntosh
  - (C) OPERATING SYSTEM: MacIntosh 6.0
  - (D) SOFTWARE: Microsoft Word 4.0
- (vi) CURRENT APPLICATION DATA:
  - (A) APPLICATION NUMBER:
  - (B) FILING DATE:
  - (C) CLASSIFICATION:
- (vii) PRIOR APPLICATION DATA:
  - (A) APPLICATION NUMBER: U.S. 07/977,339
  - (B) FILING DATE: 17-NOV-1992
- (viii) ATTORNEY/AGENT INFORMATION:
  - (A) NAME: Siegell, Barbara C.
  - (B) REGISTRATION NUMBER: 30,684
  - (C) REFERENCE/DOCKET NUMBER: BB-1043-B
- (ix) TELECOMMUNICATION INFORMATION:
  - (A) TELEPHONE: (302) 992-4927
  - (B) TELEFAX: (302) 892-7949
  - (C) TELEX: 835420

009207 625950

## (2) INFORMATION FOR SEQ ID NO:1:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 1372 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: double  
 (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: cDNA
- (iii) HYPOTHETICAL: NO
- (iv) ANTI-SENSE: NO
- (vi) ORIGINAL SOURCE:  
 (A) ORGANISM: *Arabidopsis thaliana*
- (vii) IMMEDIATE SOURCE:  
 (B) CLONE: p92103
- (ix) FEATURE:  
 (A) NAME/KEY: CDS  
 (B) LOCATION: 93..1244
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

```

AGAGAGAGAG ATTCTGCGGA GGAGCTTCTT CTTCGTAGGG TGTCATCGT TATTAACGTT      60
ATCGCCCCCTA CGTCAGCTCC ATCTCCAGAA AC ATG GGT GCA GGT GGA AGA ATG      113
                               Met Gly Ala Gly Gly Arg Met
                               1                               5
CCG GTT CCT ACT TCT TCC AAG AAA TCG GAA ACC GAC ACC ACA AAG CGT      161
Pro Val Pro Thr Ser Ser Lys Lys Ser Glu Thr Asp Thr Thr Lys Arg
          10                      15                      20
GTG CCG TGC GAG AAA CCG CCT TTC TCG GTG GGA GAT CTG AAG AAA GCA      209
Val Pro Cys Glu Lys Pro Pro Phe Ser Val Gly Asp Leu Lys Lys Ala
          25                      30                      35
ATC CCG CCG CAT TGT TTC AAA CGC TCA ATC CCT CGC TCT TTC TCC TAC      257
Ile Pro Pro His Cys Phe Lys Arg Ser Ile Pro Arg Ser Phe Ser Tyr
          40                      45                      50
CTT ATC AGT GAC ATC ATT ATA GCC TCA TGC TTC TAC TAC GTC GCC ACC      305
Leu Ile Ser Asp Ile Ile Ile Ala Ser Cys Phe Tyr Tyr Val Ala Thr
          60                      65                      70
AAT TAC TTC TCT CTC CTC CCT CAG CCT CTC TCT TAC TTG GCT TGG CCA      353
Asn Tyr Phe Ser Leu Leu Pro Gln Pro Leu Ser Tyr Leu Ala Trp Pro
          75                      80                      85
CTC TAT TGG GCC TGT CAA GGC TGT GTC CTA ACT GGT ATC TGG GTC ATA      401
Leu Tyr Trp Ala Cys Gln Gly Cys Val Leu Thr Gly Ile Trp Val Ile
          90                      95                      100

```

009201" 6226950

GCC Ala	CAC His	GAA Glu	TGC Cys	GGT Gly	CAC His	CAC His	GCA Ala	TTC Phe	AGC Ser	GAC Asp	TAC Tyr	CAA Gln	TGG Trp	CTG Leu	GAT Asp	449
105					110					115						
GAC Asp	ACA Thr	GTT Val	GGT Gly	CTT Leu	ATC Ile	TTC Phe	CAT His	TCC Ser	TTC Phe	CTC Leu	CTC Leu	GTC Val	CCT Pro	TAC Tyr	TTC Phe	497
120					125					130					135	
TCC Ser	TGG Trp	AAG Lys	TAT Tyr	AGT Ser	CAT His	CGC Arg	CGT Arg	CAC His	CAT His	TCC Ser	AAC Asn	ACT Thr	GGA Gly	TCC Ser	CTC Leu	545
				140					145					150		
GAA Glu	AGA Arg	GAT Asp	GAA Glu	GTA Val	TTT Phe	GTC Val	CCA Pro	AAG Lys	CAG Gln	AAA Lys	TCA Ser	GCA Ala	ATC Ile	AAG Lys	TGG Trp	593
			155					160					165			
TAC Tyr	GGG Gly	AAA Lys	TAC Tyr	CTC Leu	AAC Asn	AAC Asn	CCT Pro	CTT Leu	GGA Gly	CGC Arg	ATC Ile	ATG Met	ATG Met	TTA Leu	ACC Thr	641
			170				175					180				
GTC Val	CAG Gln	TTT Phe	GTC Val	CTC Leu	GGG Gly	TGG Trp	CCC Pro	TTG Leu	TAC Tyr	TTA Leu	GCC Ala	TTT Phe	AAC Asn	GTC Val	TCT Ser	689
	185					190					195					
GGC Gly	AGA Arg	CCG Pro	TAT Tyr	GAC Asp	GGG Gly	TTC Phe	GCT Ala	TGC Cys	CAT His	TTC Phe	TTC Phe	CCC Pro	AAC Asn	GCT Ala	CCC Pro	737
200					205					210					215	
ATC Ile	TAC Tyr	AAT Asn	GAC Asp	CGA Arg	GAA Glu	CGC Arg	CTC Leu	CAG Gln	ATA Ile	TAC Tyr	CTC Leu	TCT Ser	GAT Asp	GCG Ala	GGT Gly	785
				220					225					230		
ATT Ile	CTA Leu	GCC Ala	GTC Val	TGT Cys	TTT Phe	GGT Gly	CTT Leu	TAC Tyr	CGT Arg	TAC Tyr	GCT Ala	GCT Ala	GCA Ala	CAA Gln	GGG Gly	833
			235					240					245			
ATG Met	GCC Ala	TCG Ser	ATG Met	ATC Ile	TGC Cys	CTC Leu	TAC Tyr	GGA Gly	GTA Val	CCG Pro	CTT Leu	CTG Leu	ATA Ile	GTG Val	AAT Asn	881
		250				255						260				
GCG Ala	TTC Phe	CTC Leu	GTC Val	TTG Leu	ATC Ile	ACT Thr	TAC Tyr	TTG Leu	CAG Gln	CAC His	ACT Thr	CAT His	CCC Pro	TCG Ser	TTG Leu	929
	265					270					275					
CCT Pro	CAC His	TAC Tyr	GAT Asp	TCA Ser	TCA Ser	GAG Glu	TGG Trp	GAC Asp	TGG Trp	CTC Leu	AGG Arg	GGA Gly	GCT Ala	TTG Leu	GCT Ala	977
280					285					290					295	
ACC Thr	GTA Val	GAC Asp	AGA Arg	GAC Asp	TAC Tyr	GGA Gly	ATC Ile	TTG Leu	AAC Asn	AAG Lys	GTG Val	TTC Phe	CAC His	AAC Asn	ATT Ile	1025
				300					305					310		
ACA Thr	GAC Asp	ACA Thr	CAC His	GTG Val	GCT Ala	CAT His	CAC His	CTG Leu	TTC Phe	TCG Ser	ACA Thr	ATG Met	CCG Pro	CAT His	TAT Tyr	1073
			315					320					325			

AAC GCA ATG GAA GCT ACA AAG GCG ATA AAG CCA ATT CTG GGA GAC TAT 1121  
 Asn Ala Met Glu Ala Thr Lys Ala Ile Lys Pro Ile Leu Gly Asp Tyr  
 330 335 340  
 TAC CAG TTC GAT GGA ACA CCG TGG TAT GTA GCG ATG TAT AGG GAG GCA 1169  
 Tyr Gln Phe Asp Gly Thr Pro Trp Tyr Val Ala Met Tyr Arg Glu Ala  
 345 350 355  
 AAG GAG TGT ATC TAT GTA GAA CCG GAC AGG GAA GGT GAC AAG AAA GGT 1217  
 Lys Glu Cys Ile Tyr Val Glu Pro Asp Arg Glu Gly Asp Lys Lys Gly  
 360 365 370 375  
 GTG TAC TGG TAC AAC AAT AAG TTA TGAGCATGAT GGTGAAGAAA TTGTCGACCT 1271  
 Val Tyr Trp Tyr Asn Asn Lys Leu  
 380  
 TTCTCTTGTC TGTTTGTCTT TTGTTAAAGA AGCTATGCTT CGTTTAAATA ATCTTATTGT 1331  
 CCATTTTGTG GTGTTATGAC ATTTTGGCTG CTCATTATGT T 1372

## (2) INFORMATION FOR SEQ ID NO:2:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 383 amino acids  
 (B) TYPE: amino acid  
 (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Met Gly Ala Gly Gly Arg Met Pro Val Pro Thr Ser Ser Lys Lys Ser  
 1 5 10 15  
 Glu Thr Asp Thr Thr Lys Arg Val Pro Cys Glu Lys Pro Pro Phe Ser  
 20 25 30  
 Val Gly Asp Leu Lys Lys Ala Ile Pro Pro His Cys Phe Lys Arg Ser  
 35 40 45  
 Ile Pro Arg Ser Phe Ser Tyr Leu Ile Ser Asp Ile Ile Ile Ala Ser  
 50 55 60  
 Cys Phe Tyr Tyr Val Ala Thr Asn Tyr Phe Ser Leu Leu Pro Gln Pro  
 65 70 75 80  
 Leu Ser Tyr Leu Ala Trp Pro Leu Tyr Trp Ala Cys Gln Gly Cys Val  
 85 90 95  
 Leu Thr Gly Ile Trp Val Ile Ala His Glu Cys Gly His His Ala Phe  
 100 105 110  
 Ser Asp Tyr Gln Trp Leu Asp Asp Thr Val Gly Leu Ile Phe His Ser  
 115 120 125  
 Phe Leu Leu Val Pro Tyr Phe Ser Trp Lys Tyr Ser His Arg Arg His  
 130 135 140

009201" 52E25950



His Ser Asn Thr Gly Ser Leu Glu Arg Asp Glu Val Phe Val Pro Lys  
 145 150 155 160  
 Gln Lys Ser Ala Ile Lys Trp Tyr Gly Lys Tyr Leu Asn Asn Pro Leu  
 165 170 175  
 Gly Arg Ile Met Met Leu Thr Val Gln Phe Val Leu Gly Trp Pro Leu  
 180 185 190  
 Tyr Leu Ala Phe Asn Val Ser Gly Arg Pro Tyr Asp Gly Phe Ala Cys  
 195 200 205  
 His Phe Phe Pro Asn Ala Pro Ile Tyr Asn Asp Arg Glu Arg Leu Gln  
 210 215 220  
 Ile Tyr Leu Ser Asp Ala Gly Ile Leu Ala Val Cys Phe Gly Leu Tyr  
 225 230 235 240  
 Arg Tyr Ala Ala Ala Gln Gly Met Ala Ser Met Ile Cys Leu Tyr Gly  
 245 250 255  
 Val Pro Leu Leu Ile Val Asn Ala Phe Leu Val Leu Ile Thr Tyr Leu  
 260 265 270  
 Gln His Thr His Pro Ser Leu Pro His Tyr Asp Ser Ser Glu Trp Asp  
 275 280 285  
 Trp Leu Arg Gly Ala Leu Ala Thr Val Asp Arg Asp Tyr Gly Ile Leu  
 290 295 300  
 Asn Lys Val Phe His Asn Ile Thr Asp Thr His Val Ala His His Leu  
 305 310 315 320  
 Phe Ser Thr Met Pro His Tyr Asn Ala Met Glu Ala Thr Lys Ala Ile  
 325 330 335  
 Lys Pro Ile Leu Gly Asp Tyr Tyr Gln Phe Asp Gly Thr Pro Trp Tyr  
 340 345 350  
 Val Ala Met Tyr Arg Glu Ala Lys Glu Cys Ile Tyr Val Glu Pro Asp  
 355 360 365  
 Arg Glu Gly Asp Lys Lys Gly Val Tyr Trp Tyr Asn Asn Lys Leu  
 370 375 380

## (2) INFORMATION FOR SEQ ID NO:3:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1394 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: cDNA

009207 526950

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Brassica napus

(vii) IMMEDIATE SOURCE:

(B) CLONE: pCF2-165D

(ix) FEATURE:

(A) NAME/KEY: CDS

(B) LOCATION: 99..1250

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

GAGAGGAGAC AGAGACAGAG AGAGAGTTGA GAGAGCTCTC GTAGGTTATC GTATTAACGT	60
AATCTTCAAT CCCCCCTACG TCAGCCAGCT CAAGAAAC ATG GGT GCA GGT GGA	113
Met Gly Ala Gly Gly	
1 5	
AGA ATG CAA GTG TCT CCT CCC TCC AAA AAG TCT GAA ACC GAC AAC ATC	161
Arg Met Gln Val Ser Pro Pro Ser Lys Lys Ser Glu Thr Asp Asn Ile	
10 15 20	
AAG CGC GTA CCC TGC GAG ACA CCG CCC TTC ACT GTC GGA GAA CTC AAG	209
Lys Arg Val Pro Cys Glu Thr Pro Pro Phe Thr Val Gly Glu Leu Lys	
25 30 35	
AAA GCA ATC CCA CCG CAC TGT TTC AAG CGC TCG ATC CCT CGC TCT TTC	257
Lys Ala Ile Pro Pro His Cys Phe Lys Arg Ser Ile Pro Arg Ser Phe	
40 45 50	
TCC CAC CTC ATC TGG GAC ATC ATC ATA GCC TCC TGC TTC TAC TAC GTC	305
Ser His Leu Ile Trp Asp Ile Ile Ile Ala Ser Cys Phe Tyr Tyr Val	
55 60 65	
GCC ACC ACT TAC TTC CCT CTC CTC CCT AAC CCT CTC TCC TAC TTC GCC	353
Ala Thr Thr Tyr Phe Pro Leu Leu Pro Asn Pro Leu Ser Tyr Phe Ala	
70 75 80 85	
TGG CCT CTC TAC TGG GCC TGC CAG GGC TGC GTC CTA ACC GGC GTC TGG	401
Trp Pro Leu Tyr Trp Ala Cys Gln Gly Cys Val Leu Thr Gly Val Trp	
90 95 100	
GTC ATA GCC CAC GAG TGC GGC CAC GCA GCC TTC AGC GAC TAC CAG TGG	449
Val Ile Ala His Glu Cys Gly His Ala Ala Phe Ser Asp Tyr Gln Trp	
105 110 115	
CTG GAC GAC ACC GTC GGC CTC ATC TTC CAC TCC TTC CTC CTC GTC CCT	497
Leu Asp Asp Thr Val Gly Leu Ile Phe His Ser Phe Leu Leu Val Pro	
120 125 130	

TAC	TTC	TCC	TGG	AAG	TAC	AGT	CAT	CGA	CGC	CAC	CAT	TCC	AAC	ACT	GGC	545
Tyr	Phe	Ser	Trp	Lys	Tyr	Ser	His	Arg	Arg	His	His	Ser	Asn	Thr	Gly	
	135					140					145					
TCC	CTC	GAG	AGA	GAC	GAA	GTG	TTT	GTC	CCA	AGA	AGA	AGT	CAG	ACA	TCA	593
Ser	Leu	Glu	Arg	Asp	Glu	Val	Phe	Val	Pro	Arg	Arg	Ser	Gln	Thr	Ser	
150					155					160					165	
AGT	GGT	ACG	GCA	AGT	ACC	TCA	ACA	ACC	TTT	GGA	CGC	ACC	GTG	ATG	TTA	641
Ser	Gly	Thr	Ala	Ser	Thr	Ser	Thr	Thr	Phe	Gly	Arg	Thr	Val	Met	Leu	
				170					175					180		
ACG	GTT	CAG	TTC	ACT	CTC	GGC	TGG	CCT	TTG	TAC	TTA	GCC	TTC	AAC	GTC	689
Thr	Val	Gln	Phe	Thr	Leu	Gly	Trp	Pro	Leu	Tyr	Leu	Ala	Phe	Asn	Val	
			185					190					195			
TCG	GGG	AGA	CCT	TAC	GAC	GGC	GGC	TTC	GCT	TGC	CAT	TTC	CAC	CCC	AAC	737
Ser	Gly	Arg	Pro	Tyr	Asp	Gly	Gly	Phe	Ala	Cys	His	Phe	His	Pro	Asn	
		200				205						210				
GCT	CCC	ATC	TAC	AAC	GAC	CGT	GAG	CGT	CTC	CAG	ATA	TAC	ATC	TCC	GAC	785
Ala	Pro	Ile	Tyr	Asn	Asp	Arg	Glu	Arg	Leu	Gln	Ile	Tyr	Ile	Ser	Asp	
	215					220					225					
GCT	GGC	ATC	CTC	GCC	GTC	TGC	TAC	GGT	CTG	CTA	CCG	TAC	GCT	GCT	GTC	833
Ala	Gly	Ile	Leu	Ala	Val	Cys	Tyr	Gly	Leu	Leu	Pro	Tyr	Ala	Ala	Val	
230					235					240					245	
CAA	GGA	GTT	GCC	TCG	ATG	GTC	TGC	TTC	CTA	CGA	GTT	CCT	CTT	CTG	ATT	881
Gln	Gly	Val	Ala	Ser	Met	Val	Cys	Phe	Leu	Arg	Val	Pro	Leu	Leu	Ile	
				250					255					260		
GTC	AAC	GGG	TTC	TTA	GTT	TTG	ATC	ACT	TAC	TTG	CAG	CAC	ACG	CAT	CCT	929
Val	Asn	Gly	Phe	Leu	Val	Leu	Ile	Thr	Tyr	Leu	Gln	His	Thr	His	Pro	
			265					270					275			
TCC	CTG	CCT	CAC	TAT	GAC	TCG	TCT	GAG	TGG	GAT	TGG	TTG	AGG	GGA	GCT	977
Ser	Leu	Pro	His	Tyr	Asp	Ser	Ser	Glu	Trp	Asp	Trp	Leu	Arg	Gly	Ala	
		280					285					290				
TTG	GCC	ACC	GTT	GAC	AGA	GAC	TAC	GGA	ATC	TTG	AAC	CAA	GGC	TTC	CAC	1025
Leu	Ala	Thr	Val	Asp	Arg	Asp	Tyr	Gly	Ile	Leu	Asn	Gln	Gly	Phe	His	
	295					300					305					
AAT	ATC	ACG	GAC	ACG	CAC	GAG	GCG	CAT	CAC	CTG	TTC	TCG	ACC	ATG	CCG	1073
Asn	Ile	Thr	Asp	Thr	His	Glu	Ala	His	His	Leu	Phe	Ser	Thr	Met	Pro	
310					315					320					325	
CAT	TAT	CAT	GCG	ATG	GAA	GCT	ACG	AAG	GCG	ATA	AAG	CCG	ATA	CTG	GGA	1121
His	Tyr	His	Ala	Met	Glu	Ala	Thr	Lys	Ala	Ile	Lys	Pro	Ile	Leu	Gly	
				330					335					340		
GAG	TAT	TAT	CAG	TTC	GAT	GGG	ACG	CCG	GTG	GTT	AAG	GCG	ATG	TGG	AGG	1169
Glu	Tyr	Tyr	Gln	Phe	Asp	Gly	Thr	Pro	Val	Val	Lys	Ala	Met	Trp	Arg	
			345					350					355			

GAG GCG AAG GAG TGT ATC TAT GTG GAA CCG GAC AGG CAA GGT GAG AAG 1217  
 Glu Ala Lys Glu Cys Ile Tyr Val Glu Pro Asp Arg Gln Gly Glu Lys  
           360                          365                          370

AAA GGT GTG TTC TGG TAC AAC AAT AAG TTA TGAAGCAAAG AAGAACTGA 1267  
 Lys Gly Val Phe Trp Tyr Asn Asn Lys Leu  
           375                          380

ACCTTTCTCT TCTATCAATT GTCTTTGTTT AAGAAGCTAT GTTTCTGTTT CAATAATCTT 1327

AATTATCCAT TTTGTTGTGT TTTCTGACAT TTTGGCTAAA ATTATGTGAT GTTGGAAGTT 1387

AGTGTCT 1394

## (2) INFORMATION FOR SEQ ID NO:4:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 383 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

Met Gly Ala Gly Gly Arg Met Gln Val Ser Pro Pro Ser Lys Lys Ser  
   1                          5                          10                          15

Glu Thr Asp Asn Ile Lys Arg Val Pro Cys Glu Thr Pro Pro Phe Thr  
           20                          25                          30

Val Gly Glu Leu Lys Lys Ala Ile Pro Pro His Cys Phe Lys Arg Ser  
           35                          40                          45

Ile Pro Arg Ser Phe Ser His Leu Ile Trp Asp Ile Ile Ile Ala Ser  
           50                          55                          60

Cys Phe Tyr Tyr Val Ala Thr Thr Tyr Phe Pro Leu Leu Pro Asn Pro  
           65                          70                          75                          80

Leu Ser Tyr Phe Ala Trp Pro Leu Tyr Trp Ala Cys Gln Gly Cys Val  
                           85                          90                          95

Leu Thr Gly Val Trp Val Ile Ala His Glu Cys Gly His Ala Ala Phe  
           100                          105                          110

Ser Asp Tyr Gln Trp Leu Asp Asp Thr Val Gly Leu Ile Phe His Ser  
           115                          120                          125

Phe Leu Leu Val Pro Tyr Phe Ser Trp Lys Tyr Ser His Arg Arg His  
           130                          135                          140

His Ser Asn Thr Gly Ser Leu Glu Arg Asp Glu Val Phe Val Pro Arg  
           145                          150                          155                          160

Arg Ser Gln Thr Ser Ser Gly Thr Ala Ser Thr Ser Thr Thr Phe Gly  
                           165                          170                          175

009207" 6226950

Arg Thr Val Met Leu Thr Val Gln Phe Thr Leu Gly Trp Pro Leu Tyr  
 180 185 190  
 Leu Ala Phe Asn Val Ser Gly Arg Pro Tyr Asp Gly Gly Phe Ala Cys  
 195 200 205  
 His Phe His Pro Asn Ala Pro Ile Tyr Asn Asp Arg Glu Arg Leu Gln  
 210 215 220  
 Ile Tyr Ile Ser Asp Ala Gly Ile Leu Ala Val Cys Tyr Gly Leu Leu  
 225 230 235 240  
 Pro Tyr Ala Ala Val Gln Gly Val Ala Ser Met Val Cys Phe Leu Arg  
 245 250 255  
 Val Pro Leu Leu Ile Val Asn Gly Phe Leu Val Leu Ile Thr Tyr Leu  
 260 265 270  
 Gln His Thr His Pro Ser Leu Pro His Tyr Asp Ser Ser Glu Trp Asp  
 275 280 285  
 Trp Leu Arg Gly Ala Leu Ala Thr Val Asp Arg Asp Tyr Gly Ile Leu  
 290 295 300  
 Asn Gln Gly Phe His Asn Ile Thr Asp Thr His Glu Ala His His Leu  
 305 310 315 320  
 Phe Ser Thr Met Pro His Tyr His Ala Met Glu Ala Thr Lys Ala Ile  
 325 330 335  
 Lys Pro Ile Leu Gly Glu Tyr Tyr Gln Phe Asp Gly Thr Pro Val Val  
 340 345 350  
 Lys Ala Met Trp Arg Glu Ala Lys Glu Cys Ile Tyr Val Glu Pro Asp  
 355 360 365  
 Arg Gln Gly Glu Lys Lys Gly Val Phe Trp Tyr Asn Asn Lys Leu  
 370 375 380

(2) INFORMATION FOR SEQ ID NO:5:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 1462 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: double
  - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: cDNA
- (iii) HYPOTHETICAL: NO
- (iv) ANTI-SENSE: NO
- (vi) ORIGINAL SOURCE:
  - (A) ORGANISM: Glycine max

009201" 6/26/95

(vii) IMMEDIATE SOURCE:  
(B) CLONE: pSF2-165K

(ix) FEATURE:  
(A) NAME/KEY: CDS  
(B) LOCATION: 108..1247

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

CCATATACTA ATATTTGCTT GTATTGATAG CCCCTCCGTT CCCAAGAGTA TAAAACTGCA	60
TCGAATAATA CAAGCCACTA GGCATGGGTC TAGCAAAGGA AACAACA ATG GGA GGT	116
Met Gly Gly	
1	
AGA GGT CGT GTG GCC AAA GTG GAA GTT CAA GGG AAG AAG CCT CTC TCA	164
Arg Gly Arg Val Ala Lys Val Glu Val Gln Gly Lys Lys Pro Leu Ser	
5 10 15	
AGG GTT CCA AAC ACA AAG CCA CCA TTC ACT GTT GGC CAA CTC AAG AAA	212
Arg Val Pro Asn Thr Lys Pro Pro Phe Thr Val Gly Gln Leu Lys Lys	
20 25 30 35	
GCA ATT CCA CCA CAC TGC TTT CAG CGC TCC CTC CTC ACT TCA TTC TCC	260
Ala Ile Pro Pro His Cys Phe Gln Arg Ser Leu Leu Thr Ser Phe Ser	
40 45 50	
TAT GTT GTT TAT GAC CTT TCA TTT GCC TTC ATT TTC TAC ATT GCC ACC	308
Tyr Val Val Tyr Asp Leu Ser Phe Ala Phe Ile Phe Tyr Ile Ala Thr	
55 60 65	
ACC TAC TTC CAC CTC CTT CCT CAA CCC TTT TCC CTC ATT GCA TGG CCA	356
Thr Tyr Phe His Leu Leu Pro Gln Pro Phe Ser Leu Ile Ala Trp Pro	
70 75 80	
ATC TAT TGG GTT CTC CAA GGT TGC CTT CTC ACT GGT GTG TGG GTG ATT	404
Ile Tyr Trp Val Leu Gln Gly Cys Leu Leu Thr Gly Val Trp Val Ile	
85 90 95	
GCT CAC GAG TGT GGT CAC CAT GCC TTC AGC AAG TAC CAA TGG GTT GAT	452
Ala His Glu Cys Gly His His Ala Phe Ser Lys Tyr Gln Trp Val Asp	
100 105 110 115	
GAT GTT GTG GGT TTG ACC CTT CAC TCA ACA CTT TTA GTC CCT TAT TTC	500
Asp Val Val Gly Leu Thr Leu His Ser Thr Leu Leu Val Pro Tyr Phe	
120 125 130	
TCA TGG AAA ATA AGC CAT CGC CGC CAT CAC TCC AAC ACA GGT TCC CTT	548
Ser Trp Lys Ile Ser His Arg Arg His His Ser Asn Thr Gly Ser Leu	
135 140 145	
GAC CGT GAT GAA GTG TTT GTC CCA AAA CCA AAA TCC AAA GTT GCA TGG	596
Asp Arg Asp Glu Val Phe Val Pro Lys Pro Lys Ser Lys Val Ala Trp	
150 155 160	

TTT	TCC	AAG	TAC	TTA	AAC	AAC	CCT	CTA	GGA	AGG	GCT	GTT	TCT	CTT	CTC	644
Phe	Ser	Lys	Tyr	Leu	Asn	Asn	Pro	Leu	Gly	Arg	Ala	Val	Ser	Leu	Leu	
165						170					175					
GTC	ACA	CTC	ACA	ATA	GGG	TGG	CCT	ATG	TAT	TTA	GCC	TTC	AAT	GTC	TCT	692
Val	Thr	Leu	Thr	Ile	Gly	Trp	Pro	Met	Tyr	Leu	Ala	Phe	Asn	Val	Ser	
180					185					190					195	
GGT	AGA	CCC	TAT	GAT	AGT	TTT	GCA	AGC	CAC	TAC	CAC	CCT	TAT	GCT	CCC	740
Gly	Arg	Pro	Tyr	Asp	Ser	Phe	Ala	Ser	His	Tyr	His	Pro	Tyr	Ala	Pro	
				200					205					210		
ATA	TAT	TCT	AAC	CGT	GAG	AGG	CTT	CTG	ATC	TAT	GTC	TCT	GAT	GTT	GCT	788
Ile	Tyr	Ser	Asn	Arg	Glu	Arg	Leu	Leu	Ile	Tyr	Val	Ser	Asp	Val	Ala	
			215					220					225			
TTG	TTT	TCT	GTG	ACT	TAC	TCT	CTC	TAC	CGT	GTT	GCA	ACC	CTG	AAA	GGG	836
Leu	Phe	Ser	Val	Thr	Tyr	Ser	Leu	Tyr	Arg	Val	Ala	Thr	Leu	Lys	Gly	
		230					235					240				
TTG	GTT	TGG	CTG	CTA	TGT	GTT	TAT	GGG	GTG	CCT	TTG	CTC	ATT	GTG	AAC	884
Leu	Val	Trp	Leu	Leu	Cys	Val	Tyr	Gly	Val	Pro	Leu	Leu	Ile	Val	Asn	
	245					250					255					
GGT	TTT	CTT	GTG	ACT	ATC	ACA	TAT	TTG	CAG	CAC	ACA	CAC	TTT	GCC	TTG	932
Gly	Phe	Leu	Val	Thr	Ile	Thr	Tyr	Leu	Gln	His	Thr	His	Phe	Ala	Leu	
260					265					270					275	
CCT	CAT	TAC	GAT	TCA	TCA	GAA	TGG	GAC	TGG	CTG	AAG	GGA	GCT	TTG	GCA	980
Pro	His	Tyr	Asp	Ser	Ser	Glu	Trp	Asp	Trp	Leu	Lys	Gly	Ala	Leu	Ala	
				280					285					290		
ACT	ATG	GAC	AGA	GAT	TAT	GGG	ATT	CTG	AAC	AAG	GTG	TTT	CAT	CAC	ATA	1028
Thr	Met	Asp	Arg	Asp	Tyr	Gly	Ile	Leu	Asn	Lys	Val	Phe	His	His	Ile	
			295					300					305			
ACT	GAT	ACT	CAT	GTG	GCT	CAC	CAT	CTC	TTC	TCT	ACA	ATG	CCA	CAT	TAC	1076
Thr	Asp	Thr	His	Val	Ala	His	His	Leu	Phe	Ser	Thr	Met	Pro	His	Tyr	
		310					315					320				
CAT	GCA	ATG	GAG	GCA	ACC	AAT	GCA	ATC	AAG	CCA	ATA	TTG	GGT	GAG	TAC	1124
His	Ala	Met	Glu	Ala	Thr	Asn	Ala	Ile	Lys	Pro	Ile	Leu	Gly	Glu	Tyr	
	325					330					335					
TAC	CAA	TTT	GAT	GAC	ACA	CCA	TTT	TAC	AAG	GCA	CTG	TGG	AGA	GAA	GCG	1172
Tyr	Gln	Phe	Asp	Asp	Thr	Pro	Phe	Tyr	Lys	Ala	Leu	Trp	Arg	Glu	Ala	
340					345					350					355	
AGA	GAG	TGC	CTC	TAT	GTG	GAG	CCA	GAT	GAA	GGA	ACA	TCC	GAG	AAG	GGC	1220
Arg	Glu	Cys	Leu	Tyr	Val	Glu	Pro	Asp	Glu	Gly	Thr	Ser	Glu	Lys	Gly	
				360					365					370		
GTG	TAT	TGG	TAC	AGG	AAC	AAG	TAT	TGATGGAGCA	ACCAATGGGC	CATAGTGGGA						1274
Val	Tyr	Trp	Tyr	Arg	Asn	Lys	Tyr									
			375				380									
GTTATGGAAG	TTTTGT	CATG	TATTAGTACA	TAATTAGTAG	AATGTTATAA	ATAAGTGGAT										1334

TTGCCGCGTA ATGACTTTGT GTGTATTGTG AAACAGCTTG TTGCGATCAT GGTTATAATG 1394  
TAAAAATAAT TCTGGTATTA ATTACATGTG GAAAGTGTTT TGCTTATAGC TTTCTGCCTA 1454  
AAAAAATA 1462

## (2) INFORMATION FOR SEQ ID NO:6:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 379 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

Met Gly Gly Arg Gly Arg Val Ala Lys Val Glu Val Gln Gly Lys Lys  
1 5 10 15  
Pro Leu Ser Arg Val Pro Asn Thr Lys Pro Pro Phe Thr Val Gly Gln  
20 25 30  
Leu Lys Lys Ala Ile Pro Pro His Cys Phe Gln Arg Ser Leu Leu Thr  
35 40 45  
Ser Phe Ser Tyr Val Val Tyr Asp Leu Ser Phe Ala Phe Ile Phe Tyr  
50 55 60  
Ile Ala Thr Thr Tyr Phe His Leu Leu Pro Gln Pro Phe Ser Leu Ile  
65 70 75 80  
Ala Trp Pro Ile Tyr Trp Val Leu Gln Gly Cys Leu Leu Thr Gly Val  
85 90 95  
Trp Val Ile Ala His Glu Cys Gly His His Ala Phe Ser Lys Tyr Gln  
100 105 110  
Trp Val Asp Asp Val Val Gly Leu Thr Leu His Ser Thr Leu Leu Val  
115 120 125  
Pro Tyr Phe Ser Trp Lys Ile Ser His Arg Arg His His Ser Asn Thr  
130 135 140  
Gly Ser Leu Asp Arg Asp Glu Val Phe Val Pro Lys Pro Lys Ser Lys  
145 150 155 160  
Val Ala Trp Phe Ser Lys Tyr Leu Asn Asn Pro Leu Gly Arg Ala Val  
165 170 175  
Ser Leu Leu Val Thr Leu Thr Ile Gly Trp Pro Met Tyr Leu Ala Phe  
180 185 190  
Asn Val Ser Gly Arg Pro Tyr Asp Ser Phe Ala Ser His Tyr His Pro  
195 200 205

00920T 52E25950



Tyr Ala Pro Ile Tyr Ser Asn Arg Glu Arg Leu Leu Ile Tyr Val Ser  
 210 215 220  
 Asp Val Ala Leu Phe Ser Val Thr Tyr Ser Leu Tyr Arg Val Ala Thr  
 225 230 235 240  
 Leu Lys Gly Leu Val Trp Leu Leu Cys Val Tyr Gly Val Pro Leu Leu  
 245 250 255  
 Ile Val Asn Gly Phe Leu Val Thr Ile Thr Tyr Leu Gln His Thr His  
 260 265 270  
 Phe Ala Leu Pro His Tyr Asp Ser Ser Glu Trp Asp Trp Leu Lys Gly  
 275 280 285  
 Ala Leu Ala Thr Met Asp Arg Asp Tyr Gly Ile Leu Asn Lys Val Phe  
 290 295 300  
 His His Ile Thr Asp Thr His Val Ala His His Leu Phe Ser Thr Met  
 305 310 315 320  
 Pro His Tyr His Ala Met Glu Ala Thr Asn Ala Ile Lys Pro Ile Leu  
 325 330 335  
 Gly Glu Tyr Tyr Gln Phe Asp Asp Thr Pro Phe Tyr Lys Ala Leu Trp  
 340 345 350  
 Arg Glu Ala Arg Glu Cys Leu Tyr Val Glu Pro Asp Glu Gly Thr Ser  
 355 360 365  
 Glu Lys Gly Val Tyr Trp Tyr Arg Asn Lys Tyr  
 370 375

(2) INFORMATION FOR SEQ ID NO:7:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 1790 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: double
  - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: cDNA
- (iii) HYPOTHETICAL: NO
- (iv) ANTI-SENSE: NO
- (vi) ORIGINAL SOURCE:
  - (A) ORGANISM: Zea mays
- (vii) IMMEDIATE SOURCE:
  - (B) CLONE: pFad2#1
- (ix) FEATURE:
  - (A) NAME/KEY: CDS
  - (B) LOCATION: 165..1328

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

CGGCCTCTCC CCTCCCTCCT CCCTGCAAAT CCTGCAGACA CCACCGCTCG TTTTCTCTC	60
CGGGACAGGA GAAAAGGGGA GAGAGAGGTG AGGCGCGGTG TCCGCCCGAT CTGCTCTGCC	120
CCGACGCAGC TGTTACGACC TCCTCAGTCT CAGTCAGGAG CAAG ATG GGT GCC GGC	176
Met Gly Ala Gly	
1	
GGC AGG ATG ACC GAG AAG GAG CGG GAG AAG CAG GAG CAG CTC GCC CGA	224
Gly Arg Met Thr Glu Lys Glu Arg Glu Lys Gln Glu Gln Leu Ala Arg	
5 10 15 20	
GCT ACC GGT GGC GCC GCG ATG CAG CGG TCG CCG GTG GAG AAG CCT CCG	272
Ala Thr Gly Gly Ala Ala Met Gln Arg Ser Pro Val Glu Lys Pro Pro	
25 30 35	
TTC ACT CTG GGT CAG ATC AAG AAG GCC ATC CCG CCA CAC TGC TTC GAG	320
Phe Thr Leu Gly Gln Ile Lys Lys Ala Ile Pro Pro His Cys Phe Glu	
40 45 50	
CGC TCG GTG CTC AAG TCC TTC TCG TAC GTG GTC CAC GAC CTG GTG ATC	368
Arg Ser Val Leu Lys Ser Phe Ser Tyr Val Val His Asp Leu Val Ile	
55 60 65	
GCC GCG GCG CTC CTC TAC TTC GCG CTG GCC ATC ATA CCG GCG CTC CCA	416
Ala Ala Ala Leu Leu Tyr Phe Ala Leu Ala Ile Ile Pro Ala Leu Pro	
70 75 80	
AGC CCG CTC CGC TAC GCC GCC TGG CCG CTG TAC TGG ATC GCG CAG GGG	464
Ser Pro Leu Arg Tyr Ala Ala Trp Pro Leu Tyr Trp Ile Ala Gln Gly	
85 90 95 100	
TGC GTG TGC ACC GGC GTG TGG GTC ATC GCG CAC GAG TGC GGC CAC CAC	512
Cys Val Cys Thr Gly Val Trp Val Ile Ala His Glu Cys Gly His His	
105 110 115	
GCC TTC TCG GAC TAC TCG CTC CTG GAC GAC GTG GTC GGC CTG GTG CTG	560
Ala Phe Ser Asp Tyr Ser Leu Leu Asp Asp Val Val Gly Leu Val Leu	
120 125 130	
CAC TCG TCG CTC ATG GTG CCC TAC TTC TCG TGG AAG TAC AGC CAC CGG	608
His Ser Ser Leu Met Val Pro Tyr Phe Ser Trp Lys Tyr Ser His Arg	
135 140 145	
CGC CAC CAC TCC AAC ACG GGG TCC CTG GAG CGC GAC GAG GTG TTC GTG	656
Arg His His Ser Asn Thr Gly Ser Leu Glu Arg Asp Glu Val Phe Val	
150 155 160	
CCC AAG AAG AAG GAG GCG CTG CCG TGG TAC ACC CCG TAC GTG TAC AAC	704
Pro Lys Lys Lys Glu Ala Leu Pro Trp Tyr Thr Pro Tyr Val Tyr Asn	
165 170 175 180	

AAC CCG GTC GGC CGG GTG GTG CAC ATC GTG GTG CAG CTC ACC CTC GGG Asn Pro Val Gly Arg Val Val His Ile Val Val Gln Leu Thr Leu Gly 185 190 195	752
TGG CCG CTG TAC CTG GCG ACC AAC GCG TCG GGG CGG CCG TAC CCG CGC Trp Pro Leu Tyr Leu Ala Thr Asn Ala Ser Gly Arg Pro Tyr Pro Arg 200 205 210	800
TTC GCC TGC CAC TTC GAC CCC TAC GGC CCC ATC TAC AAC GAC CGG GAG Phe Ala Cys His Phe Asp Pro Tyr Gly Pro Ile Tyr Asn Asp Arg Glu 215 220 225	848
CGC GCC CAG ATC TTC GTC TCG GAC GCC GGC GTC GTG GCC GTG GCG TTC Arg Ala Gln Ile Phe Val Ser Asp Ala Gly Val Val Ala Val Ala Phe 230 235 240	896
GGG CTG TAC AAG CTG GCG GCG GCG TTC GGG GTC TGG TGG GTG GTG CGC Gly Leu Tyr Lys Leu Ala Ala Ala Phe Gly Val Trp Trp Val Val Arg 245 250 255 260	944
GTG TAC GCC GTG CCG CTG CTG ATC GTG AAC GCG TGG CTG GTG CTC ATC Val Tyr Ala Val Pro Leu Leu Ile Val Asn Ala Trp Leu Val Leu Ile 265 270 275	992
ACC TAC CTG CAG CAC ACC CAC CCG TCG CTC CCC CAC TAC GAC TCG AGC Thr Tyr Leu Gln His Thr His Pro Ser Leu Pro His Tyr Asp Ser Ser 280 285 290	1040
GAG TGG GAC TGG CTG CGC GGC GCG CTG GCC ACC ATG GAC CGC GAC TAC Glu Trp Asp Trp Leu Arg Gly Ala Leu Ala Thr Met Asp Arg Asp Tyr 295 300 305	1088
GGC ATC CTC AAC CGC GTG TTC CAC AAC ATC ACG GAC ACG CAC GTC GCG Gly Ile Leu Asn Arg Val Phe His Asn Ile Thr Asp Thr His Val Ala 310 315 320	1136
CAC CAC CTC TTC TCC ACC ATG CCG CAC TAC CAC GCC ATG GAG GCC ACC His His Leu Phe Ser Thr Met Pro His Tyr His Ala Met Glu Ala Thr 325 330 335 340	1184
AAG GCG ATC AGG CCC ATC CTC GGC GAC TAC TAC CAC TTC GAC CCG ACC Lys Ala Ile Arg Pro Ile Leu Gly Asp Tyr Tyr His Phe Asp Pro Thr 345 350 355	1232
CCT GTC GCC AAG GCG ACC TGG CGC GAG GCC GGG GAA TGC ATC TAC GTC Pro Val Ala Lys Ala Thr Trp Arg Glu Ala Gly Glu Cys Ile Tyr Val 360 365 370	1280
GAG CCC GAG GAC CGC AAG GGC GTC TTC TGG TAC AAC AAG AAG TTC TAGCCGCCGC Glu Pro Glu Asp Arg Lys Gly Val Phe Trp Tyr Asn Lys Lys Phe 375 380 385	1335
CGCTCGCAGA GCTGAGGACG CTACCGTAGG AATGGGAGCA GAAACCAGGA GGAGGAGACG	1395
GTACTCGCCC CAAAGTCTCC GTCAACCTAT CTAATCGTTA GTCGTCAGTC TTTTAGACGG	1455
GAAGAGAGAT CATTTGGGCA CAGAGACGAA GGCTTACTGC AGTGCCATCG CTAGAGCTGC	1515

CATCAAGTAC AAGTAGGCAA ATTCGTCAAC TTAGTGTGTC CCATGTTGTT TTTCTTAGTC 1575  
 GTCCGCTGCT GTAGGCTTTC CGGCGGCGGT CGTTTGTGTG GTTGGCATCC GTGGCCATGC 1635  
 CTGTGCGTGC GTGGCCGCGC TTGTCGTGTG CGTCTGTCGT CGCGTTGGCG TCGTCTCTTC 1695  
 GTGCTCCCCG TGTGTTGTTG TAAACAAGA AGATGTTTTTC TGGTGTCTTT GCGGAATAA 1755  
 CAGATCGTCC GAACGAAAAA AAAAAAAAAA AAAAA 1790

(2) INFORMATION FOR SEQ ID NO:8:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 387 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

Met	Gly	Ala	Gly	Gly	Arg	Met	Thr	Glu	Lys	Glu	Arg	Glu	Lys	Gln	Glu	1	5	10	15
Gln	Leu	Ala	Arg	Ala	Thr	Gly	Gly	Ala	Ala	Met	Gln	Arg	Ser	Pro	Val	20	25	30	
Glu	Lys	Pro	Pro	Phe	Thr	Leu	Gly	Gln	Ile	Lys	Lys	Ala	Ile	Pro	Pro	35	40	45	
His	Cys	Phe	Glu	Arg	Ser	Val	Leu	Lys	Ser	Phe	Ser	Tyr	Val	Val	His	50	55	60	
Asp	Leu	Val	Ile	Ala	Ala	Ala	Leu	Leu	Tyr	Phe	Ala	Leu	Ala	Ile	Ile	65	70	75	80
Pro	Ala	Leu	Pro	Ser	Pro	Leu	Arg	Tyr	Ala	Ala	Trp	Pro	Leu	Tyr	Trp	85	90	95	
Ile	Ala	Gln	Gly	Cys	Val	Cys	Thr	Gly	Val	Trp	Val	Ile	Ala	His	Glu	100	105	110	
Cys	Gly	His	His	Ala	Phe	Ser	Asp	Tyr	Ser	Leu	Leu	Asp	Asp	Val	Val	115	120	125	
Gly	Leu	Val	Leu	His	Ser	Ser	Leu	Met	Val	Pro	Tyr	Phe	Ser	Trp	Lys	130	135	140	
Tyr	Ser	His	Arg	Arg	His	His	Ser	Asn	Thr	Gly	Ser	Leu	Glu	Arg	Asp	145	150	155	160
Glu	Val	Phe	Val	Pro	Lys	Lys	Lys	Glu	Ala	Leu	Pro	Trp	Tyr	Thr	Pro	165	170	175	
Tyr	Val	Tyr	Asn	Asn	Pro	Val	Gly	Arg	Val	Val	His	Ile	Val	Val	Gln	180	185	190	

009201" 52E25950

Leu Thr Leu Gly Trp Pro Leu Tyr Leu Ala Thr Asn Ala Ser Gly Arg  
 195 200 205  
 Pro Tyr Pro Arg Phe Ala Cys His Phe Asp Pro Tyr Gly Pro Ile Tyr  
 210 215 220  
 Asn Asp Arg Glu Arg Ala Gln Ile Phe Val Ser Asp Ala Gly Val Val  
 225 230 235 240  
 Ala Val Ala Phe Gly Leu Tyr Lys Leu Ala Ala Ala Phe Gly Val Trp  
 245 250 255  
 Trp Val Val Arg Val Tyr Ala Val Pro Leu Leu Ile Val Asn Ala Trp  
 260 265 270  
 Leu Val Leu Ile Thr Tyr Leu Gln His Thr His Pro Ser Leu Pro His  
 275 280 285  
 Tyr Asp Ser Ser Glu Trp Asp Trp Leu Arg Gly Ala Leu Ala Thr Met  
 290 295 300  
 Asp Arg Asp Tyr Gly Ile Leu Asn Arg Val Phe His Asn Ile Thr Asp  
 305 310 315 320  
 Thr His Val Ala His His Leu Phe Ser Thr Met Pro His Tyr His Ala  
 325 330 335  
 Met Glu Ala Thr Lys Ala Ile Arg Pro Ile Leu Gly Asp Tyr Tyr His  
 340 345 350  
 Phe Asp Pro Thr Pro Val Ala Lys Ala Thr Trp Arg Glu Ala Gly Glu  
 355 360 365  
 Cys Ile Tyr Val Glu Pro Glu Asp Arg Lys Gly Val Phe Trp Tyr Asn  
 370 375 380  
 Lys Lys Phe  
 385

## (2) INFORMATION FOR SEQ ID NO:9:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 673 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Ricinus communis

(vii) IMMEDIATE SOURCE:  
(B) CLONE: prf2-1C

(ix) FEATURE:  
(A) NAME/KEY: CDS  
(B) LOCATION: 1..673

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

TGG	GTG	ATG	GCG	CAT	GAT	TGT	GGG	CAC	CAT	GCC	TTC	AGT	GAC	TAT	CAA	48
Trp	Val	Met	Ala	His	Asp	Cys	Gly	His	His	Ala	Phe	Ser	Asp	Tyr	Gln	
1				5					10					15		
TTG	CTT	GAT	GAT	GTA	GTT	GGT	CTT	ATC	CTA	CAC	TCC	TGT	CTC	CTT	GTC	96
Leu	Leu	Asp	Asp	Val	Val	Gly	Leu	Ile	Leu	His	Ser	Cys	Leu	Leu	Val	
			20					25					30			
CCT	TAT	TTT	TCA	TGG	AAA	CAC	AGC	CAT	CGC	CGA	CAT	CAT	TCC	AAC	ACA	144
Pro	Tyr	Phe	Ser	Trp	Lys	His	Ser	His	Arg	Arg	His	His	Ser	Asn	Thr	
		35					40					45				
GGG	TCC	CTG	GAA	CGG	GAT	GAA	GTG	TTT	GTT	CCC	AAG	AAG	AAA	TCT	AGT	192
Gly	Ser	Leu	Glu	Arg	Asp	Glu	Val	Phe	Val	Pro	Lys	Lys	Lys	Ser	Ser	
	50					55				60						
ATC	CGT	TGG	TAT	TCC	AAA	TAC	CTC	AAC	AAC	CCT	CCA	GGT	CGT	ATC	ATG	240
Ile	Arg	Trp	Tyr	Ser	Lys	Tyr	Leu	Asn	Asn	Pro	Pro	Gly	Arg	Ile	Met	
65					70					75					80	
ACA	ATT	GCC	GTC	ACA	CTT	TCA	CTT	GGC	TGG	CCT	CTG	TAC	CTA	GCA	TTC	288
Thr	Ile	Ala	Val	Thr	Leu	Ser	Leu	Gly	Trp	Pro	Leu	Tyr	Leu	Ala	Phe	
				85					90					95		
AAT	GTT	TCA	GGC	AGG	CCA	TAT	GAT	CGG	TTC	GCC	TGC	CAC	TAT	GAC	CCA	336
Asn	Val	Ser	Gly	Arg	Pro	Tyr	Asp	Arg	Phe	Ala	Cys	His	Tyr	Asp	Pro	
			100					105					110			
TAT	GGC	CCG	ATC	TAC	AAT	GAT	CGC	GAG	CGA	ATC	GAG	ATA	TTC	ATA	TCA	384
Tyr	Gly	Pro	Ile	Tyr	Asn	Asp	Arg	Glu	Arg	Ile	Glu	Ile	Phe	Ile	Ser	
		115					120					125				
GAT	GCT	GGT	GTT	CTT	GCT	GTC	ACT	TTT	GGT	CTC	TAC	CAA	CTT	GCT	ATA	432
Asp	Ala	Gly	Val	Leu	Ala	Val	Thr	Phe	Gly	Leu	Tyr	Gln	Leu	Ala	Ile	
	130					135					140					
GCG	AAG	GGG	CTT	GCT	TGG	GTT	GTC	TGT	GTA	TAT	GGA	GTG	CCA	TTG	TTG	480
Ala	Lys	Gly	Leu	Ala	Trp	Val	Val	Cys	Val	Tyr	Gly	Val	Pro	Leu	Leu	
145					150					155				160		
GTG	GTG	AAT	TCA	TTC	CTT	GTT	CTG	ATC	ACA	TTT	CTG	CAG	CAT	ACT	CAC	528
Val	Val	Asn	Ser	Phe	Leu	Val	Leu	Ile	Thr	Phe	Leu	Gln	His	Thr	His	
				165					170					175		
CCT	GCA	TTG	CCA	CAT	TAT	GAT	TCG	TCG	GAG	TGG	GAC	TGG	CTA	AGA	GGA	576
Pro	Ala	Leu	Pro	His	Tyr	Asp	Ser	Ser	Glu	Trp	Asp	Trp	Leu	Arg	Gly	
			180					185					190			

009201" 52E25950

GCT CTA GCA ACT GTT GAC AGA GAT TAC GGG ATC TTG AAC AAG GTG TTC 624  
 Ala Leu Ala Thr Val Asp Arg Asp Tyr Gly Ile Leu Asn Lys Val Phe  
           195                                  200                                  205

CAT AAC ATA ACG GAC ACT CAA GTA GCT CAC CAC CTT TTC ACC ATG CCC C 673  
 His Asn Ile Thr Asp Thr Gln Val Ala His His Leu Phe Thr Met Pro  
           210                                  215                                  220

(2) INFORMATION FOR SEQ ID NO:10:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 224 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

Trp Val Met Ala His Asp Cys Gly His His Ala Phe Ser Asp Tyr Gln  
   1                                  5                                  10                                  15  
 Leu Leu Asp Asp Val Val Gly Leu Ile Leu His Ser Cys Leu Leu Val  
                                   20                                  25                                  30  
 Pro Tyr Phe Ser Trp Lys His Ser His Arg Arg His His Ser Asn Thr  
                                   35                                  40                                  45  
 Gly Ser Leu Glu Arg Asp Glu Val Phe Val Pro Lys Lys Lys Ser Ser  
                                   50                                  55                                  60  
 Ile Arg Trp Tyr Ser Lys Tyr Leu Asn Asn Pro Pro Gly Arg Ile Met  
   65                                  70                                  75                                  80  
 Thr Ile Ala Val Thr Leu Ser Leu Gly Trp Pro Leu Tyr Leu Ala Phe  
                                   85                                  90                                  95  
 Asn Val Ser Gly Arg Pro Tyr Asp Arg Phe Ala Cys His Tyr Asp Pro  
                                   100                                  105                                  110  
 Tyr Gly Pro Ile Tyr Asn Asp Arg Glu Arg Ile Glu Ile Phe Ile Ser  
                                   115                                  120                                  125  
 Asp Ala Gly Val Leu Ala Val Thr Phe Gly Leu Tyr Gln Leu Ala Ile  
                                   130                                  135                                  140  
 Ala Lys Gly Leu Ala Trp Val Val Cys Val Tyr Gly Val Pro Leu Leu  
   145                                  150                                  155                                  160  
 Val Val Asn Ser Phe Leu Val Leu Ile Thr Phe Leu Gln His Thr His  
                                   165                                  170                                  175  
 Pro Ala Leu Pro His Tyr Asp Ser Ser Glu Trp Asp Trp Leu Arg Gly  
                                   180                                  185                                  190  
 Ala Leu Ala Thr Val Asp Arg Asp Tyr Gly Ile Leu Asn Lys Val Phe  
                                   195                                  200                                  205

009207" 62625960

His Asn Ile Thr Asp Thr Gln Val Ala His His Leu Phe Thr Met Pro  
 210 215 220

## (2) INFORMATION FOR SEQ ID NO:11:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1369 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: double  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

## (vi) ORIGINAL SOURCE:

(A) ORGANISM: Ricinus communis

## (vii) IMMEDIATE SOURCE:

(B) CLONE: pRF197c-42

## (ix) FEATURE:

- (A) NAME/KEY: CDS  
 (B) LOCATION: 184..1347

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

CGGCCGGGAT TCCGGTTTTT CACTAATTT GCAAAAAATG CATGATTTCA CCTCAAATCA	60
AACACCACAC CTTATAACTT AGTCTTAAGA GAGAGAGAGA GAGGAGACAT TTCTCTTCTC	120
TGAGATGAGC ACTTCTCTTC CAGACATCGA AGCCTCAGGA AAGTGCTTGA GAAGAGCTTG	180
AGA ATG GGA GGT GGT GGT CGC ATG TCT ACT GTC ATA ATC AGC AAC AAC	228
Met Gly Gly Gly Gly Arg Met Ser Thr Val Ile Ile Ser Asn Asn	
1 5 10 15	
AGT GAG AAG AAA GGA GGA AGC AGC CAC CTG GAG CGA GCG CCG CAC ACG	276
Ser Glu Lys Lys Gly Gly Ser Ser His Leu Glu Arg Ala Pro His Thr	
20 25 30	
AAG CCT CCT TAC ACA CTT GGT AAC CTC AAG AGA GCC ATC CCA CCC CAT	324
Lys Pro Pro Tyr Thr Leu Gly Asn Leu Lys Arg Ala Ile Pro Pro His	
35 40 45	
TGC TTT GAA CGC TCT TTT GTG CGC TCA TTC TCC AAT TTT GCC TAT AAT	372
Cys Phe Glu Arg Ser Phe Val Arg Ser Phe Ser Asn Phe Ala Tyr Asn	
50 55 60	
TTC TGC TTA AGT TTT CTT TCC TAC TCG ATC GCC ACC AAC TTC TTC CCT	420
Phe Cys Leu Ser Phe Leu Ser Tyr Ser Ile Ala Thr Asn Phe Phe Pro	
65 70 75	

00920T"52E/5950



TAC	ATC	TCT	TCT	CCG	CTC	TCG	TAT	GTC	GCT	TGG	CTG	GTT	TAC	TGG	CTC	468
Tyr	Ile	Ser	Ser	Pro	Leu	Ser	Tyr	Val	Ala	Trp	Leu	Val	Tyr	Trp	Leu	
80					85					90					95	
TTC	CAA	GGC	TGC	ATT	CTC	ACT	GGT	CTT	TGG	GTC	ATC	GGC	CAT	GAA	TGT	516
Phe	Gln	Gly	Cys	Ile	Leu	Thr	Gly	Leu	Trp	Val	Ile	Gly	His	Glu	Cys	
				100					105					110		
GGC	CAT	CAT	GCT	TTT	AGT	GAG	TAT	CAG	CTG	GCT	GAT	GAC	ATT	GTT	GGC	564
Gly	His	His	Ala	Phe	Ser	Glu	Tyr	Gln	Leu	Ala	Asp	Asp	Ile	Val	Gly	
			115					120					125			
CTA	ATT	GTC	CAT	TCT	GCA	CTT	CTG	GTT	CCA	TAT	TTT	TCA	TGG	AAA	TAT	612
Leu	Ile	Val	His	Ser	Ala	Leu	Leu	Val	Pro	Tyr	Phe	Ser	Trp	Lys	Tyr	
		130					135					140				
AGC	CAT	CGC	CGC	CAC	CAT	TCT	AAC	ATA	GGA	TCT	CTC	GAG	CGA	GAC	GAA	660
Ser	His	Arg	Arg	His	His	Ser	Asn	Ile	Gly	Ser	Leu	Glu	Arg	Asp	Glu	
	145					150					155					
GTG	TTC	GTC	CCG	AAA	TCA	AAG	TCG	AAA	ATT	TCA	TGG	TAT	TCT	AAG	TAC	708
Val	Phe	Val	Pro	Lys	Ser	Lys	Ser	Lys	Ile	Ser	Trp	Tyr	Ser	Lys	Tyr	
160					165					170					175	
TTA	AAC	AAC	CCG	CCA	GGT	CGA	GTT	TTG	ACA	CTT	GCT	GCC	ACG	CTC	CTC	756
Leu	Asn	Asn	Pro	Pro	Gly	Arg	Val	Leu	Thr	Leu	Ala	Ala	Thr	Leu	Leu	
				180					185					190		
CTT	GGC	TGG	CCT	TTA	TAT	TTA	GCT	TTC	AAT	GTC	TCT	GGT	AGA	CCT	TAC	804
Leu	Gly	Trp	Pro	Leu	Tyr	Leu	Ala	Phe	Asn	Val	Ser	Gly	Arg	Pro	Tyr	
			195					200					205			
GAT	CGC	TTT	GCT	TGC	CAT	TAT	GAT	CCC	TAT	GGC	CCA	ATA	TTT	TCC	GAA	852
Asp	Arg	Phe	Ala	Cys	His	Tyr	Asp	Pro	Tyr	Gly	Pro	Ile	Phe	Ser	Glu	
		210					215					220				
AGA	GAA	AGG	CTT	CAG	ATT	TAC	ATT	GCT	GAC	CTC	GGA	ATC	TTT	GCC	ACA	900
Arg	Glu	Arg	Leu	Gln	Ile	Tyr	Ile	Ala	Asp	Leu	Gly	Ile	Phe	Ala	Thr	
	225					230					235					
ACG	TTT	GTG	CTT	TAT	CAG	GCT	ACA	ATG	GCA	AAA	GGG	TTG	GCT	TGG	GTA	948
Thr	Phe	Val	Leu	Tyr	Gln	Ala	Thr	Met	Ala	Lys	Gly	Leu	Ala	Trp	Val	
240					245					250					255	
ATG	CGT	ATC	TAT	GGG	GTG	CCA	TTG	CTT	ATT	GTT	AAC	TGT	TTC	CTT	GTT	996
Met	Arg	Ile	Tyr	Gly	Val	Pro	Leu	Leu	Ile	Val	Asn	Cys	Phe	Leu	Val	
				260					265					270		
ATG	ATC	ACA	TAC	TTG	CAG	CAC	ACT	CAC	CCA	GCT	ATT	CCA	CGC	TAT	GGC	1044
Met	Ile	Thr	Tyr	Leu	Gln	His	Thr	His	Pro	Ala	Ile	Pro	Arg	Tyr	Gly	
			275					280					285			
TCA	TCG	GAA	TGG	GAT	TGG	CTC	CGG	GGA	GCA	ATG	GTG	ACT	GTC	GAT	AGA	1092
Ser	Ser	Glu	Trp	Asp	Trp	Leu	Arg	Gly	Ala	Met	Val	Thr	Val	Asp	Arg	
		290					295					300				

009201" 52E45960

GAT TAT GGG GTG TTG AAT AAA GTA TTC CAT AAC ATT GCA GAC ACT CAT 1140  
 Asp Tyr Gly Val Leu Asn Lys Val Phe His Asn Ile Ala Asp Thr His  
 305 310 315  
 GTA GCT CAT CAT CTC TTT GCT ACA GTG CCA CAT TAC CAT GCA ATG GAG 1188  
 Val Ala His His Leu Phe Ala Thr Val Pro His Tyr His Ala Met Glu  
 320 325 330 335  
 GCC ACT AAA GCA ATC AAG CCT ATA ATG GGT GAG TAT TAC CGG TAT GAT 1236  
 Ala Thr Lys Ala Ile Lys Pro Ile Met Gly Glu Tyr Tyr Arg Tyr Asp  
 340 345 350  
 GGT ACC CCA TTT TAC AAG GCA TTG TGG AGG GAG GCA AAG GAG TGC TTG 1284  
 Gly Thr Pro Phe Tyr Lys Ala Leu Trp Arg Glu Ala Lys Glu Cys Leu  
 355 360 365  
 TTC GTC GAG CCA GAT GAA GGA GCT CCT ACA CAA GGC GTT TTC TGG TAC 1332  
 Phe Val Glu Pro Asp Glu Gly Ala Pro Thr Gln Gly Val Phe Trp Tyr  
 370 375 380  
 CGG AAC AAG TAT TAAAAAAGTG TCATGTAGCC TGCCG 1369  
 Arg Asn Lys Tyr  
 385

## (2) INFORMATION FOR SEQ ID NO:12:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 387 amino acids  
 (B) TYPE: amino acid  
 (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

Met Gly Gly Gly Gly Arg Met Ser Thr Val Ile Ile Ser Asn Asn Ser  
 1 5 10 15  
 Glu Lys Lys Gly Gly Ser Ser His Leu Glu Arg Ala Pro His Thr Lys  
 20 25 30  
 Pro Pro Tyr Thr Leu Gly Asn Leu Lys Arg Ala Ile Pro Pro His Cys  
 35 40 45  
 Phe Glu Arg Ser Phe Val Arg Ser Phe Ser Asn Phe Ala Tyr Asn Phe  
 50 55 60  
 Cys Leu Ser Phe Leu Ser Tyr Ser Ile Ala Thr Asn Phe Phe Pro Tyr  
 65 70 75 80  
 Ile Ser Ser Pro Leu Ser Tyr Val Ala Trp Leu Val Tyr Trp Leu Phe  
 85 90 95  
 Gln Gly Cys Ile Leu Thr Gly Leu Trp Val Ile Gly His Glu Cys Gly  
 100 105 110

[illegible]

## (2) INFORMATION FOR SEQ ID NO:13:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 23 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: cDNA
- (iii) HYPOTHETICAL: NO
- (iv) ANTI-SENSE: NO
- (ix) FEATURE:
  - (A) NAME/KEY: misc\_feature
  - (B) LOCATION: 1..23
  - (D) OTHER INFORMATION: /product=  
"synthetic  
oligonucleotide"
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

TGGGTATGCC AYGANTGYGG NCA

23

## (2) INFORMATION FOR SEQ ID NO:14:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 22 base pairs
  - (B) TYPE: nucleic acid
  - (C) TRANDEDNESS: single
  - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: cDNA
- (iii) HYPOTHETICAL: NO
- (iv) ANTI-SENSE: NO
- (ix) FEATURE:
  - (A) NAME/KEY: misc\_feature
  - (B) LOCATION: 1..22
  - (D) CTHER INFORMATION: /product=  
"synthetic  
oligonucleotide"
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

AAARTGRTGG CACRTGNGTR TC

22

## (2) INFORMATION FOR SEQ ID NO:15:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 2973 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: double
  - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: DNA (genomic)
- (iii) HYPOTHETICAL: NO
- (iv) ANTI-SENSE: NO
- (vi) ORIGINAL SOURCE:
  - (A) ORGANISM: *Arabidopsis thaliana*
- (vii) IMMEDIATE SOURCE:
  - (B) CLONE: pAGF2-6
- (ix) FEATURE:
  - (A) NAME/KEY: exon
  - (B) LOCATION: 433..520
- (ix) FEATURE:
  - (A) NAME/KEY: intron
  - (B) LOCATION: 521..1654
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

```

ATTCGGTAAT TCCTACATAT TTTAGAGATT AGTTTGAGTT TCCATCCATA CTTTACTAGT    60
GATTATAAAT TTAAATACG TACTTTTCGA CTATAAAGTG AAATAAGTA AATTAGAACG    120
TGATATTAAA AAGTTAATGT TCACTGTTAT ATTTTTTTC CAAGTAAAAA ATGGGTTATT    180
TGCGGTAAAT AAAAATACCA GATATTTTGA ATTGATTAAA AAGGTTGAAA TAAGAGAGGA    240
GGGGAAAGAA AAGAAGGTGG GGGCCAGTA TGAAAGGGAA AGGTGTCATC AAATCATCTC    300
TCTCTCTCTC TACCTTCGAC CCACGGGCCG TGTCCATTTA AAGCCCTGTC TCTTGCCATT    360
CCCCATCTGA CCACCAGAAG AAGAGCCACA CACTCACAAA TAAAAAGAG AGAGAGAGAG    420
AGAGAGACAG AGAGAGAGAG AGATTCTGCG GAGGAGCTTC TTCTTCGTAG GGTGTTTCATC    480
GTTATTAACG TTATCGCCCC TACGTCAGCT CCATCTCCAG GTCCGTCGCT TCTCTTCCAT    540
TTCTTCTCAT TTTCGATTTT GATTCTTATT TCTTCCAGT AGCTCCTGCT CTGTGAATTT    600
CTCCGCTCAC GATAGATCTG CTTATACTCC TTACATTCAA CCTTAGATCT GGTCTCGATT    660
CTCTGTTTCT CTGTTTTTTT CTTTTGGTCG AGAATCTGAT GTTTGTTTAT GTTCTGTCAC    720
CATTAAATAAT GATGAACTCT CTCATTCTA CAATGATTAG TTTCTCTCGT CTACCAAACG    780

```

ATATGTTGCA TTTTCACTTT TCTTCTTTTT TTCTAAGATG ATTTGCTTTG ACCAATTTGT 840  
 TTAGATCTTT ATTTTATTTT ATTTTCTGGT GGGTTGGTGG AAATTGAAAA AAAAAAAAAA 900  
 AAAAGCATAA ATTGTTATTT GTTAATGTAT TCATTTTTTG GCTATTTGTT CTGGGTAAAA 960  
 ATCTGCTTCT ACTGTTGAAT CTTTCCTGGA TTTTTTACTC CTATTGGGTT TTTATAGTAA 1020  
 AAATACATAA TAAAAGGAAA ACAAAGTTT TATAGATTCT CTTAAACCCC TTACGATAAA 1080  
 AGTTGGAATC AAAATAATTC AGGATCAGAT GCTCTTTGAT TGATTCAGAT GCGATTACAG 1140  
 TTGCATGGAA AATTTTCTAG ATCCGTCGTC ACATTTTATT TTCTGTTTAA ATATCTAAAT 1200  
 CTGATATATG ATGTCGACAA ATTCTGGTGG CTTATACATC ACTTCAACTG TTTTCTTTTG 1260  
 GCTTTGTTTG TCAACTTGGT TTTCAATACG ATTTGTGATT TCGATCGCTG AATTTTAAAT 1320  
 ACAAGCAAAC TGATGTTAAC CACAAGCAAG AGATGTGACC TGCCTTATTA ACATCGTATT 1380  
 ACTTACTACT AGTCGTATTC TCAACGCAAT CGTTTTTGTA TTTCTCACAT TATGCCGCTT 1440  
 CTCTACTCTT TATTCCTTTT GGTCCACGCA TTTTCTATTT GTGGCAATCC CTTTCACAAC 1500  
 CTGATTTCCC ACTTTGGATC ATTTGTCTGA AGACTCTCTT GAATCGTTAC CACTTGTTTC 1560  
 TTGTGCATGC TCTGTTTTTT AGAATTAATG ATAAACTAT TCCATAGTCT TGAGTTTTCA 1620  
 GCTTGTTGAT TCTTTTGCTT TTGGTTTTCT GCAGAAACAT GGGTGCAGGT GGAAGAATGC 1680  
 CGGTTCCCTAC TTCTTCCAAG AAATCGGAAA CCGACACCAC AAAGCGTGTG CCGTGCAGAGA 1740  
 AACCGCCTTT CTCGGTGGGA GATCTGAAGA AAGCAATCCC GCCGCATTGT TTCAAACGCT 1800  
 CAATCCCTCG CTCTTCTCC TACCTTATCA GTGACATCAT TATAGCCTCA TGCTTCTACT 1860  
 ACGTCGCCAC CAATTACTTC TCTCTCCTCC CTCAGCCTCT CTCTTACTTG GCTTGGCCAC 1920  
 TCTATTGGGC CTGTCAAGGC TGTGTCTTAA CTGGTATCTG GGTATAGCC CACGAATGCG 1980  
 GTCACCACGC ATTCAGCGAC TACCAATGGC TGGATGACAC AGTTGGTCTT ATCTTCCATT 2040  
 CCTTCCTCCT CGTCCCTTAC TTCTCCTGGA AGTATAGTCA TCGCCGTCAC CATTCCAACA 2100  
 CTGGATCCCT CGAAAGAGAT GAAGTATTTG TCCCAAAGCA GAAATCAGCA ATCAAGTGGT 2160  
 ACGGGAAATA CCTCAACAAC CCTCTTGGAC GCATCATGAT GTTAACCGTC CAGTTTGTCC 2220  
 TCGGGTGGCC CTTGTACTTA GCCTTTAACG TCTCTGGCAG ACCGTATGAC GGGTTCGCTT 2280  
 GCCATTTCTT CCCCACGCT CCCATCTACA ATGACCGAGA ACGCCTCCAG ATATACCTCT 2340  
 CTGATGCGGG TATTCTAGCC GTCTGTTTTG GTCTTTACCG TTACGCTGCT GCACAAGGGA 2400  
 TGGCCTCGAT GATCTGCCTC TACGGAGTAC CGCTTCTGAT AGTGAATGCG TTCCTCGTCT 2460  
 TGATCACTTA CTTGCAGCAC ACTCATCCCT CGTTGCCTCA CTACGATTCA TCAGAGTGGG 2520

ACTGGCTCAG GGGAGCTTTG GCTACCGTAG ACAGAGACTA CGGAATCTTG AACAAAGGTGT 2580  
 TCCACAACAT TACAGACACA CACGTGGCTC ATCACCTGTT CTCGACAATG CCGCATTATA 2640  
 ACGCAATGGA AGCTACAAAG GCGATAAAGC CAATTCTGGG AGACTATTAC CAGTTCGATG 2700  
 GAACACCGTG GTATGTGGCG ATGTATAGGG AGGCAAAGGA GTGTATCTAT GTAGAACCGG 2760  
 ACAGGGAAGG TGACAAGAAA GGTGTGTACT GGTACAACAA TAAGTTATGA GGATGATGGT 2820  
 GAAGAAATTG TCGACTTTTC TCTTGTCTGT TTGTCTTTTG TTAAAGAAGC TATGCTTCGT 2880  
 TTTAATAATC TTATTGTCCA TTTTGTGTGT TTATGACATT TTGGCTGCTC ATTATGTTAT 2940  
 GTGGGAAGTT AGCGTTCAAA TGTTTTGGGT CGG 2973

(2) INFORMATION FOR SEQ ID NO:16:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 23 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(ix) FEATURE:

- (A) NAME/KEY: misc\_feature
- (B) LOCATION: 1..23
- (D) OTHER INFORMATION: /product=  
"synthetic  
oligonucleotide"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

GGGCATGTNG ARAANARRTG RTG 23

(2) INFORMATION FOR SEQ ID NO:17:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 23 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

009201"5275950

(iv) ANTI-SENSE: NO

(ix) FEATURE:

(A) NAME/KEY: misc feature

(B) LOCATION: 1..23

(D) OTHER INFORMATION: /product=  
"synthetic  
oligonucleotide"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

GGGCATGTRC TRAANARRTG RTG

23

003201" 6/6/6950



CLAIMS

1. An isolated nucleic acid fragment comprising a nucleic acid sequence encoding a fatty acid desaturase or a fatty acid desaturase-related enzyme with an amino acid identity of 50% or greater to the polypeptide encoded by SEQ ID NOS:1, 3, 5, 7, 9, 11, or 15.
2. The isolated nucleic acid fragment of Claim 1 wherein the amino acid identity is 60% or greater to the polypeptide encoded by SEQ ID NOS:1, 3, 5, 7, 9, 11, or 15.
3. The isolated nucleic acid fragment of Claim 1 wherein the nucleic acid identity is 90% or greater to SEQ ID NOS:1, 3, 5, 7, 9, 11, or 15.
4. The isolated nucleic acid fragment of Claim 1 wherein said fragment is isolated from an oil-producing plant species.
5. An isolated nucleic acid fragment comprising a nucleic acid sequence encoding a delta-12 fatty acid hydroxylase.
6. A chimeric gene capable of causing altered levels of ricinoleic acid in a transformed plant cell, said chimeric gene comprising a nucleic acid fragment of Claim 5, said fragment operably linked to suitable regulatory sequences.
7. A chimeric gene capable of causing altered levels of fatty acids in a transformed plant cell, said chimeric gene comprising a nucleic acid fragment of any of Claims 1, 2, 3, said fragment operably linked to suitable regulatory sequences.
8. Plants containing a chimeric gene of Claim 6 or Claim 7.
9. Oil obtained from seeds of the plants of Claim 8.
10. A method of producing seed oil containing altered levels of unsaturated fatty acids comprising:

(a) transforming a plant cell of an oil-producing species with a chimeric gene of Claim 5;

(b) growing fertile plants from the transformed plant cells of step (a);

5 (c) screening progeny seeds from the fertile plants of step (b) for the desired levels of unsaturated fatty acids; and

(d) processing the progeny seed of step (c) to obtain seed oil containing altered levels of  
10 unsaturated fatty acids.

11. A method of molecular breeding to obtain altered levels of a fatty acid in seed oil of oil-producing plant species comprising:

(a) making a cross between two varieities of  
15 oil-producing species differing in the fatty acid trait;

(b) making a Southern blot of restriction enzyme digested genomic DNA isolated from several progeny plants resulting from the cross of step (a); and

(c) hybridizing the Southern blot with the  
20 radiolabelled nucleic acid fragment of Claim 1.

12. A method of RFLP mapping comprising:

(a) making a cross between two varieties of plants;

(b) making a Southern blot of restriction  
25 enzyme digested genomic DNA isolated from several progeny plants resulting from the cross of step (a); and

(c) hybridizing the Southern blot with the radiolabelled nucleic acid fragments of Claim 1.

13. A method to isolate nucleic acid fragments  
30 encoding fatty acid desaturases and related enzymes, comprising:

(a) comparing SEQ ID NOS:2, 4, 6, 8, 10, or 12 and other fatty acid desaturase polypeptide sequences;

0969774-102600

(b) identifying the conserved sequences of 4 or more amino acids obtained in step a;

(c) designing degenerate oligomers based on the conserved sequences identified in step b; and

5 (d) using the degenerate oligomers of step c to isolate sequences encoding fatty acid desaturases and desaturase-related enzymes by sequence dependent protocols.

14. An isolated nucleic acid fragment of Claim 1  
10 comprising a nucleic acid sequence encoding a plant microsomal delta-12 fatty acid desaturase.

15. A method for altering fatty acids composition in seeds comprising:

(a) making a cross between a mutant line with  
15 altered fatty acid composition with a plant containing the chimeric gene of Claim 7;

(b) growing fertile plants from seeds obtained from the cross; and

(c) screening progeny seeds from the fertile  
20 plants of step (b) for seeds containing altered fatty acid levels.

16. A method for reducing polyunsaturated fatty acids in rapeseed oil comprising:

(a) making a cross between a rapeseed variety  
25 with increased oleic acid content or reduced linolenic acid content with a plant containing the chimeric gene of Claim 7;

(b) growing fertile plants from seeds obtained from the cross; and

(c) screening progeny seeds from the fertile  
30 plants of step (b) for seeds containing reduced polyunsaturated fatty acids.

17. The method of Claim 16 wherein the cross in  
(a) is between a progeny plant derived from a seed  
35 comprising the Brassica variety having an oleic acid

content of about 69% to 77%, based upon total extractable oil and belonging to a line in which the said oleic acid content has been stabilized for both the generation to which the seed belongs and its parent  
5 generation.

18. A method for reducing saturated fatty acids in rapeseed seeds comprising:

- (a) making a cross between a rapeseed variety with increased oleic acid content with a plant  
10 containing the chimeric gene of Claim 7;
- (b) growing fertile plants from seeds obtained from the cross; and
- (c) screening progeny seeds from the fertile plants of step (b) for seeds containing reduced  
15 saturated fatty acids.

19. A method for reducing polyunsaturated fatty acids in soybean oil comprising:

- (a) making a cross between a soybean variety with increased oleic acid content or reduced linolenic  
20 acid content with a plant containing the chimeric gene of Claim 7;
- (b) growing fertile plants from seeds obtained from the cross; and
- (c) screening progeny seeds from the fertile  
25 plants of step (b) for seeds containing reduced polyunsaturated fatty acids.

20. A method for reducing saturated fatty acids in soybean seeds comprising:

- (a) making a cross between a soybean variety  
30 with increased oleic acid content with a plant containing the chimeric gene of Claim 7;
- (b) growing fertile plants from seeds obtained from the cross; and

00920T 5265950

(c) screening progeny seeds from the fertile plants of step (b) for seeds containing reduced saturated fatty acids.

21. A *Brassica* sp. plant with seed palmitic acid  
5 of 2.7% or lower of total fatty acid.

22. A *Brassica* sp. plant with seed stearic acid of 1.1% or lower of total fatty acid.

23. A *Brassica* sp. plant with a combined seed  
10 palmitic acid and stearic acid content of 3.9% or lower of total fatty acids.

24. A soybean plant with seed palmitic acid of 6.7% or lower of total fatty acid.

25. A soybean plant with seed stearic acid of 2.1% or lower of total fatty acid.

15 26. A soybean plant with a combined seed palmitic acid and stearic acid content of 9.2% or lower total of fatty acids.

27. Oil obtained from the plants of Claims 21-26.

24 21/2

TITLE

GENES FOR MICROSOMAL DELTA-12 FATTY ACID  
DESATURASES AND RELATED ENZYMES FROM PLANTS

ABSTRACT OF THE DISCLOSURE

5       The preparation and use of nucleic acid fragments  
encoding fatty acid desaturase enzymes are described.  
The invention permits alteration of plant lipid  
composition. Chimeric genes incorporating such nucleic  
acid fragments with suitable regulatory sequences may be  
10       used to create transgenic plants with altered levels of  
unsaturated fatty acids.

15

20

25

30

35   BCS/bjm

009207 626369

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i) APPLICANT: JONATHAN EDWARD LIGHTNER  
JOHN JOSEPH OKULEY

(ii) TITLE OF INVENTION: GENES FOR MICROSOMAL  
FATTY ACID DELTA-12  
DESATURASES AND RELATED  
ENZYMES FROM PLANTS

(iii) NUMBER OF SEQUENCES: 17

(iv) CORRESPONDENCE ADDRESS:

(A) ADDRESSEE: E. I. DU PONT DE NEMOURS  
AND COMPANY

(B) STREET: 1007 MARKET STREET

(C) CITY: WILMINGTON

(D) STATE: DELAWARE

(E) COUNTRY: U.S.A.

(F) ZIP: 19898

(v) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk

(B) COMPUTER: MacIntosh

(C) OPERATING SYSTEM: Microsoft Windows 95

(D) SOFTWARE: Microsoft Word Version 7.0

(vi) CURRENT APPLICATION DATA:

(A) APPLICATION NUMBER:

(B) FILING DATE:

(C) CLASSIFICATION:

(vii) PRIOR APPLICATION DATA:

(A) APPLICATION NUMBER: U.S. 07/977,339

(B) FILING DATE: 17-NOV-1992

(A) APPLICATION NUMBER: U.S. 09/133,962

(B) FILING DATE: 14-AUG-1998

(viii) ATTORNEY/AGENT INFORMATION:

(A) NAME: Christenbury, Lynne M..

(B) REGISTRATION NUMBER: 30,971

(C) REFERENCE/DOCKET NUMBER: BB1043 US DIV

(ix) TELECOMMUNICATION INFORMATION:

(A) TELEPHONE: (302) 992-5481

(B) TELEFAX: (302) 892-1026

(C) TELEX: 835420

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 1372 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: double  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:  
 (A) ORGANISM: *Arabidopsis thaliana*

(vii) IMMEDIATE SOURCE:  
 (B) CLONE: p92103

(ix) FEATURE:  
 (A) NAME/KEY: CDS  
 (B) LOCATION: 93..1244

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

AGAGAGAGAG ATTCTGCGGA GGAGCTTCTT CTTCGTAGGG TGTTTCATCGT TATTAACGTT  
 60

ATCGCCCCTA CGTCAGCTCC ATCTCCAGAA AC ATG GGT GCA GGT GGA AGA ATG  
 113

Met Gly Ala Gly Gly Arg Met  
 1 5

CCG GTT CCT ACT TCT TCC AAG AAA TCG GAA ACC GAC ACC ACA AAG CGT  
 161

Pro Val Pro Thr Ser Ser Lys Lys Ser Glu Thr Asp Thr Thr Lys Arg  
 10 15 20

GTG CCG TGC GAG AAA CCG CCT TTC TCG GTG GGA GAT CTG AAG AAA GCA  
 209

Val Pro Cys Glu Lys Pro Pro Phe Ser Val Gly Asp Leu Lys Lys Ala  
 25 30 35

ATC CCG CCG CAT TGT TTC AAA CGC TCA ATC CCT CGC TCT TTC TCC TAC  
 257

Ile Pro Pro His Cys Phe Lys Arg Ser Ile Pro Arg Ser Phe Ser Tyr  
 40 45 50 55

CTT ATC AGT GAC ATC ATT ATA GCC TCA TGC TTC TAC TAC GTC GCC ACC  
 305

Leu Ile Ser Asp Ile Ile Ile Ala Ser Cys Phe Tyr Tyr Val Ala Thr  
 60 65 70

AAT TAC TTC TCT CTC CTC CCT CAG CCT CTC TCT TAC TTG GCT TGG CCA  
 353

Asn Tyr Phe Ser Leu Leu Pro Gln Pro Leu Ser Tyr Leu Ala Trp Pro  
 75 80 85

CTC TAT TGG GCC TGT CAA GGC TGT GTC CTA ACT GGT ATC TGG GTC ATA  
 401



```

Leu Tyr Trp Ala Cys Gln Gly Cys Val Leu Thr Gly Ile Trp Val Ile
    90                                95                        100

GCC CAC GAA TGC GGT CAC CAC GCA TTC AGC GAC TAC CAA TGG CTG GAT
449
Ala His Glu Cys Gly His His Ala Phe Ser Asp Tyr Gln Trp Leu Asp
    105                                110                        115

GAC ACA GTT GGT CTT ATC TTC CAT TCC TTC CTC CTC GTC CCT TAC TTC
497
Asp Thr Val Gly Leu Ile Phe His Ser Phe Leu Leu Val Pro Tyr Phe
120                                125                        130                        135

TCC TGG AAG TAT AGT CAT CGC CGT CAC CAT TCC AAC ACT GGA TCC CTC
545
Ser Trp Lys Tyr Ser His Arg Arg His His Ser Asn Thr Gly Ser Leu
    140                                145                        150

GAA AGA GAT GAA GTA TTT GTC CCA AAG CAG AAA TCA GCA ATC AAG TGG
593
Glu Arg Asp Glu Val Phe Val Pro Lys Gln Lys Ser Ala Ile Lys Trp
    155                                160                        165

TAC GGG AAA TAC CTC AAC AAC CCT CTT GGA CGC ATC ATG ATG TTA ACC
641
Tyr Gly Lys Tyr Leu Asn Asn Pro Leu Gly Arg Ile Met Met Leu Thr
    170                                175                        180

GTC CAG TTT GTC CTC GGG TGG CCC TTG TAC TTA GCC TTT AAC GTC TCT
689
Val Gln Phe Val Leu Gly Trp Pro Leu Tyr Leu Ala Phe Asn Val Ser
    185                                190                        195

GGC AGA CCG TAT GAC GGG TTC GCT TGC CAT TTC TTC CCC AAC GCT CCC
737
Gly Arg Pro Tyr Asp Gly Phe Ala Cys His Phe Phe Pro Asn Ala Pro
200                                205                        210                        215

ATC TAC AAT GAC CGA GAA CGC CTC CAG ATA TAC CTC TCT GAT GCG GGT
785
Ile Tyr Asn Asp Arg Glu Arg Leu Gln Ile Tyr Leu Ser Asp Ala Gly
    220                                225                        230

ATT CTA GCC GTC TGT TTT GGT CTT TAC CGT TAC GCT GCT GCA CAA GGG
833
Ile Leu Ala Val Cys Phe Gly Leu Tyr Arg Tyr Ala Ala Ala Gln Gly
    235                                240                        245

ATG GCC TCG ATG ATC TGC CTC TAC GGA GTA CCG CTT CTG ATA GTG AAT
881
Met Ala Ser Met Ile Cys Leu Tyr Gly Val Pro Leu Leu Ile Val Asn
    250                                255                        260

GCG TTC CTC GTC TTG ATC ACT TAC TTG CAG CAC ACT CAT CCC TCG TTG
929
Ala Phe Leu Val Leu Ile Thr Tyr Leu Gln His Thr His Pro Ser Leu

```

265                      270                      275  
 CCT CAC TAC GAT TCA TCA GAG TGG GAC TGG CTC AGG GGA GCT TTG GCT  
 977  
 Pro His Tyr Asp Ser Ser Glu Trp Asp Trp Leu Arg Gly Ala Leu Ala  
 280                      285                      290                      295  
  
 ACC GTA GAC AGA GAC TAC GGA ATC TTG AAC AAG GTG TTC CAC AAC ATT  
 1025  
 Thr Val Asp Arg Asp Tyr Gly Ile Leu Asn Lys Val Phe His Asn Ile  
                     300                      305                      310  
  
 ACA GAC ACA CAC GTG GCT CAT CAC CTG TTC TCG ACA ATG CCG CAT TAT  
 1073  
 Thr Asp Thr His Val Ala His His Leu Phe Ser Thr Met Pro His Tyr  
                     315                      320                      325  
  
 AAC GCA ATG GAA GCT ACA AAG GCG ATA AAG CCA ATT CTG GGA GAC TAT  
 1121  
 Asn Ala Met Glu Ala Thr Lys Ala Ile Lys Pro Ile Leu Gly Asp Tyr  
                     330                      335                      340  
  
 TAC CAG TTC GAT GGA ACA CCG TGG TAT GTA GCG ATG TAT AGG GAG GCA  
 1169  
 Tyr Gln Phe Asp Gly Thr Pro Trp Tyr Val Ala Met Tyr Arg Glu Ala  
                     345                      350                      355  
  
 AAG GAG TGT ATC TAT GTA GAA CCG GAC AGG GAA GGT GAC AAG AAA GGT  
 1217  
 Lys Glu Cys Ile Tyr Val Glu Pro Asp Arg Glu Gly Asp Lys Lys Gly  
 360                      365                      370                      375  
  
 GTG TAC TGG TAC AAC AAT AAG TTA TGAGCATGAT GGTGAAGAAA TTGTCGACCT  
 1271  
 Val Tyr Trp Tyr Asn Asn Lys Leu  
                     380  
  
 TTCTCTTGTC TGTTTGTCTT TTGTTAAAGA AGCTATGCTT CGTTTTAATA ATCTTATTGT  
 1331  
  
 CCATTTTGTT GTGTTATGAC ATTTTGGCTG CTCATTATGT T  
 1372

## (2) INFORMATION FOR SEQ ID NO:2:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 383 amino acids  
 (B) TYPE: amino acid  
 (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Met Gly Ala Gly Gly Arg Met Pro Val Pro Thr Ser Ser Lys Lys Ser  
 1                      5                      10                      15

```

Glu Thr Asp Thr Thr Lys Arg Val Pro Cys Glu Lys Pro Pro Phe Ser
      20                      25                      30
Val Gly Asp Leu Lys Lys Ala Ile Pro Pro His Cys Phe Lys Arg Ser
      35                      40                      45
Ile Pro Arg Ser Phe Ser Tyr Leu Ile Ser Asp Ile Ile Ile Ala Ser
      50                      55                      60
Cys Phe Tyr Tyr Val Ala Thr Asn Tyr Phe Ser Leu Leu Pro Gln Pro
      65                      70                      75                      80
Leu Ser Tyr Leu Ala Trp Pro Leu Tyr Trp Ala Cys Gln Gly Cys Val
      85                      90                      95
Leu Thr Gly Ile Trp Val Ile Ala His Glu Cys Gly His His Ala Phe
      100                      105                      110
Ser Asp Tyr Gln Trp Leu Asp Asp Thr Val Gly Leu Ile Phe His Ser
      115                      120                      125
Phe Leu Leu Val Pro Tyr Phe Ser Trp Lys Tyr Ser His Arg Arg His
      130                      135                      140
His Ser Asn Thr Gly Ser Leu Glu Arg Asp Glu Val Phe Val Pro Lys
      145                      150                      155                      160
Gln Lys Ser Ala Ile Lys Trp Tyr Gly Lys Tyr Leu Asn Asn Pro Leu
      165                      170                      175
Gly Arg Ile Met Met Leu Thr Val Gln Phe Val Leu Gly Trp Pro Leu
      180                      185                      190
Tyr Leu Ala Phe Asn Val Ser Gly Arg Pro Tyr Asp Gly Phe Ala Cys
      195                      200                      205
His Phe Phe Pro Asn Ala Pro Ile Tyr Asn Asp Arg Glu Arg Leu Gln
      210                      215                      220
Ile Tyr Leu Ser Asp Ala Gly Ile Leu Ala Val Cys Phe Gly Leu Tyr
      225                      230                      235                      240
Arg Tyr Ala Ala Ala Gln Gly Met Ala Ser Met Ile Cys Leu Tyr Gly
      245                      250                      255
Val Pro Leu Leu Ile Val Asn Ala Phe Leu Val Leu Ile Thr Tyr Leu
      260                      265                      270
Gln His Thr His Pro Ser Leu Pro His Tyr Asp Ser Ser Glu Trp Asp
      275                      280                      285
Trp Leu Arg Gly Ala Leu Ala Thr Val Asp Arg Asp Tyr Gly Ile Leu
      290                      295                      300
Asn Lys Val Phe His Asn Ile Thr Asp Thr His Val Ala His His Leu

```

```

305          310          315          320
Phe Ser Thr Met Pro His Tyr Asn Ala Met Glu Ala Thr Lys Ala Ile
          325          330          335
Lys Pro Ile Leu Gly Asp Tyr Tyr Gln Phe Asp Gly Thr Pro Trp Tyr
          340          345          350
Val Ala Met Tyr Arg Glu Ala Lys Glu Cys Ile Tyr Val Glu Pro Asp
          355          360          365
Arg Glu Gly Asp Lys Lys Gly Val Tyr Trp Tyr Asn Asn Lys Leu
          370          375          380

```

## (2) INFORMATION FOR SEQ ID NO:3:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1426 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: cDNA to mRNA

## (iii) HYPOTHETICAL: NO

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Brassica napus

## (ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 130..1284

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

```

GGCACGAGCT CGTGCCGAAT TCGGCACGAG AGGAGACAGA GAGAGAGTTT GAGGAGGAGC 60
TTCTTCGTAG GGTTTCATCGT TATTAACGTT AAATCTTCAT CCCCCCTAC GTCAGCCAGC 120
TCAAGAAAC ATG GGT GCA GGT GGA AGA ATG CAA GTG TCT CCT CCC TCC 168
      Met Gly Ala Gly Gly Arg Met Gln Val Ser Pro Pro Ser
        1          5          10
AAA AAG TCT GAA ACC GAC AAC ATC AAG CGC GTA CCC TGC GAG ACA CCG 216
Lys Lys Ser Glu Thr Asp Asn Ile Lys Arg Val Pro Cys Glu Thr Pro
      15          20          25
CCC TTC ACT GTC GGA GAA CTC AAG AAA GCA ATC CCA CCG CAC TGT TTC 264
Pro Phe Thr Val Gly Glu Leu Lys Lys Ala Ile Pro Pro His Cys Phe
      30          35          40
AAA CGC TCG ATC CCT CGC TCT TTC TCC TAC CTC ATC TGG GAC ATC ATC 312
Lys Arg Ser Ile Pro Arg Ser Phe Ser Tyr Leu Ile Trp Asp Ile Ile
          50          55          60
ATA GCC TCC TGC TTC TAC TAC GTC GCC ACC ACT TAC TTC CCT CTC CTC 360

```

Ile	Ala	Ser	Cys	Phe	Tyr	Tyr	Val	Ala	Thr	Thr	Tyr	Phe	Pro	Leu	Leu	
			65					70					75			
CCT	CAC	CCT	CTC	TCC	TAC	TTC	GCC	TGG	CCT	CTC	TAC	TGG	GCC	TGC	CAG	408
Pro	His	Pro	Leu	Ser	Tyr	Phe	Ala	Trp	Pro	Leu	Tyr	Trp	Ala	Cys	Gln	
		80					85					90				
GGC	TGC	GTC	CTA	ACC	GGC	GTC	TGG	GTC	ATA	GCC	CAC	GAG	TGC	GGC	CAC	456
Gly	Cys	Val	Leu	Thr	Gly	Val	Trp	Val	Ile	Ala	His	Glu	Cys	Gly	His	
	95					100					105					
CAC	GCC	TTC	AGC	GAC	TAC	CAG	TGG	CTG	GAC	GAC	ACC	GTC	GGC	CTC	ATC	504
His	Ala	Phe	Ser	Asp	Tyr	Gln	Trp	Leu	Asp	Asp	Thr	Val	Gly	Leu	Ile	
110					115					120					125	
TTC	CAC	TCC	TTC	CTC	CTC	GTC	CCT	TAC	TTC	TCC	TGG	AAG	TAC	AGT	CAT	552
Phe	His	Ser	Phe	Leu	Leu	Val	Pro	Tyr	Phe	Ser	Trp	Lys	Tyr	Ser	His	
				130					135					140		
CGA	CGC	CAC	CAT	TCC	AAC	ACT	GGC	TCC	CTC	GAG	AGA	GAC	GAA	GTG	TTT	600
Arg	Arg	His	His	Ser	Asn	Thr	Gly	Ser	Leu	Glu	Arg	Asp	Glu	Val	Phe	
			145					150					155			
GTC	CCC	AAG	AAG	AAG	TCA	GAC	ATC	AAG	TGG	TAC	GGC	AAG	TAC	CTC	AAC	648
Val	Pro	Lys	Lys	Lys	Ser	Asp	Ile	Lys	Trp	Tyr	Gly	Lys	Tyr	Leu	Asn	
		160					165					170				
AAC	CCT	TTG	GGA	CGC	ACC	GTG	ATG	TTA	ACG	GTT	CAG	TTC	ACT	CTC	GGC	696
Asn	Pro	Leu	Gly	Arg	Thr	Val	Met	Leu	Thr	Val	Gln	Phe	Thr	Leu	Gly	
	175					180					185					
TGG	CCT	TTG	TAC	TTA	GCC	TTC	AAC	GTC	TCG	GGG	AGA	CCT	TAC	GAC	GGC	744
Trp	Pro	Leu	Tyr	Leu	Ala	Phe	Asn	Val	Ser	Gly	Arg	Pro	Tyr	Asp	Gly	
190					195					200					205	
GGC	TTC	GCT	TGC	CAT	TTC	CAC	CCC	AAC	GCT	CCC	ATC	TAC	AAC	GAC	CGT	792
Gly	Phe	Ala	Cys	His	Phe	His	Pro	Asn	Ala	Pro	Ile	Tyr	Asn	Asp	Arg	
				210					215					220		
GAG	CGT	CTC	CAG	ATA	TAC	ATC	TCC	GAC	GCT	GGC	ATC	CTC	GCC	GTC	TGC	840
Glu	Arg	Leu	Gln	Ile	Tyr	Ile	Ser	Asp	Ala	Gly	Ile	Leu	Ala	Val	Cys	
			225					230					235			
TAC	GGT	CTC	TAC	CGC	TAC	GCT	GCT	GTC	CAA	GGA	GTT	GCC	TCG	ATG	GTC	888
Tyr	Gly	Leu	Tyr	Arg	Tyr	Ala	Ala	Val	Gln	Gly	Val	Ala	Ser	Met	Val	
		240				245						250				
TGC	TTC	TAC	GGA	GTT	CCT	CTT	CTG	ATT	GTC	AAC	GGG	TTC	TTA	GTT	TTG	936
Cys	Phe	Tyr	Gly	Val	Pro	Leu	Leu	Ile	Val	Asn	Gly	Phe	Leu	Val	Leu	
	255					260					265					
ATC	ACT	TAC	TTG	CAG	CAC	ACG	CAT	CCT	TCC	CTG	CCT	CAC	TAT	GAC	TCG	984
Ile	Thr	Tyr	Leu	Gln	His	Thr	His	Pro	Ser	Leu	Pro	His	Tyr	Asp	Ser	
270					275					280					285	
TCT	GAG	TGG	GAT	TGG	TTG	AGG	GGA	GCT	TTG	GCC	ACC	GTT	GAC	AGA	GAC	1032

Ser Glu Trp Asp Trp Leu Arg Gly Ala Leu Ala Thr Val Asp Arg Asp  
 290 295 300  
 TAC GGA ATC TTG AAC AAG GTC TTC CAC AAT ATC ACG GAC ACG CAC GTG 1080  
 Tyr Gly Ile Leu Asn Lys Val Phe His Asn Ile Thr Asp Thr His Val  
 305 310 315  
 GCG CAT CAC CTG TTC TCG ACC ATG CCG CAT TAT CAT GCG ATG GAA GCT 1128  
 Ala His His Leu Phe Ser Thr Met Pro His Tyr His Ala Met Glu Ala  
 320 325 330  
 ACG AAG GCG ATA AAG CCG ATA CTG GGA GAG TAT TAT CAG TTC GAT GGG 1176  
 Thr Lys Ala Ile Lys Pro Ile Leu Gly Glu Tyr Tyr Gln Phe Asp Gly  
 335 340 345  
 ACG CCG GTG GTT AAG GCG ATG TGG AGG GAG GCG AAG GAG TGT ATC TAT 12  
 24  
 Thr Pro Val Val Lys Ala Met Trp Arg Glu Ala Lys Glu Cys Ile Tyr  
 350 355 360 365  
 GTG GAA CCG GAC AGG CAA GGT GAG AAG AAA GGT GTG TTC TGG TAC AAC 12  
 72  
 Val Glu Pro Asp Arg Gln Gly Glu Lys Lys Gly Val Phe Trp Tyr Asn  
 370 375 380  
 AAT AAG TTA TGA AGCAAAGAAG AACTGAACC TTTCTCTTCT ATGATTGTCT 13  
 24  
 Asn Lys Leu  
 TTGTTTAAGA AGCTATGTTT CTGTTTCAAT AATCTTAATT ATCCATTTTG TTGTGTTTTT 13  
 84  
 TGACATTTTG GCTAAAATTA TGTGATGTTG GAAGTTAGTG TC 14  
 26

## (2) INFORMATION FOR SEQ ID NO:4:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 384 amino acids  
 (B) TYPE: amino acid  
 (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

Met Gly Ala Gly Gly Arg Met Gln Val Ser Pro Pro Ser Lys Lys Ser  
 1 5 10  
 Glu Thr Asp Asn Ile Lys Arg Val Pro Cys Glu Thr Pro Pro Phe Thr  
 20 25 30  
 Val Gly Glu Leu Lys Lys Ala Ile Pro Pro His Cys Phe Lys Arg Ser  
 35 40 45

Ile Pro Arg Ser Phe Ser Tyr Leu Ile Trp Asp Ile Ile Ile Ala Ser  
 50 55 60  
 Cys Phe Tyr Tyr Val Ala Thr Thr Tyr Phe Pro Leu Leu Pro His Pro  
 65 70 75 80  
 Leu Ser Tyr Phe Ala Trp Pro Leu Tyr Trp Ala Cys Gln Gly Cys Val  
 85 90 95  
 Leu Thr Gly Val Trp Val Ile Ala His Glu Cys Gly His His Ala Phe  
 100 105 110  
 Ser Asp Tyr Gln Trp Leu Asp Asp Thr Val Gly Leu Ile Phe His Ser  
 115 120 125  
 Phe Leu Leu Val Pro Tyr Phe Ser Trp Lys Tyr Ser His Arg Arg His  
 130 135 140  
 His Ser Asn Thr Gly Ser Leu Glu Arg Asp Glu Val Phe Val Pro Lys  
 145 150 155 160  
 Lys Lys Ser Asp Ile Lys Trp Tyr Gly Lys Tyr Leu Asn Asn Pro Leu  
 165 170 175  
 Gly Arg Thr Val Met Leu Thr Val Gln Phe Thr Leu Gly Trp Pro Leu  
 180 185 190  
 Tyr Leu Ala Phe Asn Val Ser Gly Arg Pro Tyr Asp Gly Gly Phe Ala  
 195 200 205  
 Cys His Phe His Pro Asn Ala Pro Ile Tyr Asn Asp Arg Glu Arg Leu  
 210 215 220  
 Gln Ile Tyr Ile Ser Asp Ala Gly Ile Leu Ala Val Cys Tyr Gly Leu  
 225 230 235 240  
 Tyr Arg Tyr Ala Ala Val Gln Gly Val Ala Ser Met Val Cys Phe Tyr  
 245 250 255  
 Gly Val Pro Leu Leu Ile Val Asn Gly Phe Leu Val Leu Ile Thr Tyr  
 260 265 270  
 Leu Gln His Thr His Pro Ser Leu Pro His Tyr Asp Ser Ser Glu Trp  
 275 280 285  
 Asp Trp Leu Arg Gly Ala Leu Ala Thr Val Asp Arg Asp Tyr Gly Ile  
 290 295 300  
 Leu Asn Lys Val Phe His Asn Ile Thr Asp Thr His Val Ala His His  
 305 310 315 320  
 Leu Phe Ser Thr Met Pro His Tyr His Ala Met Glu Ala Thr Lys Ala  
 325 330 335  
 Ile Lys Pro Ile Leu Gly Glu Tyr Tyr Gln Phe Asp Gly Thr Pro Val  
 340 345 350

Val Lys Ala Met Trp Arg Glu Ala Lys Glu Cys Ile Tyr Val Glu Pro  
 355 360 365  
 Asp Arg Gln Gly Glu Lys Lys Gly Val Phe Trp Tyr Asn Asn Lys Leu  
 370 375 380

## (2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 1462 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: double  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Glycine max

(vii) IMMEDIATE SOURCE:

(B) CLONE: pSF2-165K

(ix) FEATURE:

(A) NAME/KEY: CDS

(B) LOCATION: 108..1247

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

CCATATACTA ATATTTGCTT GTATTGATAG CCCCTCCGTT CCCAAGAGTA TAAAACTGCA  
 60

TCGAATAATA CAAGCCACTA GGCATGGGTC TAGCAAAGGA AACAACA ATG GGA GGT  
 116

Met Gly Gly  
 1

AGA GGT CGT GTG GCC AAA GTG GAA GTT CAA GGG AAG AAG CCT CTC TCA  
 164

Arg Gly Arg Val Ala Lys Val Glu Val Gln Gly Lys Lys Pro Leu Ser  
 5 10 15

AGG GTT CCA AAC ACA AAG CCA CCA TTC ACT GTT GGC CAA CTC AAG AAA  
 212

Arg Val Pro Asn Thr Lys Pro Pro Phe Thr Val Gly Gln Leu Lys Lys  
 20 25 30 35

GCA ATT CCA CCA CAC TGC TTT CAG CGC TCC CTC CTC ACT TCA TTC TCC  
 260

Ala Ile Pro Pro His Cys Phe Gln Arg Ser Leu Leu Thr Ser Phe Ser  
 40 45 50



TAT GTT GTT TAT GAC CTT TCA TTT GCC TTC ATT TTC TAC ATT GCC ACC  
 308  
 Tyr Val Val Tyr Asp Leu Ser Phe Ala Phe Ile Phe Tyr Ile Ala Thr  
 55 60 65

ACC TAC TTC CAC CTC CTT CCT CAA CCC TTT TCC CTC ATT GCA TGG CCA  
 356  
 Thr Tyr Phe His Leu Leu Pro Gln Pro Phe Ser Leu Ile Ala Trp Pro  
 70 75 80

ATC TAT TGG GTT CTC CAA GGT TGC CTT CTC ACT GGT GTG TGG GTG ATT  
 404  
 Ile Tyr Trp Val Leu Gln Gly Cys Leu Leu Thr Gly Val Trp Val Ile  
 85 90 95

GCT CAC GAG TGT GGT CAC CAT GCC TTC AGC AAG TAC CAA TGG GTT GAT  
 452  
 Ala His Glu Cys Gly His His Ala Phe Ser Lys Tyr Gln Trp Val Asp  
 100 105 110 115

GAT GTT GTG GGT TTG ACC CTT CAC TCA ACA CTT TTA GTC CCT TAT TTC  
 500  
 Asp Val Val Gly Leu Thr Leu His Ser Thr Leu Leu Val Pro Tyr Phe  
 120 125 130

TCA TGG AAA ATA AGC CAT CGC CGC CAT CAC TCC AAC ACA GGT TCC CTT  
 548  
 Ser Trp Lys Ile Ser His Arg Arg His His Ser Asn Thr Gly Ser Leu  
 135 140 145

GAC CGT GAT GAA GTG TTT GTC CCA AAA CCA AAA TCC AAA GTT GCA TGG  
 596  
 Asp Arg Asp Glu Val Phe Val Pro Lys Pro Lys Ser Lys Val Ala Trp  
 150 155 160

TTT TCC AAG TAC TTA AAC AAC CCT CTA GGA AGG GCT GTT TCT CTT CTC  
 644  
 Phe Ser Lys Tyr Leu Asn Asn Pro Leu Gly Arg Ala Val Ser Leu Leu  
 165 170 175

GTC ACA CTC ACA ATA GGG TGG CCT ATG TAT TTA GCC TTC AAT GTC TCT  
 692  
 Val Thr Leu Thr Ile Gly Trp Pro Met Tyr Leu Ala Phe Asn Val Ser  
 180 185 190 195

GGT AGA CCC TAT GAT AGT TTT GCA AGC CAC TAC CAC CCT TAT GCT CCC  
 740  
 Gly Arg Pro Tyr Asp Ser Phe Ala Ser His Tyr His Pro Tyr Ala Pro  
 200 205 210

ATA TAT TCT AAC CGT GAG AGG CTT CTG ATC TAT GTC TCT GAT GTT GCT  
 788  
 Ile Tyr Ser Asn Arg Glu Arg Leu Leu Ile Tyr Val Ser Asp Val Ala  
 215 220 225

TTG TTT TCT GTG ACT TAC TCT CTC TAC CGT GTT GCA ACC CTG AAA GGG

836  
 Leu Phe Ser Val Thr Tyr Ser Leu Tyr Arg Val Ala Thr Leu Lys Gly  
 230 235 240  
  
 TTG GTT TGG CTG CTA TGT GTT TAT GGG GTG CCT TTG CTC ATT GTG AAC  
 884  
 Leu Val Trp Leu Leu Cys Val Tyr Gly Val Pro Leu Leu Ile Val Asn  
 245 250 255  
  
 GGT TTT CTT GTG ACT ATC ACA TAT TTG CAG CAC ACA CAC TTT GCC TTG  
 932  
 Gly Phe Leu Val Thr Ile Thr Tyr Leu Gln His Thr His Phe Ala Leu  
 260 265 270 275  
  
 CCT CAT TAC GAT TCA TCA GAA TGG GAC TGG CTG AAG GGA GCT TTG GCA  
 980  
 Pro His Tyr Asp Ser Ser Glu Trp Asp Trp Leu Lys Gly Ala Leu Ala  
 280 285 290  
  
 ACT ATG GAC AGA GAT TAT GGG ATT CTG AAC AAG GTG TTT CAT CAC ATA  
 1028  
 Thr Met Asp Arg Asp Tyr Gly Ile Leu Asn Lys Val Phe His His Ile  
 295 300 305  
  
 ACT GAT ACT CAT GTG GCT CAC CAT CTC TTC TCT ACA ATG CCA CAT TAC  
 1076  
 Thr Asp Thr His Val Ala His His Leu Phe Ser Thr Met Pro His Tyr  
 310 315 320  
  
 CAT GCA ATG GAG GCA ACC AAT GCA ATC AAG CCA ATA TTG GGT GAG TAC  
 1124  
 His Ala Met Glu Ala Thr Asn Ala Ile Lys Pro Ile Leu Gly Glu Tyr  
 325 330 335  
  
 TAC CAA TTT GAT GAC ACA CCA TTT TAC AAG GCA CTG TGG AGA GAA GCG  
 1172  
 Tyr Gln Phe Asp Asp Thr Pro Phe Tyr Lys Ala Leu Trp Arg Glu Ala  
 340 345 350 355  
  
 AGA GAG TGC CTC TAT GTG GAG CCA GAT GAA GGA ACA TCC GAG AAG GGC  
 1220  
 Arg Glu Cys Leu Tyr Val Glu Pro Asp Glu Gly Thr Ser Glu Lys Gly  
 360 365 370  
  
 GTG TAT TGG TAC AGG AAC AAG TAT TGATGGAGCA ACCAATGGGC CATAGTGGGA  
 1274  
 Val Tyr Trp Tyr Arg Asn Lys Tyr 380  
  
 GTTATGGAAG TTTTGTCATG TATTAGTACA TAATTAGTAG AATGTTATAA ATAAGTGGAT  
 1334  
  
 TTGCCGCGTA ATGACTTTGT GTGTATTGTG AAACAGCTTG TTGCGATCAT GGTTATAATG  
 1394  
  
 TAAAAATAAT TCTGGTATTA ATTACATGTG GAAAGTGTTT TGCTTATAGC TTTCTGCCTA

1454

AAAAAAA

1462

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 379 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

```

Met Gly Gly Arg Gly Arg Val Ala Lys Val Glu Val Gln Gly Lys Lys
 1           5           10           15

Pro Leu Ser Arg Val Pro Asn Thr Lys Pro Pro Phe Thr Val Gly Gln
          20           25           30

Leu Lys Lys Ala Ile Pro Pro His Cys Phe Gln Arg Ser Leu Leu Thr
      35           40           45

Ser Phe Ser Tyr Val Val Tyr Asp Leu Ser Phe Ala Phe Ile Phe Tyr
      50           55           60

Ile Ala Thr Thr Tyr Phe His Leu Leu Pro Gln Pro Phe Ser Leu Ile
      65           70           75           80

Ala Trp Pro Ile Tyr Trp Val Leu Gln Gly Cys Leu Leu Thr Gly Val
          85           90           95

Trp Val Ile Ala His Glu Cys Gly His His Ala Phe Ser Lys Tyr Gln
          100          105          110

Trp Val Asp Asp Val Val Gly Leu Thr Leu His Ser Thr Leu Leu Val
          115          120          125

Pro Tyr Phe Ser Trp Lys Ile Ser His Arg Arg His His Ser Asn Thr
          130          135          140

Gly Ser Leu Asp Arg Asp Glu Val Phe Val Pro Lys Pro Lys Ser Lys
145           150           155           160

Val Ala Trp Phe Ser Lys Tyr Leu Asn Asn Pro Leu Gly Arg Ala Val
          165          170          175

Ser Leu Leu Val Thr Leu Thr Ile Gly Trp Pro Met Tyr Leu Ala Phe
          180          185          190

Asn Val Ser Gly Arg Pro Tyr Asp Ser Phe Ala Ser His Tyr His Pro
          195          200          205

Tyr Ala Pro Ile Tyr Ser Asn Arg Glu Arg Leu Leu Ile Tyr Val Ser

```

(2) INFORMATION FOR SEQ ID NO:7:

(A) LENGTH: 1790 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: double  
(D) TOPOLOGY: linear

(iii) HYPOTHETICAL: NO

(vi) ORIGINAL SOURCE:

(vii) IMMEDIATE SOURCE:

(ix) FEATURE:

(B) LOCATION: 165..1328

CGGCTCTCC CCTCCCTCCT CCCTGCAAAT CCTGCAGACA CCACCGCTCG TTTTCTCTC  
60

CGGGACAGGA GAAAAGGGGA GAGAGAGGTG AGGCGCGGTG TCCGCCCCGAT CTGCTCTGCC  
120

CCGACGCAGC TGTTACGACC TCCTCAGTCT CAGTCAGGAG CAAG ATG GGT GCC GGC  
176

Met Gly Ala Gly  
1

GGC AGG ATG ACC GAG AAG GAG CGG GAG AAG CAG GAG CAG CTC GCC CGA  
224

224  
Gly Arg Met Thr Glu Lys Glu Arg Glu Lys Gln Glu Gln Leu Ala Arg  
5 10 15 20

GCT ACC GGT GGC GCC GCG ATG CAG CGG TCG CCG GTG GAG AAG CCT CCG  
272

Ala Thr Gly Gly Ala Ala Met Gln Arg Ser Pro Val Glu Lys Pro Pro  
25 30 35

TTC ACT CTG GGT CAG ATC AAG AAG GCC ATC CCG CCA CAC TGC TTC GAG  
320

Phe Thr Leu Gly Gln Ile Lys Lys Ala Ile Pro Pro His Cys Phe Glu  
40 45 50

CGC TCG GTG CTC AAG TCC TTC TCG TAC GTG GTC CAC GAC CTG GTG ATC  
368

Arg Ser Val Leu Lys Ser Phe Ser Tyr Val Val His Asp Leu Val Ile  
55 60 65

GCC GCG GCG CTC CTC TAC TTC GCG CTG GCC ATC ATA CCG GCG CTC CCA  
416

Ala Ala Ala Leu Leu Tyr Phe Ala Leu Ala Ile Ile Pro Ala Leu Pro  
70 75 80

AGC CCG CTC CGC TAC GCC GCC TGG CCG CTG TAC TGG ATC GCG CAG GGG  
464

Ser Pro Leu Arg Tyr Ala Ala Trp Pro Leu Tyr Trp Ile Ala Gln Gly  
85 90 95 100

TGC GTG TGC ACC GGC GTG TGG GTC ATC GCG CAC GAG TGC GGC CAC CAC  
512

Cys Val Cys Thr Gly Val Trp Val Ile Ala His Glu Cys Gly His His  
105 110 115

GCC TTC TCG GAC TAC TCG CTC CTG GAC GAC GTG GTC GGC CTG GTG CTG  
560

Ala Phe Ser Asp Tyr Ser Leu Leu Asp Asp Val Val Gly Leu Val Leu  
120 125 130

CAC TCG TCG CTC ATG GTG CCC TAC TTC TCG TGG AAG TAC AGC CAC CGG  
608

His Ser Ser Leu Met Val Pro Tyr Phe Ser Trp Lys Tyr Ser His Arg

135	140	145
CGC CAC CAC TCC AAC ACG GGG TCC CTG GAG CGC GAC GAG GTG TTC GTG		
656		
Arg His His Ser Asn Thr Gly Ser Leu Glu Arg Asp Glu Val Phe Val		
150	155	160
CCC AAG AAG AAG GAG GCG CTG CCG TGG TAC ACC CCG TAC GTG TAC AAC		
704		
Pro Lys Lys Lys Glu Ala Leu Pro Trp Tyr Thr Pro Tyr Val Tyr Asn		
165	170	175
AAC CCG GTC GGC CGG GTG GTG CAC ATC GTG GTG CAG CTC ACC CTC GGG		
752		
Asn Pro Val Gly Arg Val Val His Ile Val Val Gln Leu Thr Leu Gly		
	185	190
TGG CCG CTG TAC CTG GCG ACC AAC GCG TCG GGG CGG CCG TAC CCG CGC		
800		
Trp Pro Leu Tyr Leu Ala Thr Asn Ala Ser Gly Arg Pro Tyr Pro Arg		
	200	205
TTC GCC TGC CAC TTC GAC CCC TAC GGC CCC ATC TAC AAC GAC CGG GAG		
848		
Phe Ala Cys His Phe Asp Pro Tyr Gly Pro Ile Tyr Asn Asp Arg Glu		
	215	220
CGC GCC CAG ATC TTC GTC TCG GAC GCC GGC GTC GTG GCC GTG GCG TTC		
896		
Arg Ala Gln Ile Phe Val Ser Asp Ala Gly Val Val Ala Val Ala Phe		
	230	235
GGG CTG TAC AAG CTG GCG GCG GCG TTC GGG GTC TGG TGG GTG GTG CGC		
944		
Gly Leu Tyr Lys Leu Ala Ala Ala Phe Gly Val Trp Trp Val Val Arg		
245	250	255
GTG TAC GCC GTG CCG CTG CTG ATC GTG AAC GCG TGG CTG GTG CTC ATC		
992		
Val Tyr Ala Val Pro Leu Leu Ile Val Asn Ala Trp Leu Val Leu Ile		
	265	270
ACC TAC CTG CAG CAC ACC CAC CCG TCG CTC CCC CAC TAC GAC TCG AGC		
1040		
Thr Tyr Leu Gln His Thr His Pro Ser Leu Pro His Tyr Asp Ser Ser		
	280	285
GAG TGG GAC TGG CTG CGC GGC GCG CTG GCC ACC ATG GAC CGC GAC TAC		
1088		
Glu Trp Asp Trp Leu Arg Gly Ala Leu Ala Thr Met Asp Arg Asp Tyr		
	295	300
GGC ATC CTC AAC CGC GTG TTC CAC AAC ATC ACG GAC ACG CAC GTC GCG		
1136		
Gly Ile Leu Asn Arg Val Phe His Asn Ile Thr Asp Thr His Val Ala		
	310	315
		320

CAC CAC CTC TTC TCC ACC ATG CCG CAC TAC CAC GCC ATG GAG GCC ACC  
1184  
His His Leu Phe Ser Thr Met Pro His Tyr His Ala Met Glu Ala Thr  
325 330 335 340  
AAG GCG ATC AGG CCC ATC CTC GGC GAC TAC TAC CAC TTC GAC CCG ACC  
1232  
Lys Ala Ile Arg Pro Ile Leu Gly Asp Tyr Tyr His Phe Asp Pro Thr  
345 350 355  
CCT GTC GCC AAG GCG ACC TGG CGC GAG GCC GGG GAA TGC ATC TAC GTC  
1280  
Pro Val Ala Lys Ala Thr Trp Arg Glu Ala Gly Glu Cys Ile Tyr Val  
360 365 370  
GAG CCC GAG GAC CGC AAG GGC GTC TTC TGG TAC AAC AAG AAG TTC TAGCCGCCG  
C 1335  
Glu Pro Glu Asp Arg Lys Gly Val Phe Trp Tyr Asn Lys Lys Phe  
375 380 385  
CGCTCGCAGA GCTGAGGACG CTACCGTAGG AATGGGAGCA GAAACCAGGA GGAGGAGACG  
1395  
GTACTCGCCC CAAAGTCTCC GTCAACCTAT CTAATCGTTA GTCGTCAGTC TTTTAGACGG  
1455  
GAAGAGAGAT CATTTGGGCA CAGAGACGAA GGCTTACTGC AGTGCCATCG CTAGAGCTGC  
1515  
CATCAAGTAC AAGTAGGCAA ATTCGTCAAC TTAGTGTGTC CCATGTTGTT TTTCTTAGTC  
1575  
GTCCGCTGCT GTAGGCTTTC CGGCGGCGGT CGTTTGTGTG GTTGGCATCC GTGGCCATGC  
1635  
CTGTGCGTGC GTGGCCGCGC TTGTCGTGTG CGTCTGTCGT CGCGTTGGCG TCGTCTCTTC  
1695  
GTGCTCCCCG TGTGTTGTTG TAAACAAGA AGATGTTTTT TGGTGTCTTT GGCGGAATAA  
1755  
CAGATCGTCC GAACGAAAAA AAAAAAAAAA AAAAA  
1790

## (2) INFORMATION FOR SEQ ID NO:8:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 387 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

```

Met Gly Ala Gly Gly Arg Met Thr Glu Lys Glu Arg Glu Lys Gln Glu
 1          5          10          15

Gln Leu Ala Arg Ala Thr Gly Gly Ala Ala Met Gln Arg Ser Pro Val
          20          25          30

Glu Lys Pro Pro Phe Thr Leu Gly Gln Ile Lys Lys Ala Ile Pro Pro
          35          40          45

His Cys Phe Glu Arg Ser Val Leu Lys Ser Phe Ser Tyr Val Val His
 50          55          60

Asp Leu Val Ile Ala Ala Ala Leu Leu Tyr Phe Ala Leu Ala Ile Ile
 65          70          75          80

Pro Ala Leu Pro Ser Pro Leu Arg Tyr Ala Ala Trp Pro Leu Tyr Trp
          85          90          95

Ile Ala Gln Gly Cys Val Cys Thr Gly Val Trp Val Ile Ala His Glu
          100          105          110

Cys Gly His His Ala Phe Ser Asp Tyr Ser Leu Leu Asp Asp Val Val
          115          120          125

Gly Leu Val Leu His Ser Ser Leu Met Val Pro Tyr Phe Ser Trp Lys
          130          135          140

Tyr Ser His Arg Arg His His Ser Asn Thr Gly Ser Leu Glu Arg Asp
145          150          155          160

Glu Val Phe Val Pro Lys Lys Lys Glu Ala Leu Pro Trp Tyr Thr Pro
          165          170          175

Tyr Val Tyr Asn Asn Pro Val Gly Arg Val Val His Ile Val Val Gln
          180          185          190

Leu Thr Leu Gly Trp Pro Leu Tyr Leu Ala Thr Asn Ala Ser Gly Arg
          195          200          205

Pro Tyr Pro Arg Phe Ala Cys His Phe Asp Pro Tyr Gly Pro Ile Tyr
          210          215          220

Asn Asp Arg Glu Arg Ala Gln Ile Phe Val Ser Asp Ala Gly Val Val
225          230          235          240

Ala Val Ala Phe Gly Leu Tyr Lys Leu Ala Ala Ala Phe Gly Val Trp
          245          250          255

Trp Val Val Arg Val Tyr Ala Val Pro Leu Leu Ile Val Asn Ala Trp
          260          265          270

Leu Val Leu Ile Thr Tyr Leu Gln His Thr His Pro Ser Leu Pro His
          275          280          285

Tyr Asp Ser Ser Glu Trp Asp Trp Leu Arg Gly Ala Leu Ala Thr Met
          290          295          300

```



Asp Arg Asp Tyr Gly Ile Leu Asn Arg Val Phe His Asn Ile Thr Asp  
 305 310 315 320  
 Thr His Val Ala His His Leu Phe Ser Thr Met Pro His Tyr His Ala  
 325 330 335  
 Met Glu Ala Thr Lys Ala Ile Arg Pro Ile Leu Gly Asp Tyr Tyr His  
 340 345 350  
 Phe Asp Pro Thr Pro Val Ala Lys Ala Thr Trp Arg Glu Ala Gly Glu  
 355 360 365  
 Cys Ile Tyr Val Glu Pro Glu Asp Arg Lys Gly Val Phe Trp Tyr Asn  
 370 375 380  
 Lys Lys Phe  
 385

## (2) INFORMATION FOR SEQ ID NO:9:

(i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 673 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: double  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:  
 (A) ORGANISM: Ricinus communis

(vii) IMMEDIATE SOURCE:  
 (B) CLONE: pRF2-1C

(ix) FEATURE:  
 (A) NAME/KEY: CDS  
 (B) LOCATION: 1..673

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

TGG GTG ATG GCG CAT GAT TGT GGG CAC CAT GCC TTC AGT GAC TAT CAA  
 48  
 Trp Val Met Ala His Asp Cys Gly His His Ala Phe Ser Asp Tyr Gln  
 1 5 10 15  
 TTG CTT GAT GAT GTA GTT GGT CTT ATC CTA CAC TCC TGT CTC CTT GTC  
 96  
 Leu Leu Asp Asp Val Val Gly Leu Ile Leu His Ser Cys Leu Leu Val  
 20 25 30  
 CCT TAT TTT TCA TGG AAA CAC AGC CAT CGC CGA CAT CAT TCC AAC ACA

144  
 Pro Tyr Phe Ser Trp Lys His Ser His Arg Arg His His Ser Asn Thr  
           35                                  40                                  45

GGG TCC CTG GAA CGG GAT GAA GTG TTT GTT CCC AAG AAG AAA TCT AGT  
 192  
 Gly Ser Leu Glu Arg Asp Glu Val Phe Val Pro Lys Lys Lys Ser Ser  
           50                                  55                                  60

ATC CGT TGG TAT TCC AAA TAC CTC AAC AAC CCT CCA GGT CGT ATC ATG  
 240  
 Ile Arg Trp Tyr Ser Lys Tyr Leu Asn Asn Pro Pro Gly Arg Ile Met  
           65                                  70                                  75                                  80

ACA ATT GCC GTC ACA CTT TCA CTT GGC TGG CCT CTG TAC CTA GCA TTC  
 288  
 Thr Ile Ala Val Thr Leu Ser Leu Gly Trp Pro Leu Tyr Leu Ala Phe  
                                   85                                  90                                  95

AAT GTT TCA GGC AGG CCA TAT GAT CGG TTC GCC TGC CAC TAT GAC CCA  
 336  
 Asn Val Ser Gly Arg Pro Tyr Asp Arg Phe Ala Cys His Tyr Asp Pro  
                                   100                                  105                                  110

TAT GGC CCG ATC TAC AAT GAT CGC GAG CGA ATC GAG ATA TTC ATA TCA  
 384  
 Tyr Gly Pro Ile Tyr Asn Asp Arg Glu Arg Ile Glu Ile Phe Ile Ser  
           115                                  120                                  125

GAT GCT GGT GTT CTT GCT GTC ACT TTT GGT CTC TAC CAA CTT GCT ATA  
 432  
 Asp Ala Gly Val Leu Ala Val Thr Phe Gly Leu Tyr Gln Leu Ala Ile  
           130                                  135                                  140

GCG AAG GGG CTT GCT TGG GTT GTC TGT GTA TAT GGA GTG CCA TTG TTG  
 480  
 Ala Lys Gly Leu Ala Trp Val Val Cys Val Tyr Gly Val Pro Leu Leu  
           145                                  150                                  155                                  160

GTG GTG AAT TCA TTC CTT GTT CTG ATC ACA TTT CTG CAG CAT ACT CAC  
 528  
 Val Val Asn Ser Phe Leu Val Leu Ile Thr Phe Leu Gln His Thr His  
                                   165                                  170                                  175

CCT GCA TTG CCA CAT TAT GAT TCG TCG GAG TGG GAC TGG CTA AGA GGA  
 576  
 Pro Ala Leu Pro His Tyr Asp Ser Ser Glu Trp Asp Trp Leu Arg Gly  
                                   180                                  185                                  190

GCT CTA GCA ACT GTT GAC AGA GAT TAC GGG ATC TTG AAC AAG GTG TTC  
 624  
 Ala Leu Ala Thr Val Asp Arg Asp Tyr Gly Ile Leu Asn Lys Val Phe  
           195                                  200                                  205

CAT AAC ATA ACG GAC ACT CAA GTA GCT CAC CAC CTT TTC ACC ATG CCC C  
 673

His Asn Ile Thr Asp Thr Gln Val Ala His His Leu Phe Thr Met Pro  
 210 215 220

(2) INFORMATION FOR SEQ ID NO:10:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 224 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

Trp Val Met Ala His Asp Cys Gly His His Ala Phe Ser Asp Tyr Gln  
 1 5 10 15  
 Leu Leu Asp Asp Val Val Gly Leu Ile Leu His Ser Cys Leu Leu Val  
 20 25 30  
 Pro Tyr Phe Ser Trp Lys His Ser His Arg Arg His His Ser Asn Thr  
 35 40 45  
 Gly Ser Leu Glu Arg Asp Glu Val Phe Val Pro Lys Lys Lys Ser Ser  
 50 55 60  
 Ile Arg Trp Tyr Ser Lys Tyr Leu Asn Asn Pro Pro Gly Arg Ile Met  
 65 70 75 80  
 Thr Ile Ala Val Thr Leu Ser Leu Gly Trp Pro Leu Tyr Leu Ala Phe  
 85 90 95  
 Asn Val Ser Gly Arg Pro Tyr Asp Arg Phe Ala Cys His Tyr Asp Pro  
 100 105 110  
 Tyr Gly Pro Ile Tyr Asn Asp Arg Glu Arg Ile Glu Ile Phe Ile Ser  
 115 120 125  
 Asp Ala Gly Val Leu Ala Val Thr Phe Gly Leu Tyr Gln Leu Ala Ile  
 130 135 140  
 Ala Lys Gly Leu Ala Trp Val Val Cys Val Tyr Gly Val Pro Leu Leu  
 145 150 155 160  
 Val Val Asn Ser Phe Leu Val Leu Ile Thr Phe Leu Gln His Thr His  
 165 170 175  
 Pro Ala Leu Pro His Tyr Asp Ser Ser Glu Trp Asp Trp Leu Arg Gly  
 180 185 190  
 Ala Leu Ala Thr Val Asp Arg Asp Tyr Gly Ile Leu Asn Lys Val Phe  
 195 200 205  
 His Asn Ile Thr Asp Thr Gln Val Ala His His Leu Phe Thr Met Pro  
 210 215 220

## (2) INFORMATION FOR SEQ ID NO:11:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1369 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: double  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Ricinus communis

(vii) IMMEDIATE SOURCE:

(B) CLONE: pRF197c-42

(ix) FEATURE:

- (A) NAME/KEY: CDS  
 (B) LOCATION: 184..1347

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

CGGCCGGGAT TCCGGTTTTTC AACTAATTT GCAAAAAATG CATGATTTC CCTCAAATCA  
 60

AACACCACAC CTTATAACTT AGTCTTAAGA GAGAGAGAGA GAGGAGACAT TTCTCTTCTC  
 120

TGAGATGAGC ACTTCTCTTC CAGACATCGA AGCCTCAGGA AAGTGCTTGA GAAGAGCTTG  
 180

AGA ATG GGA GGT GGT GGT CGC ATG TCT ACT GTC ATA ATC AGC AAC AAC  
 228

Met Gly Gly Gly Gly Arg Met Ser Thr Val Ile Ile Ser Asn Asn  
 1 5 10 15

AGT GAG AAG AAA GGA GGA AGC AGC CAC CTG GAG CGA GCG CCG CAC ACG  
 276

Ser Glu Lys Lys Gly Gly Ser Ser His Leu Glu Arg Ala Pro His Thr  
 20 25 30

AAG CCT CCT TAC ACA CTT GGT AAC CTC AAG AGA GCC ATC CCA CCC CAT  
 324

Lys Pro Pro Tyr Thr Leu Gly Asn Leu Lys Arg Ala Ile Pro Pro His  
 35 40 45

TGC TTT GAA CGC TCT TTT GTG CGC TCA TTC TCC AAT TTT GCC TAT AAT  
 372

Cys Phe Glu Arg Ser Phe Val Arg Ser Phe Ser Asn Phe Ala Tyr Asn  
 50 55 60

TTC TGC TTA AGT TTT CTT TCC TAC TCG ATC GCC ACC AAC TTC TTC CCT

420  
 Phe Cys Leu Ser Phe Leu Ser Tyr Ser Ile Ala Thr Asn Phe Phe Pro  
     65                    70                    75  
  
 TAC ATC TCT TCT CCG CTC TCG TAT GTC GCT TGG CTG GTT TAC TGG CTC  
 468  
 Tyr Ile Ser Ser Pro Leu Ser Tyr Val Ala Trp Leu Val Tyr Trp Leu  
     80                    85                    90                    95  
  
 TTC CAA GGC TGC ATT CTC ACT GGT CTT TGG GTC ATC GGC CAT GAA TGT  
 516  
 Phe Gln Gly Cys Ile Leu Thr Gly Leu Trp Val Ile Gly His Glu Cys  
                     100                    105                    110  
  
 GGC CAT CAT GCT TTT AGT GAG TAT CAG CTG GCT GAT GAC ATT GTT GGC  
 564  
 Gly His His Ala Phe Ser Glu Tyr Gln Leu Ala Asp Asp Ile Val Gly  
                     115                    120                    125  
  
 CTA ATT GTC CAT TCT GCA CTT CTG GTT CCA TAT TTT TCA TGG AAA TAT  
 612  
 Leu Ile Val His Ser Ala Leu Leu Val Pro Tyr Phe Ser Trp Lys Tyr  
                     130                    135                    140  
  
 AGC CAT CGC CGC CAC CAT TCT AAC ATA GGA TCT CTC GAG CGA GAC GAA  
 660  
 Ser His Arg Arg His His Ser Asn Ile Gly Ser Leu Glu Arg Asp Glu  
                     145                    150                    155  
  
 GTG TTC GTC CCG AAA TCA AAG TCG AAA ATT TCA TGG TAT TCT AAG TAC  
 708  
 Val Phe Val Pro Lys Ser Lys Ser Lys Ile Ser Trp Tyr Ser Lys Tyr  
     160                    165                    170                    175  
  
 TTA AAC AAC CCG CCA GGT CGA GTT TTG ACA CTT GCT GCC ACG CTC CTC  
 756  
 Leu Asn Asn Pro Pro Gly Arg Val Leu Thr Leu Ala Ala Thr Leu Leu  
                     180                    185                    190  
  
 CTT GGC TGG CCT TTA TAT TTA GCT TTC AAT GTC TCT GGT AGA CCT TAC  
 804  
 Leu Gly Trp Pro Leu Tyr Leu Ala Phe Asn Val Ser Gly Arg Pro Tyr  
                     195                    200                    205  
  
 GAT CGC TTT GCT TGC CAT TAT GAT CCC TAT GGC CCA ATA TTT TCC GAA  
 852  
 Asp Arg Phe Ala Cys His Tyr Asp Pro Tyr Gly Pro Ile Phe Ser Glu  
                     210                    215                    220  
  
 AGA GAA AGG CTT CAG ATT TAC ATT GCT GAC CTC GGA ATC TTT GCC ACA  
 900  
 Arg Glu Arg Leu Gln Ile Tyr Ile Ala Asp Leu Gly Ile Phe Ala Thr  
     225                    230                    235  
  
 ACG TTT GTG CTT TAT CAG GCT ACA ATG GCA AAA GGG TTG GCT TGG GTA  
 948

Thr Phe Val Leu Tyr Gln Ala Thr Met Ala Lys Gly Leu Ala Trp Val  
 240 245 250 255  
 ATG CGT ATC TAT GGG GTG CCA TTG CTT ATT GTT AAC TGT TTC CTT GTT  
 996  
 Met Arg Ile Tyr Gly Val Pro Leu Leu Ile Val Asn Cys Phe Leu Val  
 260 265 270  
 ATG ATC ACA TAC TTG CAG CAC ACT CAC CCA GCT ATT CCA CGC TAT GGC  
 1044  
 Met Ile Thr Tyr Leu Gln His Thr His Pro Ala Ile Pro Arg Tyr Gly  
 275 280 285  
 TCA TCG GAA TGG GAT TGG CTC CGG GGA GCA ATG GTG ACT GTC GAT AGA  
 1092  
 Ser Ser Glu Trp Asp Trp Leu Arg Gly Ala Met Val Thr Val Asp Arg  
 290 295 300  
 GAT TAT GGG GTG TTG AAT AAA GTA TTC CAT AAC ATT GCA GAC ACT CAT  
 1140  
 Asp Tyr Gly Val Leu Asn Lys Val Phe His Asn Ile Ala Asp Thr His  
 305 310 315  
 GTA GCT CAT CAT CTC TTT GCT ACA GTG CCA CAT TAC CAT GCA ATG GAG  
 1188  
 Val Ala His His Leu Phe Ala Thr Val Pro His Tyr His Ala Met Glu  
 320 325 330 335  
 GCC ACT AAA GCA ATC AAG CCT ATA ATG GGT GAG TAT TAC CGG TAT GAT  
 1236  
 Ala Thr Lys Ala Ile Lys Pro Ile Met Gly Glu Tyr Tyr Arg Tyr Asp  
 340 345 350  
 GGT ACC CCA TTT TAC AAG GCA TTG TGG AGG GAG GCA AAG GAG TGC TTG  
 1284  
 Gly Thr Pro Phe Tyr Lys Ala Leu Trp Arg Glu Ala Lys Glu Cys Leu  
 355 360 365  
 TTC GTC GAG CCA GAT GAA GGA GCT CCT ACA CAA GGC GTT TTC TGG TAC  
 1332  
 Phe Val Glu Pro Asp Glu Gly Ala Pro Thr Gln Gly Val Phe Trp Tyr  
 370 375 380  
 CGG AAC AAG TAT TAAAAAAGTG TCATGTAGCC TGCCG  
 1369  
 Arg Asn Lys Tyr  
 385

## (2) INFORMATION FOR SEQ ID NO:12:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 387 amino acids
  - (B) TYPE: amino acid
  - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

```

Met Gly Gly Gly Gly Arg Met Ser Thr Val Ile Ile Ser Asn Asn Ser
 1           5           10           15

Glu Lys Lys Gly Gly Ser Ser His Leu Glu Arg Ala Pro His Thr Lys
           20           25           30

Pro Pro Tyr Thr Leu Gly Asn Leu Lys Arg Ala Ile Pro Pro His Cys
           35           40           45

Phe Glu Arg Ser Phe Val Arg Ser Phe Ser Asn Phe Ala Tyr Asn Phe
 50           55           60

Cys Leu Ser Phe Leu Ser Tyr Ser Ile Ala Thr Asn Phe Phe Pro Tyr
 65           70           75           80

Ile Ser Ser Pro Leu Ser Tyr Val Ala Trp Leu Val Tyr Trp Leu Phe
           85           90           95

Gln Gly Cys Ile Leu Thr Gly Leu Trp Val Ile Gly His Glu Cys Gly
           100          105          110

His His Ala Phe Ser Glu Tyr Gln Leu Ala Asp Asp Ile Val Gly Leu
          115          120          125

Ile Val His Ser Ala Leu Leu Val Pro Tyr Phe Ser Trp Lys Tyr Ser
          130          135          140

His Arg Arg His His Ser Asn Ile Gly Ser Leu Glu Arg Asp Glu Val
          145          150          155          160

Phe Val Pro Lys Ser Lys Ser Lys Ile Ser Trp Tyr Ser Lys Tyr Leu
           165          170          175

Asn Asn Pro Pro Gly Arg Val Leu Thr Leu Ala Ala Thr Leu Leu Leu
          180          185          190

Gly Trp Pro Leu Tyr Leu Ala Phe Asn Val Ser Gly Arg Pro Tyr Asp
          195          200          205

Arg Phe Ala Cys His Tyr Asp Pro Tyr Gly Pro Ile Phe Ser Glu Arg
          210          215          220

Glu Arg Leu Gln Ile Tyr Ile Ala Asp Leu Gly Ile Phe Ala Thr Thr
          225          230          235          240

Phe Val Leu Tyr Gln Ala Thr Met Ala Lys Gly Leu Ala Trp Val Met
          245          250          255

Arg Ile Tyr Gly Val Pro Leu Leu Ile Val Asn Cys Phe Leu Val Met
          260          265          270

Ile Thr Tyr Leu Gln His Thr His Pro Ala Ile Pro Arg Tyr Gly Ser
          275          280          285

```

Ser Glu Trp Asp Trp Leu Arg Gly Ala Met Val Thr Val Asp Arg Asp  
 290 295 300

Tyr Gly Val Leu Asn Lys Val Phe His Asn Ile Ala Asp Thr His Val  
 305 310 315 320

Ala His His Leu Phe Ala Thr Val Pro His Tyr His Ala Met Glu Ala  
 325 330 335

Thr Lys Ala Ile Lys Pro Ile Met Gly Glu Tyr Tyr Arg Tyr Asp Gly  
 340 345 350

Thr Pro Phe Tyr Lys Ala Leu Trp Arg Glu Ala Lys Glu Cys Leu Phe  
 355 360 365

Val Glu Pro Asp Glu Gly Ala Pro Thr Gln Gly Val Phe Trp Tyr Arg  
 370 375 380

Asn Lys Tyr  
 385

## (2) INFORMATION FOR SEQ ID NO:13:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 23 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

## (ix) FEATURE:

- (A) NAME/KEY: misc\_feature
- (B) LOCATION: 1..23
- (D) OTHER INFORMATION: /product="synthetic oligonucleotide"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

TGGGTATGCC AYGANTGYGG NCA

23

## (2) INFORMATION FOR SEQ ID NO:14:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 22 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA



(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(ix) FEATURE:

(A) NAME/KEY: misc\_feature

(B) LOCATION: 1..22

(D) OTHER INFORMATION: /product=  
"synthetic  
oligonucleotide"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

AAARTGRTGG CACRTGNGTR TC

22

(2) INFORMATION FOR SEQ ID NO:15:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 2973 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: double

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Arabidopsis thaliana

(vii) IMMEDIATE SOURCE:

(B) CLONE: pAGF2-6

(ix) FEATURE:

(A) NAME/KEY: exon

(B) LOCATION: 433..520

(ix) FEATURE:

(A) NAME/KEY: intron

(B) LOCATION: 521..1654

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

ATTCGGTAAT TCCTACATAT TTTAGAGATT AGTTTGAGTT TCCATCCATA CTTTACTAGT  
60

GATTATAAAT TTAAATACG TACTTTTCGA CTATAAAGTG AACTAAGTA AATTAGAACG 1  
20

TGATATTAAA AAGTTAATGT TCACTGTTAT ATTTTTTTCA CAAGTAAAAA ATGGGTTATT 1  
80

TGCGGTAAAT AAAAATACCA GATATTTTGA ATTGATTAAA AAGGTTGAAA TAAGAGAGGA 2

40

GGGGAAAGAA AAGAAGGTGG GGGCCCAGTA TGAAAGGGAA AGGTGTCATC AAATCATCTC 3  
00

TCTCTCTCTC TACCTTCGAC CCACGGGCCG TGTCCATTTA AAGCCCTGTC TCTTGCCATT 3  
60

CCCCATCTGA CCACCAGAAG AAGAGCCACA CACTCACAAA TTAAAAAGAG AGAGAGAGAG 4  
20

AGAGAGACAG AGAGAGAGAG AGATTCTGCG GAGGAGCTTC TTCTTCGTAG GGTGTTTCATC 4  
80

GTTATTAACG TTATCGCCCC TACGTCAGCT CCATCTCCAG GTCCGTCGCT TCTCTTCCAT 5  
40

TTCTTCTCAT TTTTCGATTTT GATTCTTATT TCTTTCCAGT AGCTCCTGCT CTGTGAATTT 6  
00

CTCCGCTCAC GATAGATCTG CTTATACTCC TTACATTCAA CCTTAGATCT GGTCTCGATT 6  
60

CTCTGTTTCT CTGTTTTTTT CTTTTGGTCG AGAATCTGAT GTTTGTTTAT GTTCTGTCAC 7  
20

CATTAATAAT GATGAACTCT CTCATTCATA CAATGATTAG TTTCTCTCGT CTACCAAACG 7  
80

ATATGTTGCA TTTTCACTTT TCTTCTTTTT TTCTAAGATG ATTTGCTTTG ACCAATTTGT 8  
40

TTAGATCTTT ATTTTATTTT ATTTTCTGGT GGGTTGGTGG AAATTGAAAA AAAAAAAAAA 9  
00

AAAAGCATAA ATTGTTATTT GTTAATGTAT TCATTTTTTG GCTATTTGTT CTGGGTAAAA 9  
60

ATCTGCTTCT ACTGTTGAAT CTTTCCTGGA TTTTTTACTC CTATTGGGTT TTTATAGTAA 10  
20

AAATACATAA TAAAAGGAAA ACAAAGTTT TATAGATTCT CTAAACCCC TTACGATAAA 10  
80

AGTTGGAATC AAAATAATTC AGGATCAGAT GCTCTTTGAT TGATTCAGAT GCGATTACAG 11  
40

TTGCATGGAA AATTTTCTAG ATCCGTCGTC ACATTTTATT TTCTGTTTAA ATATCTAAAT 12  
00

CTGATATATG ATGTCGACAA ATTCTGGTGG CTTATACATC ACTTCAACTG TTTTCTTTTG 12  
60

GCTTTGTTTG TCAACTTGGT TTTCAATACG ATTTGTGATT TCGATCGCTG AATTTTTAAT 13  
20

ACAAGCAAAC TGATGTTAAC CACAAGCAAG AGATGTGACC TGCCTTATTA ACATCGTATT 13  
 80  
 ACTTACTACT AGTCGTATTC TCAACGCAAT CGTTTTTGTG TTTCTCACAT TATGCCGCTT 14  
 40  
 CTCTACTCTT TATTCCTTTT GGTCCACGCA TTTTCTATTT GTGGCAATCC CTTTCACAAC 15  
 00  
 CTGATTTCCC ACTTTGGATC ATTTGTCTGA AGACTCTCTT GAATCGTTAC CACTTGTTTC 15  
 60  
 TTGTGCATGC TCTGTTTTTT AGAATTAATG ATAAAACTAT TCCATAGTCT TGAGTTTTCA 16  
 20  
 GCTTGTTGAT TCTTTTGCTT TTGGTTTTCT GCAGAAACAT GGGTGCAGGT GGAAGAATGC 16  
 80  
 CGGTTCCCTAC TTCTTCCAAG AAATCGGAAA CCGACACCAC AAAGCGTGTG CCGTGCGAGA 17  
 40  
 AACCGCCTTT CTCGGTGGGA GATCTGAAGA AAGCAATCCC GCCGCATTGT TTCAAACGCT 18  
 00  
 CAATCCCTCG CTCTTTCTCC TACCTTATCA GTGACATCAT TATAGCCTCA TGCTTCTACT 18  
 60  
 ACGTCGCCAC CAATTACTTC TCTCTCCTCC CTCAGCCTCT CTCTTACTTG GCTTGGCCAC 19  
 20  
 TCTATTGGGC CTGTCAAGGC TGTGTCCTAA CTGGTATCTG GGTCATAGCC CACGAATGCG 19  
 80  
 GTCACCACGC ATTCAGCGAC TACCAATGGC TGGATGACAC AGTTGGTCTT ATCTTCCATT 20  
 40  
 CCTTCCTCCT CGTCCCTTAC TTCTCCTGGA AGTATAGTCA TCGCCGTCAC CATTCCAACA 21  
 00  
 CTGGATCCCT CGAAAGAGAT GAAGTATTTG TCCCAAAGCA GAAATCAGCA ATCAAGTGGT 21  
 60  
 ACGGGAAATA CCTCAACAAC CCTCTTGGAC GCATCATGAT GTTAACCGTC CAGTTTGTCC 22  
 20  
 TCGGGTGGCC CTTGTACTTA GCCTTTAACG TCTCTGGCAG ACCGTATGAC GGGTTCGCTT 22  
 80  
 GCCATTTCTT CCCCAACGCT CCCATCTACA ATGACCGAGA ACGCCTCCAG ATATACCTCT 23  
 40  
 CTGATGCGGG TATTCTAGCC GTCTGTTTTG GTCTTTACCG TTACGCTGCT GCACAAGGGA 24  
 00  
 TGGCCTCGAT GATCTGCCTC TACGGAGTAC CGCTTCTGAT AGTGAATGCG TTCCTCGTCT 24  
 60

TGATCACTTA CTTGCAGCAC ACTCATCCCT CGTTGCCTCA CTACGATTCA TCAGAGTGGG 25  
 20  
 ACTGGCTCAG GGGAGCTTTG GCTACCGTAG ACAGAGACTA CGGAATCTTG AACAAAGGTGT 25  
 80  
 TCCACAACAT TACAGACACA CACGTGGCTC ATCACCTGTT CTCGACAATG CCGCATTATA 26  
 40  
 ACGCAATGGA AGCTACAAAG GCGATAAAGC CAATTCTGGG AGACTATTAC CAGTTCGATG 27  
 00  
 GAACACCGTG GTATGTGGCG ATGTATAGGG AGGCAAAGGA GTGTATCTAT GTAGAACCGG 27  
 60  
 ACAGGGAAGG TGACAAGAAA GGTGTGTACT GGTACAACAA TAAGTTATGA GGATGATGGT 28  
 20  
 GAAGAAATTG TCGACTTTTC TCTTGTCTGT TTGTCTTTTG TTAAAGAAGC TATGCTTCGT 28  
 80  
 TTTAATAATC TTATTGTCCA TTTTGTGTGT TTATGACATT TTGGCTGCTC ATTATGTTAT 29  
 40  
 GTGGGAAGTT AGCGTTCAAA TGTTTTGGGT CGG 29  
 73

## (2) INFORMATION FOR SEQ ID NO:16:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 23 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

## (ix) FEATURE:

- (A) NAME/KEY: misc feature
- (B) LOCATION: 1..23
- (D) OTHER INFORMATION: /product="synthetic oligonucleotide"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

GGGCATGTNG ARAANARRTG RTG 23

## (2) INFORMATION FOR SEQ ID NO:17:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 23 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(ix) FEATURE:

(A) NAME/KEY: misc\_feature

(B) LOCATION: 1..23

(D) OTHER INFORMATION: /product=  
 "synthetic  
 oligonucleotide"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

GGGCATGTRC TRAANARRTG RTG

23

EXPRESS MAIL LABEL NO. EL073740691US

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

IN THE APPLICATION OF:

E. I. DUPONT DE NEMOURS AND COMPANY

CASE NO.: BB1043 US NA DIV

APPLICATION NO.: UNKNOWN

GROUP ART UNIT: UNKNOWN

FILED: CONCURRENTLY HERewith

EXAMINER: UNKNOWN

FOR: GENES FOR MICROSOMAL DELTA-12  
FATTY ACID DESATURASES AND  
RELATED ENZYMES FROM PLANTS


Assistant Commissioner for Patents  
Washington, DC 20231

Sir:

**DECLARATION IN ACCORDANCE WITH 37 CFR 1.821**

I hereby state that the content of the paper and computer readable copies of the Sequence Listing, submitted in accordance with 37 CFR 1.821(c) and (e), respectively are the same.

Respectfully submitted,



LYNNE M. CHRISTENBURY  
ATTORNEY FOR APPLICANTS  
REGISTRATION NO. 30,971  
TELEPHONE: 302-992-5481  
FACSIMILE: 302-892-1026

Dated: October 26, 2000

009207 " 54E 960

# DECLARATION and POWER OF ATTORNEY

As a below-named inventor, I declare that I believe that I am the original, first and sole inventor (if only one name is listed below) or an original first and joint inventor (if plural names are listed below) of the subject matter which is claimed in the specification identified below and for which a patent is sought on the invention as titled therein. I hereby state that I have reviewed and understand the contents of said specification including the claims, as amended by any amendment referred to herein. I acknowledge the duty to disclose to the Office all information known to me to be material to patentability as defined in 37 CFR §1.56.

Full Name of Inventor <sup>(1)</sup>	Last Name <b>LIGHTNER</b>	First Name <b>JONATHAN</b>	Middle Name <b>EDWARD</b>
Residence & Citizenship	City <b>MARIETTA</b>	State or Foreign Country <b>PENNSYLVANIA</b>	Country of Citizenship <b>U.S.A.</b>
Post Office Address	Post Office Address <b>R.D.2 BOX 101A</b>	City <b>AIRVILLE</b>	State or Country <b>PENNSYLVANIA</b> Zip Code <b>17302</b>
Full Name of Inventor <sup>(2)</sup>	Last Name <b>OKULEY</b>	First Name <b>JOHN</b>	Middle Name <b>JOSEPH</b>
Residence & Citizenship	City <b>COLUMBUS</b>	State or Foreign Country <b>OHIO</b>	Country of Citizenship <b>U.S.A.</b>
Post Office Address	Post Office Address <b>217 FALLIS ROAD</b>	City <b>COLUMBUS</b>	State or Country <b>OHIO</b> Zip Code <b>43214</b>
Full Name of Inventor <sup>(3)</sup>	Last Name	First Name	Middle Name
Residence & Citizenship	City	State or Foreign Country	Country of Citizenship
Post Office Address	Post Office Address	City	State or Country Zip Code

Title of the Invention: **Genes for Microsomal Delta-12 Fatty Acid Desaturases and Related Enzymes from Plants**

- ☒ Patent specification attached.
- ☐ Patent specification was filed on \_\_\_\_\_ as U.S. application Serial No. \_\_\_\_\_ and was amended on \_\_\_\_\_ (if applicable).
- ☐ No earlier-filed foreign applications of which priority benefit is claimed.
- ☐ Required information as to earlier-filed foreign applications of which priority benefit is claimed appears on page 2 attached.
- ☐ No earlier-filed U.S. applications of which priority benefit is claimed.
- ☒ Required information as to earlier-filed U.S. applications of which priority benefit is claimed appears on page 2 attached.

POWER OF ATTORNEY: The power to prosecute this application and transact all business in the Patent and Trademark Office connected therewith is hereby granted to the following ☒ attorney(s) ☐ agent(s):

Name: **Barbara C. Siegell**  
**Hoge T. Sutherland**

Registration No.: **30,684**  
**20,010**

Send correspondence and direct telephone calls to:  
**Barbara C. Siegell**

**E. I. du Pont de Nemours & Co.**  
**Legal - Patents**  
**Wilmington, DE 19898**

Tel. No.  
**(302) 992-4931**

I further declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

Signature of Inventor<sup>(1)</sup>

Signature of Inventor<sup>(2)</sup>

Signature of Inventor<sup>(3)</sup>

Date **20 JUNE 1994**

Date **Sept 21, 1994**

Date

☐ Required information as to earlier-filed foreign applications of which priority benefit is claimed.

I hereby claim foreign priority benefits under Title 35, United States Code, §119 of any foreign application(s) for patent or inventor's certificate listed below and have also identified below any foreign application for patent or inventor's certificate having a filing date before that of the application on which priority is claimed.

Country	Application No.	Filing Date	Priority Claimed (Yes-No)
---------	-----------------	-------------	---------------------------

☒ Required information as to earlier-filed U.S. applications of which priority benefit is claimed.

I hereby claim the benefit under Title 35, United States Code, §120 of any United States application(s) listed below and insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application in the manner provided by the first paragraph of Title 35, United States Code, §112. I acknowledge the duty to disclose to the Office all information known to me to be material to patentability as defined in 37 CFR §1.56, which became available between the filing date of the prior application and the national or PCT international filing date of this application.

U.S. Application Serial No.	U.S. Filing Date	Status (patented, pending or abandoned)
07/977,339	11/17/92	Abandoned
PCT/US93/09987	10/15/93	Pending
U.S. National Application filed 6/17/94		